

GORDON'S CHEMICALS SEVIN® 50W SPRAY

A WETTABLE POWDER

This 2 Pound Bag Treats 5,000 Square Feet of Lawn

CONTROLS: SOD WEBWORM - ANTS - FLEAS - LEAFHOPPERS
VEGETABLE, FRUIT AND ORNAMENTAL PESTS


ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate).....	50.0%
INERT INGREDIENTS.....	<u>50.0%</u>
TOTAL	100.0%

SEVIN® is the registered trademark of Rhone Poulenc for carbaryl insecticide.

KEEP OUT OF REACH OF CHILDREN


CAUTION



See side panel for Precautionary Statements
and Statement of Practical Treatment

NET WEIGHT 2 POUNDS

620 APXXXXXX
EPA REG. NO. 2217-389
EPA EST. NO. 2217-KS-1
Mfd. by PBI/GORDON CORPORATION
KANSAS CITY, MISSOURI 64101



274

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid breathing vapors [dust or spray mist]. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

User Safety Requirements:

- ◆ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
- ◆ Wear a hat and eye protection when making overhead applications.
- ◆ Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- ◆ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- ◆ Do not allow children or pets to contact treated area until sprays have dried.

Statement of Practical Treatment

IF SWALLOWED: Induce vomiting and seek medical attention immediately.

IF IN EYES OR ON SKIN: Flush eyes thoroughly with water; wash skin thoroughly with soap and water.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine sulfate is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS:

This pesticide is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold and discard bag in trash.

- ♦ Apply when air is calm to avoid drift and contact with eyes and skin.
- ♦ Start applying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.
- ♦ For trees taller than 10 feet, consider hiring a licensed professional.
- ♦ Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.
- ♦ Do not repeat applications more than once a week.
- ♦ Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

- SOD WEBWORM CONTROL -

When to spray: In late June and July where numerous moths can be seen flying low over bluegrass lawn during early evening, spraying anytime within the next week will prevent damage from hatching worms. If lawn damage was heavy the preceding year, an early spray (May 15th) will prevent spring damage from over-wintering worms.

How to spray: To get spray down to soil level, mow lawn and rake out any heavy grass. Water lawn. Spray using a garden-pressure hand operated sprayer or hose-end sprayer. Add Sevin® 50W Spray to water using at least 1 pint of water for every 2 cups insecticide. Thoroughly mix, then fill spray container with water to spray level.

Use 2 cupfuls of product in 10 to 15 gallons of water per 1000 square feet. Sprinkle after spraying to wash insecticide into soil level of turf. This 2 pound package will treat 5000 square feet, one time.

This treatment will control ants, fleas, leafhoppers, and chinch bugs.

- ORNAMENTAL INSECTS -

Use 2 pounds of product per 100 gallons of water or 2 tablespoons of product per gallon of water. Spray all foliage thoroughly. Repeat as necessary To control bagworm, blister beetle, boxelder bug, flea beetle, lace bug, leafhoppers, rose slug, and green cankerworm. Do not spray Boston Ivy or Virginia Creeper. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

- HOME ORCHARDS AND GARDEN VEGETABLES -

For best insect control through season plan on a regular spray schedule every 7 to 10 days. Two (2) pounds of product to 100 gallons of water is equivalent to 2 tablespoons of product per gallon of water.

APPLES AND PEARS: *Codling moth, plum curculio, apple aphid, green fruitworm, Japanese beetle, leaf roller, and San Jose scale* - As insects appear, spray, using mix 2 tablespoons of product per gallon of water. repeat as needed for control, up to one day before harvest. To avoid fruit thinning, delay use until at least 30 days after bloom (not before second cover). For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week. Allow one (1) day before harvest.

PEACHES, APRICOTS, PLUMS, AND CHERRIES: *Plum curculio, oriental fruit moth, lesser peach tree borer, peach twig borer, "cat-facing" bugs, and San Jose scale* - Mix 2 tablespoons of product per gallon of water. Allow one day between spraying and harvest; three days cr. Apricots. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

4 7 4

CUCUMBER, MELONS, AND SQUASH: *For pickleworm, melonworm, cucumber beetle, flea beetle, leafhoppers, and squash bug* - Mix 2 tablespoons of product per gallon of water. Avoid excessive applications since leaf injury may occur on tender foliage when wet and humidity is high. Allow 3 days before harvest. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

TOMATOES, EGGPLANTS, PEPPERS: *For tomato hornworm, and fruitworm, fall armyworm, stink bugs, and lace bug* - Mix 2 tablespoons of product per gallon of water. Thoroughly treat. Use until day of picking. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

CABBAGE, BROCCOLI, AND CAULIFLOWER: *For control of cabbage looper, flea beetle, armyworm, and corn earworm* - Mix 2 tablespoons of product per gallon of water on a 7 to 10 day spray schedule, allowing 3 days before picking. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

SWEET CORN: *Corn earworm, European corn borer, flea beetle, and grasshoppers* - Mix 2 tablespoons of product per gallon of water. Spray for earworms until silks are dry. Apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals. ~~as infestation warrants. Reapply immediately after rain.~~ For other pests, thoroughly treat entire plant every 2 weeks, starting when corn is knee high. Do not apply within 7 days of harvest if fodder is to be fed to livestock, otherwise no limitation.

Note: Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of products or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.