


GORDON'S



tomato
AND
garden dust

ACCEPTED

MAR 27 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
recommended, for the pesticide
registered under
EPA Reg. No.

DIRECTIONS FOR USE

DUST. Apply thoroughly to vegetable and flowering plants, making sure to treat both top and bottom sides of leaves, stems and fruit. Make first application when insects first appear and / or disease is anticipated. Begin early, when new growth is starting. Repeat as necessary generally at 7 day intervals.

BEANS: (Snap and Lima)—Controls aphids, red spiders, leafhoppers, Mexican bean beetles, bean leaf beetles and thrips. Diseases: downy mildew, bean rust, and anthracnose. Do not apply within 7 days of harvest.

CABBAGE, CAULIFLOWER, BROCCOLI, TURNIPS, AND CARROTS
—Controls aphids, imported cabbage worm, cabbage looper, flea beetles. Diseases: downy mildew and alternaria leaf spot. Stop use 14 days before harvest on Broccoli, 7 days on Cabbage and Cauliflower, 7 days on Turnips and Carrots, or 14 days if tops are to be used for feed or food.


CUCUMBERS AND SQUASH—Controls melon worms, striped a., cucumber beetles, flea beetles and aphids. Diseases: downy mildew, alternaria (Macrosporium blight), anthracnose and gummy stem blight. Stop use 7 days before harvest.

TOMATOES AND EGGPLANT—Controls aphids, tomato russet mite, spider mites, tomato horn worm, tomato fruit worm, flea beetles, leafhoppers and psyllid. Diseases include early and late blight, anthracnose, stemphylium and septoria leaf spots and cladosporium grey mold. Do Not Apply within 5 days of harvest.

FLOWERS AND ORNAMENTALS (On roses, mums, snapdragons, glads, hollyhocks)—Controls botrytis blight and stemphylium on glads, rust, septoria leaf spot and anthracnose on snaps, mums, carnations and hollyhocks. Black spot on roses. Kills aphids, thrips, grasshoppers and cutworms. For cutworms, dust the soil around the base of each plant.

SWEET CORN—For corn earworm, European corn borer, fall army worm and flea beetles. For earworm, dust silks at 2 to 3 day intervals until they are dry. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations. See Note.

Squeeze Buster



CAUTION: Keep Out of Reach of Children

See side panel for additional cautions

NET WEIGHT TEN OUNCES

Manufactured by

PDI GORDON CORPORATION

KANSAS CITY, KANSAS 64110

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

"Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated."

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE: Do not feed forage, fruit or pods to dairy animals or animals being finished for slaughter. Toxic to honey bees. Do not apply to open blossoms.

Refill only with Gordon's TOMATO & GARDEN DUST. Destroy package when unit becomes unserviceable.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

ACTIVE INGREDIENTS

1-Garbaryl (1-Naphthyl N-Methylcarbamate)	3.00%
2-Zeb. 2-(o-Ethylene Res.)thio Carbamate	3.90%
3-Prop. Piperonyl Butoxide**	0.50%
4-Pyrene	0.05%
INERT INGREDIENTS	92.55%

TOTAL 100.00%

*Sevin T.M. Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate, U.S. Patent 2,903,478

**Equivalent to (Butylcarbityl) (6-propylpiperonyl) ether 0.4% plus related compounds 0.1%

P.A. Reg. No. 2217-401 AA

T.M. Registered 2-8

528-674

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

"Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated."

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE: Do not feed foliage, fruit or pods to dairy animals or animals being finished for slaughter. Toxic to honey bees. Do not apply to open blossoms.

Refill only with Gordon's TOMATO & GARDEN DUST. Destroy package when unit becomes unserviceable.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

ACTIVE INGREDIENTS:

*Carbaryl (1-Naphthyl N-Methylcarbamate)	3.00 %
*Zineb (Zinc Ethylene Bisdithio Carbonate)	3.90 %
*Tech. Piperonyl Butoxide**	0.50 %
*Pyrethrins	0.05 %

INERT INGREDIENTS:

92.55 %

TOTAL 100.00%

*Sevin T.M. Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate, U.S. Patent 2,903,478

**Equivalent to (Butylcarbityl) (6-propylpiperonyl) ether 0.4% plus related compounds 0.1%

E.P.A. Reg. No. 2217-401-AA

*T.M. Registered, P.B.I.

E.P.A. EST. NO. 2217-KSH

528-674