

ACCEPTED

MAR 9 1984

the Federal Insecticide,
Fungicide, and Rodenticide Act.
No. 7217-401

TOMATO & GARDEN DUST

Contains SEVIN* and ZINEB
Controls insects and plant diseases.

ACTIVE INGREDIENTS:

Zineb (zinc ethylene bisdithiocarbamate).....	3.90%
*Carbaryl (1-naphthyl N-methylcarbamate).....	3.00%
**Technical Piperoryl Butoxide.....	0.50%
Pyrethrins.....	0.05%
<u>INERT INGREDIENTS.....</u>	<u>92.55%</u>
	TOTAL 100.00%

*SEVIN is the registered trademark of
UNION CARBIDE CORPORATION for carbaryl
insecticide.

**Equivalent to (Butylcarbityl) (6-propylpiperonyl)
ether 0.4% plus related compounds 0.1%.

KEEP OUT OF REACH OF CHILDREN ←
CAUTION ←
See side panels for additional precautionary statements.

12 pt bold
18 pt bold

NET WEIGHT 10 OUNCES

STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW
DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

-Hazards to Humans & Domestic Animals-

CAUTION: May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. When using, wear protective clothing, i.e. long sleeves, long pants, hat, gloves, covering for face and eyes. Wash thoroughly after use and launder clothing before reuse. Remove any residues at harvest by brushing, washing or peeling. Do not feed treated crops, pods, or husks to dairy animals or animals being readied for slaughter.

Note to Physician: This product contains carbaryl. Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

-Environmental Hazards-

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Bee Precaution: This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not dust when bees are active in the area or plants in full bloom. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

12 pt. bold

STORAGE & DISPOSAL

Store away from children, pets, and foodstuffs. This squeeze duster is an applicator/container package. The container may be refilled and used again and again, but only with this product. WHEN NO LONGER WANTED OR NO LONGER SERVICEABLE, WRAP AND PLACE IN TRASH COLLECTION.

528/283

EPA REG. NO. 2217-401
EPA EST. NO. 2217-PP-1

Mfd. by PBI/GORDON CORPORATION
Kansas City, Mo. 64112

Limited Warranty. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suitable for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

Exclusion of Other Warranties and Remedies. Except where such disclaimers and exclusions are specifically prohibited by applicable law, THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESS OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty, BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

HOW TO USE

Lift cap and squeeze sidewalls to dust. Apply thoroughly to vegetable and flowering plants, making sure to treat both top and bottom sides of leaves, stems, and fruit. Make first application when insects first appear and/or disease is anticipated. Begin early, when new growth is starting. Repeat as necessary, generally at 7 day intervals.

BEST DOCUMENT AVAILABLE

USES

BEANS. Snap and Lima. Controls aphids, red spiders, leafhoppers, Mexican bean beetles, bean leaf beetles and thrips. Diseases: Downy mildew, bean rust and anthracnose. Do not apply within 7 days of harvest.

CABBAGE, CAULIFLOWER, BROCCOLI, TURNIPS, AND CARROTS. Controls imported cabbage worm, cabbage looper, flea beetle. Diseases: downy mildew and alternaria leaf spot. Stop use 14 days before harvest on Broccoli, 7 days on Cabbage and Cauliflower, 7 days on Turnips, and Carrots, or 14 days if tops are to be eaten for feed or food.

MELONS, CUCUMBERS AND SQUASH. Controls cucumber beetle, striped and spotted cucumber beetles, flea beetle, and aphids. Diseases: downy mildew, Alternaria (*Alternosporium*) blight, anthracnose, stem blight. Stop use 7 days before harvest.

TOMATOES AND EGGPLANT. Controls spider mites, tomato horn worm, tomato pinworm, tobacco hornworm, caterpillars and psyllid. Diseases: net leaf, late blight, septoria, stemmy blight and septoria leaf spots and other diseases. Stop use within 5 days of harvest.

SWEET CORN. Controls European cornborer, corn earworm, and other caterpillars. Apply within 7 days of silking. Do not apply within 7 days of harvest. Do not apply during the period of silking.

FLOWERS AND ORNAMENTALS. (On ornamentals, use only on flowers.) Controls leafhoppers, beet beetles, and aphids. Diseases: downy mildew and anthracnose. Do not use on snapdragons, petunias, and geraniums. Do not use on flowers to be eaten for feed or food.

(Note to reviewer: underlines, items will be set in bold-faced type to highlight.)

Do not feed treated crop plants, or bulk to feed animals or animals being raised for slaughter.