


(Front Panel)

GORDON'S

LV 600 2,4-D WEED KILLER

A LOW VOLATILE ESTER
FOR AGRICULTURAL & SERVICE USE ONLY.



ACTIVE INGREDIENT:

*Isooctyl ester of 2,4-Dichlorophenoxyacetic acid.....87.9%

INERT INGREDIENTS.....12.1%

TOTAL 100.00%

This Product Contains:

*5.6 lbs. 2,4-Dichlorophenoxyacetic acid per gallon or 58.3%

*isomer Specific By AOAC Method No. 6.D01-5.

KEEP OUT OF REACH OF CHILDREN ← 14 pt bold

CAUTION ← 18 pt bold

Statement of Practical Treatment

If Swallowed: Do not induce vomiting. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If On Skin: Wash with soap and water. Get medical attention if irritation persists.

If In Eyes: Flush with water 15 minutes and get medical attention.

See side panels for additional precautionary statements.

NET CONTENTS ONE U.S. GALLON



READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Harmful if inhaled. Avoid breathing spray mist. Remove contaminated clothing and wash before reuse.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Use with care when applying in areas adjacent to any body of water. Do not contaminate water intended for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes.

Physical Or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL ← 14 qt. bold

This product may be stored in an unheated building.

Prohibitions: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near pesticides or seeds.


Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

Container Disposal: Triple rinse (or equivalent) and offer for recycling, or dispose of in a sanitary landfill, or by other approved State and local procedures.

USE PRECAUTIONS:

Don't overdose. Avoid spray drift to cotton, soybeans, tomatoes, tobacco, grapes, fruit trees, flowers, garden crops, ornamental plants, shrubs, trees and other hormone herbicide--sensitive desirable plants. Do not apply near these plants because small quantities of wind-drifted herbicide may cause severe injury. Do not apply when wind speed is sufficient to cause drift. Do not apply when a temperature air inversion exists. An air inversion may be detected by creating a smoke column and observing for a layering effect. Do not apply when temperature exceeds 90° F. Do not apply if rain is expected within the hour.

817/480
EPA REG. NO. 2217-413
EPA EST. NO. 2217-KS-1



KANSAS CITY, KANSAS 64114

GENERAL:

Apply LV 600 as a water or oil spray during warm weather when weeds or brush are actively growing. Application under drought conditions often will give poor results. Use low spray pressure to minimize drift. On cropland and along roadsides, do not exceed 20 psi pressure. Apply enough spray volume to provide uniform coverage of weeds and brush, usually 5 to 20 gallons per acre by ground equipment and 3 to 5 gallons by aircraft. Higher gallonage may be used if desired to improve spray coverage. Generally, the low dosages recommended on this label will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species, and, under conditions where control is more difficult, the higher dosages will be needed. For crop uses, do not mix with oil or other adjuvants unless specifically recommended on this label. Deep-rooted perennial weeds such as Canada thistle and field bindweed, and many woody plants usually require repeated applications for maximum control. Do not apply LV 600 where spray drift may contact nearby susceptible crops or other desirable plants, or may contaminate water used for irrigation or domestic purposes. Read and follow all precautions on this label. Local conditions may affect the use of herbicides. Consult your State Agricultural Experiment Station or Extension Service weed specialists for advice in selecting treatments from this label to best fit local conditions. Be sure that use of this product conforms to all applicable regulations. Apply this product only as specified on this label.

WEEDS:

ARROWHEAD	JEWELWEED	SHEPHERDSPURSE
BEGGARTICKS	JIMSONWEED	SICKLEPOD
BINDWEED	KNOTWEED	SMARTWEED
BITTERWEED	KOCHIA	SNEEZEWEED, BITTER
BLUEWEED TEXAS	LAMBSQUARTER	SOWTHISTLE, ANNUAL
BROOMWEED	LOCOWEED	SPANISHNEEDLES
BUCKBRUSH	MALLOW, VENICE	SUMAC
BUCKHORN	MANZANITA	SUNFLOWER
BUCKWHEAT	MARSHELDER	SWEETCLOVER
BURDOCK	MILKWEED	TANSYMUSTARD
BURHEAD	MILKVETCH	TANSYRAGWORT
CANADA THISTLE	MORNINGGLORY, ANNUAL	THISTLE, BULL
CARPETWEED	MUSTARDS	THISTLE, MUSK
CATNIP	NETTLES	THISTLE, RUSSIAN
CHAMISE	PENNYCRESS	TUMBLEWEED
CHICORY	PEPPERWEED, FIELD	VELVETLEAF
COCKLEBUR	PIGWEEED	VERVAINS
COFFEEWEED	PLANTAINS	VETCH
CORNFLOWER	POISON IVY	WATER PLANTAIN
COYOTEBRUSH	POORJOE	WILD CARROT
CROTON	RABBITBRUSH	WILD GARLIC
DANDELION	RAGWEED	WILD HEMP
DOCKS	RAPE, WILD	WILD ONION
DOGFENNEL	REDSTEM	WILD RADISH
ELDERBERRY	SAGE, COASTAL	WILD SWEET POTATO
FANWEED	SAGEBRUSH, BIG	WILLOW
GALINSOGA	SAGEBRUSH, SAND	WITCHWEED
GOATSBEARD	SALSIFY	WORMWOOD
HALOGETON	Sand	YELLOW ROCKET
HOARY CRESS	Shinnery Oak	YELLOW STARHISTLE
HORSE NETTLE	SHEEP SORREL	

NOTES ABOUT WIND DRIFT:

Ground Equipment--Spray drift can be lessened by: Keep the spray boom as low as possible and by applying 20 gallons or more of spray per acre. Use no more than 20 pounds spraying pressure with flat fan or flooding flat fan nozzle tips. Spray when wind velocity is low. Do not spray with oil when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone-type insecticide or other nozzles that produce a fine-droplet spray.

Aircraft Application--Spray drift can be lessened by: Apply not less than 5 gallons of spray per acre. Use no more than 20 pounds spray pressure at the nozzles. Use nozzles which produce a coarse spray pattern. Spray only when wind velocity is less than 5 miles per hour.

PREPARATION OF THE SPRAY:

With Water--Fill the spray tank about half full with water. Add the required amount of LV 600 with agitation. Then, add the rest of the water.

Note: LV 600 in water forms an emulsion which tends to separate unless the mixture is kept agitated.

With Water & Oil--Mix LV 600 and the oil first. Add this mixture to the water. However, with adequate agitation, the oil can be added after the LV 600 is mixed in the water.

With Oil--If straight oil is used, a solution is formed and separation does not occur. Do not allow any water to get into the herbicide-oil solution to avoid formation of an invert emulsion.

SMALL GRAINS (not underseeded with a legume):

Note: Do not permit dairy or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment. Do not feed treated straw to livestock.

Spring Wheat & Barley--Apply 1/3 to 3/4 pint per acre. Spray when grain is in full tiller stage (usually 4 to 8 inches tall) but before the boot stage, and when weeds are small. Do not apply before the tiller stage nor from early boot to the dough stage. Higher rates (3/4 to 1 1/2 pints per acre) may be required to control certain weeds but crop injury may occur.

Winter Wheat & Rye--Apply 1/3 to 1/2 pint per acre in the spring at the full tiller stage but before the early boot stage.

Spring Seeded Oats--Apply 1/3 pint per acre at the full tiller stage but before the early boot stage. Oats are less tolerant to 2,4-D than wheat or barley and more likely to suffer some injury.

Fall Seeded Oats (Southern) Grown for Grain--Apply 1/2 to 1 pints per acre after full tillering but before the early boot stage. Some difficult weeds may require the higher rate for maximum control but crop injury may result. Do not apply during or immediately following cold weather.

Preharvest Treatment--Apply 1/2 to 1 1/2 pints per acre when grains are in the hard dough stage to control large weeds that may interfere with harvest. Best results are obtained when soil moisture is sufficient to cause succulent weed growth.

CORN: Hybrids vary in response to 2,4-D and some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Contact seed company or your Agricultural Experiment Station or Extension Service weed specialists for this information.

Use one of the following programs for weed control in corn:

Preemergence--Apply 1 1/2 to 2 2/3 pints per acre to soil anytime after planting but before corn emerges. Do not use on light sandy soil. Do not cultivate until necessary.

Emergence--Apply 3/4 pint per acre just as corn plants are breaking ground.

Postemergence--After emergence of corn, use 1/2 pint per acre. Application of 1/2 to 3/4 pint per acre may be needed for maximum control of some weeds but such rates are more likely to injure the corn. Do not apply from the tasseling to dough stage. Do not use with oil, Atrazine or other adjuvants. Crop injury is more likely to occur if corn is growing rapidly under high temperature and high soil moisture conditions. To reduce breakage of stalks from temporary brittleness caused by 2,4-D, delay cultivation for 8 to 10 days after treatment.

Pre-Harvest Treatment--After the hard dough or denting stage, apply 1/3 to 1 1/3 pints per acre by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as bindweed, cocklebur, dogbane, Jimsonweed, ragweed, smartweed, velvetleaf, and vines that interfere with harvesting. Do not forage or feed corn fodder for 7 days following application.

SORGHUM (Milo): Apply 1/3 pint per acre when sorghum is 5 to 15 inches tall. A higher rate of 1/2 to 3/4 pint per acre may be needed to control some weeds but the chance for crop injury is likewise increased. Do not use with oil. Do not treat before the sorghum is 5 inches tall nor during the boot, flowering, or early dough stages. If sorghum is taller than 8 inches, use drop nozzles to keep the spray off the foliage as much as possible. Temporary crop injury may occur under conditions of high soil moisture and high air temperatures. Varieties vary in tolerance to 2,4-D and some hybrids are quite sensitive. Spray only varieties known to be tolerant to 2,4-D. Some hybrids are quite sensitive. Contact seed company or your Agricultural Experiment Station or Extension Service weed specialists for information.

GRASS SEED CROPS: Use 3/4 to 1 pints per acre in the amount of water required for uniform application by air or ground equipment. Apply to established stands in spring from the tiller to early boot stage. Do not spray in boot stage. New spring seedlings may be treated with the lower rate after the grasses have at least five leaves. Perennial weed regrowth may be treated in the fall.

WEED AND BRUSH CONTROL IN RANGELANDS & GRASS PASTURES: Do not graze dairy animals on treated areas within 7 days after application. Do not use on bentgrass, alfalfa, clover, or other legumes. Do not use on newly seeded areas until grass is well established. Do not use from early boot to milk stage where grass seed production is desired.

Bitterweed, Broomweed, Croton, Decks, Kochia, Marshelder, Musk-thistle and Other Broadleaf Weeds--Use 3 pints of LV 600 per acre in the amount of water needed for uniform application. If the weeds are young and growing actively, 1/2 pints per acre will provide control of some species. Deep-rooted perennial weeds may require repeated treatments in the same year or in subsequent years.

Wild Garlic and Wild Onion--Apply 3 to 4 pints per acre making three applications (fall-spring-fall) or (spring-fall-spring), starting in late fall or early spring.

Weed Control in Newly Sprigged Coastal Bermudagrass--Apply 1/2 to 3 pints per acre preemergence and/or postemergence.

Sand Shinnery Oak and Sand Sycamore--Use 1/2 pint in 5 gallons of oil or 1/4 gallon in 5 gallons of water or other

acre. Apply by aircraft between May 15 and June 15. On the Sagebrush, use 1½ pints in 3 gallons of oil per acre and apply by aircraft when foliage is fully expanded and the brush is actively growing.

Big Sagebrush and Rabbitbrush--Use 3 to 4 pints per acre in 2 to 3 gallons of oil or in 3 to 5 gallons of oil-water emulsion spray. For rabbitbrush, the 3 quart rate is usually required. Brush should be leafed out and growing actively when treated. Retreatment may be needed.

Chamise, Manzanita, Buckbrush, Coastal Sage, Coyotebrush, and Certain Other Chaparral Species--Use 3 to 4 pints per acre in 5 to 10 gallons of water. One gallon of fuel oil may be included in the spray mixture for added effectiveness. Make applications by aircraft or ground equipment to obtain uniform spray coverage. For effective control the brush must be fully leafed out and growing actively when sprayed. Retreatment may be needed.

WOODY PLANT CONTROL IN NON-CROP AREAS: To control species susceptible to 2,4-D in right-of-ways, fencerows, roadsides, and along drainage ditchbanks, spray brush up to 5 to 8 feet tall after spring foliage is well developed, using 4 to 6 pints of LV 600 in 100 gallons of water and wetting all parts of the brush including foliage, stems and bark. This may require up to 400 gallons of spray per acre for adequate coverage of solid stand of brush. Make applicaiton in such a way as to prevent drift of the spray away from the area being treated. Spraying can be effective at any time up to 3 weeks before frost as long as soil moisture is sufficient for active growth of the brush. Control will be less effective in mid-summer during hot dry weather when soil moisture is deficient and plants are not actively growing. Oil or wetting agent may be added to the spray, if needed, for increased effectiveness.

WEED CONTROL IN NON-CROP AREAS SUCH AS LAWNS, GOLF COURSES, CEMETERIES, PARKS, AIRFIELDS, ROADSIDES, VACANT LOTS, DRAINAGE DITCH BANKS: Apply 1½ to 4 pints per acre in the amount of water needed for uniform application. Usually 3 pints per acre provides good weed control under average conditions. Treat when weeds are young and growing well. Do not use on golf greens nor on dichondra or other broadleaf herbaceous ground covers. Do not use on creeping grasses such as bent and St. Augustine except for spot treating, not on newly seeded turf until grass is well established.

Reseeding of treated areas should be delayed following treatment. With spring application, reseed in the fall; with fall application, reseed in the spring. Legumes are usually damaged or killed so do not treat areas where the legumes are desired. Deep-rooted perennial weeds may require repeated treatment in the same season or in subsequent years.

TULE (BULRUSH) AND OTHER RUSSIA: Mix 1 pint of LV 600 and 1 gallon of diesel oil or kerosene, then add this mixture to 100 gallons of water. Spray to wet all foliage (400-800 gallon per acre).

Addition of a wetting agent may be advisable. Apply in the spring during flower head emergence. Respray if needed when regrowth is 3 to 5 feet tall.

SPOT TREATMENT: To control broadleaf weeds in small non-cropland areas with a hand sprayer, use $\frac{1}{2}$ pint of LV 600 in 3 gallons of water and spray to thoroughly wet all weed foliage. Keep spray mixture agitated to prevent separation.

WARRANTY:

The manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled uses when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed. Therefore, the manufacturer makes no other implied or express warranty nor is any agent of the manufacturer allowed to do so. Upon purchase of this product the buyer assumes all risks associated with use of this product. In the event of damage resulting from a breach of warranty the buyer agrees to accept a refund of the purchase price of the product as full discharge of the manufacturer's liability.