

\$2.98

Feb. 12, 1974

# fore<sup>®</sup> lawn and ornamental fungicide

### ACTIVE INGREDIENTS

A co-ordination product of zinc ion and manganese ethylene bisdithiocarbamate..... 80%

In these ingredients are

Manganese + + ..... 18%

Zinc + ..... 2%

Ethylene bisdithiocarbamate ion (C<sub>2</sub>H<sub>2</sub>N<sub>2</sub>S<sub>2</sub>) = ..... 62%

NET WEIGHTS ..... 20%

100%

U. S. Patents 2,504,404; 2,874,158 and Patent Pending  
 ® Registered Trademark Rehm and Maas Co.

NET WEIGHT 15 OUNCES

**CAUTION: Keep Out of Reach of Children**  
 See Side Panels for Additional Cautions

### PRODUCT USE

FORE Turf and Ornamental Fungicide is a unique and different fungicide which when used according to directions, provides very high fungicidal activity and is safe for use on turf grasses and many ornamentals. This new fungicide provides proven control of most common turf grass diseases and it is also effective in controlling major fungus diseases of ornamentals.

Diseases of turf grass and ornamentals can attack suddenly and unexpectedly causing severe damage and may even result in total loss of large areas of valuable turf grass and ornamental plants. The use of a regular protective spray program will minimize the risk of disease damage and can usually be accomplished with lower rates and less frequent fungicide applications. Once diseases have become established higher rates of fungicide and more frequent applications are required to bring them under control. Follow a regular protective spray program for maximum performance of your lawn and ornamentals.

### DIRECTIONS FOR USE

Apply FORE Fungicide to turf grass and ornamentals according to the directions on the label. Use a backpack sprayer or a pump-up sprayer. Apply the fungicide to the foliage of the plants. Do not apply to the soil. Do not apply to the roots of the plants. Do not apply to the stems of the plants. Do not apply to the flowers of the plants. Do not apply to the fruit of the plants. Do not apply to the seed of the plants. Do not apply to the young plants of the plants. Do not apply to the old plants of the plants. Do not apply to the dead plants of the plants. Do not apply to the plants that are already dead. Do not apply to the plants that are already dying. Do not apply to the plants that are already diseased. Do not apply to the plants that are already damaged. Do not apply to the plants that are already injured. Do not apply to the plants that are already stressed. Do not apply to the plants that are already weakened. Do not apply to the plants that are already exhausted. Do not apply to the plants that are already overgrown. Do not apply to the plants that are already undernourished. Do not apply to the plants that are already overwatered. Do not apply to the plants that are already underwatered. Do not apply to the plants that are already overfertilized. Do not apply to the plants that are already underfertilized. Do not apply to the plants that are already overpruned. Do not apply to the plants that are already underpruned. Do not apply to the plants that are already overwatered. Do not apply to the plants that are already underwatered. Do not apply to the plants that are already overfertilized. Do not apply to the plants that are already underfertilized. Do not apply to the plants that are already overpruned. Do not apply to the plants that are already underpruned.

Follow local turf disease control recommendations for details on timing of applications. Diseases on ornamentals generally occur under the same conditions favorable to turf diseases. Preventive sprays used during warm humid periods will reduce the risk of diseases. If preventive sprays are not used and disease does develop, three or more applications of FORE should be made at 3 to 5 day intervals to arrest the disease and protect new growth.

- FORE Fungicide can be used for the control of these and other diseases:
- Turf and Grass—Fusarium Blight; Red Thread; Copper Spot; Slime Molds; Helminthosporium; Rhizoctonia; Brown Patch; Rust; Pythium Blight; Algae; Dollar Spot and Fairy Ring.
  - Cornflower—Leaf Spot
  - Rose—Black Spot
  - Chrysanthemum—Botrytis Petal Spot
  - Gladiolus—Curvularia and Botrytis
  - Ornamentals—Leaf Spot
  - Ornamentals—Botrytis Blight
  - Ornamentals—Fusarium Blight
  - Ornamentals—Leaf Spot

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

**PRODUCT USE**

**FORE Turf and Ornamental Fungicide** is a unique and different fungicide which when used according to directions, provides very high fungicidal activity and is safe for use on turf grasses and many ornamentals. This new fungicide provides proven control of most common turf grass diseases and it is also effective in controlling major fungus diseases of ornamentals.

Diseases of turf grass and ornamentals can attack suddenly and unexpectedly causing severe damage and may even result in total loss of large areas of valuable turf grass and ornamental plants. The use of a regular protective spray program will minimize the risk of disease damage and can generally be accomplished with lower rates and less frequent fungicide applications. Once diseases have become established higher rates of fungicide and more frequent applications are required to bring them under control. Follow a regular protective spray program for maximum control.

**DIRECTIONS FOR USE**

Use FORE Fungicide on turf grasses and ornamentals according to the following directions. For detailed information on diseases of turf grasses and ornamentals, consult your local agricultural agent or extension office. For information on the use of FORE Fungicide on other plants, consult the label on the container. Use FORE Fungicide on turf grasses and ornamentals according to the following directions. For detailed information on diseases of turf grasses and ornamentals, consult your local agricultural agent or extension office. For information on the use of FORE Fungicide on other plants, consult the label on the container.

Follow local turf disease control recommendations for details on timing of applications. Diseases on ornamentals generally occur under the same conditions favorable to turf diseases. Preventive sprays used during warm humid periods will reduce the risk of diseases. If preventive sprays are not used and disease does develop, three or more applications of FORE should be made at 3 to 5-day intervals to arrest the disease and protect new growth.

- FORE Fungicide can be used for the control of these and other diseases:**
- Turf and Lawn—Fusarium Blight; Red Thread; Copper Spot; Slime Molds; Helminthosporium; Anthracnose; Brown Patch; Rust; Pythium Blight; Algae; Dollar Spot and Fairy Ring.
  - Cornflower—Leaf Spot
  - Rose—Black Spot
  - Shrub and Tree—Botrytis Blight
  - Chrysanthemum—Botrytis Petal Spot
  - Holly—Purple Spot
  - Gladiolus—Curvularia and Botrytis
  - Iris—Leaf Spot

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Some conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary. Seller makes no claims other than those stated on this label.

**DILUTION TABLES—TURF**

**Hand-End Sprayers:**

Use a lawn type sprayer. Measure required amount of FORE into jar and add just enough water to wet the powder. Stir until all powder is wet and smooth. Add water until concentrate in jar is up to the required level. Shake to agitate periodically while spraying.

AMOUNT FORE		Control Treatment		Fill Bottle to Mark Gallons	Area to Treat Sq. Ft.
Preventive Treatment Tbsp.	Cups	Tbsp.	Cups		
2	—	3	—	1	135
10	3/4	15	1	5	675
20	1 1/2	30	2	10	1,350
30	2	45	3	15	2,000

Use standard measuring spoons and cup. All volumes are level measure.

**Pressure Sprayers:**

AMOUNT FORE		Control Treatment		Fill Bottle to Mark Gallons	Area to Treat Sq. Feet
Preventive Treatment Tbsp.	Cup.	Tbsp.	Cup.		
1 1/2	—	2	—	1	Spray all foliage and flowers (except Gladiolus)* until spray begins to drip off leaves
3	—	4	—	2	
4 1/2	—	6	—	3	
6	—	8	—	4	
7 1/2	—	10	—	5	
9	—	12	—	6	

\*Reduce dosage by one half to spray gladiolus flower spikes.

**CAUTION:** May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. In case of skin contact wash with water; for eyes wash copiously with water and get medical attention. Fish may be killed if their waters are contaminated with this product. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not reuse container. Destroy when empty.

**IMPORTANT:** Keep away from fire and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of FORE and may also generate flammable vapors. Keep container closed when not in use.