

PM-15 2217-450  
1/24/0

194

VAPONA(R)

SHOW-COAT DAIRY CATTLE SPRAY

CONTAINS DDVP

Ready-To-Use for  
FACE FLIES -- STABLE FLIES -- HOUSE FLIES -- HORN FLIES

ACTIVE INGREDIENTS

Dichlorvos [2,2-dichlorovinyl dimethyl phosphate].....	1.0%
INERT INGREDIENTS.....	99.0%
TOTAL	100.0%

VAPONA(R) is a registered trademark of Shell Chemical Company for DDVP or Dichlorovos insecticide.

**KEEP OUT OF REACH OF CHILDREN**


**WARNING**

See side panels for Precautionary  
Statements and Statement of  
Practical Treatment

NET CONTENTS ONE U.S. GALLON (3.785 liters)

/190 AP092667  
EPA REG. NO. 2217-450  
EPA EST. NO. 2217-KS-1

Mfd by PBI/GORDON CORPORATION  
Kansas City, Kansas 66118



Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

STOP! READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.

2  
3  
4

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals**

**WARNING:** May be fatal if swallowed, inhaled, or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating or smoking.

**Statement of Practical Treatment**

**IF SWALLOWED:** Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

**IF INHALED:** Remove victim to fresh air. Apply artificial respiration if indicated.

**IF ON SKIN:** Remove contaminated clothing and wash affected areas with soap and water.

**IF IN EYES:** Flush eyes with plenty of water. Call a physician immediately.

**NOTE TO PHYSICIAN:** This product is an organic phosphate which is a cholinesterase inhibitor. Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally. If symptoms of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be used in conjunction with atropine.

**ENVIRONMENTAL HAZARDS:**

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**PHYSICAL OR CHEMICAL HAZARDS:**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### STORAGE & DISPOSAL

**STORAGE:** Do not contaminate water, food or feed by storage or disposal. Store in original container in a locked storage area inaccessible to children or pets.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**FACE FLIES, HORN FLIES, MOSQUITOES, HOUSE FLIES ON BEEF AND DAIRY CATTLE:** Use a mist type hand sprayer or automatic unit that will insure a fine spray mist for desired coverage of the animal. For Face Flies spray the forehead, taking care to avoid spraying the eyes. For Horn Flies, direct the spray mist to the back, withers, and belly of the cattle. Make 2 applications of one ounce each day. Treat in the morning and the evening.

**STABLE FLIES:** Direct the spray mist to the back, withers, belly, and leg areas of the cattle. Make 2 applications of one ounce each day. Treat in the morning and the evening.

**NOTE:** Do not exceed the maximum dose of 2 ounces per animal daily. For effective fly control, Vapona Show-Coat Dairy Cattle Spray should be used as directed. It is highly toxic to flies but loses its strength within the day and no insecticidal residue is built up. A well planned fly control program that includes the 2 spray applications per day as directed will give results.

Do not allow the spray mist to contaminate milk or milking utensils. Cover feed and feed troughs or remove before spraying.

**LIMITED WARRANTY:** Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

Exclusion of Other Warranties and Remedies. Except where such disclaimers and exclusions are specifically prohibited by applicable law. THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said

limited warranty, BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER  
FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFAC-  
TURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

953