

GORDON'S



broccoli, eggplant, peppers, and tomatoes are grown, as well as seeded areas of beans, corn, squash, and melons. Do not waste Slug-Killer. One pound will treat 1,000 to 1,500 square feet.

NOTE THESE CAUTIONS: Do not apply to or contaminate foliage or other edible plant parts. Use only on soil in which the listed crops are grown.

SLUGS, SNAILS, AND CUTWORMS: Sprinkle Slug-Killer over the

ACCEPTED

AUG 19 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT CONTROL ACT
FOR FEDERAL TOXIC POISON REGISTER

slug-killer

PBI

MANUFACTURED BY
GORDON CORPORATION

KANSAS CITY, KANSAS 64118

GORDON'S



slug-killer

children and domestic animals. Dispose of container promptly when empty.

ANTIDOTE FOR ARSENIC POISONING: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then give two tablespoons of Epsom Salts or Milk of Magnesium in water and plenty of milk and water. Have victim lie down and keep quiet.

CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING.

MANUFACTURED BY
PBI - GORDON CORPORATION

KANSAS CITY, KANSAS 64110

\$2.69

DIRECTIONS FOR USE

WHERE TO USE SLUG-KILLER: Slug-Killer may be used on the soil throughout all ornamental flower plantings, in most soil areas around home foundations and porches, and on vegetable garden soils where spring plant-sets of cabbage, cauliflower,

garden soil at spring planting time to protect plants against pest damage. If pests are present, treat the entire area where they are found. Do not apply to plants. If soil is dry, water the area lightly before or after application. Repeat treatment in three to four weeks, or as needed for control of pests. For cutworms, place Slug-Killer in the plant rows to give worms easy access to the bait. Repeat baiting as needed for control.

STRAWBERRY ROOT WEEVIL: In home gardens apply to the soil surface around the base of young plants in the spring (April through May) and again in late July through August. Do not apply directly to plants or within six days of harvest of berries. Do not contaminate edible parts.

E.P.A. Reg. No. 2217-454-AA

605-973

slug-killer

kills snails, slugs, and cutworms

one pound baits up to 1,500 sq. ft.

ACTIVE INGREDIENTS:	
Calcium Arsenate	5.00%
Metalddehyde	3.25%
INERT INGREDIENTS	91.75%
Total	
100.00%	
Arsenic expressed as metallic	
Water Soluble Arsenic (As)	
1.88%	
1.88%	

DANGER: Keep Out of Reach of Children and Domestic Animals

See Right Panel for Antidote and Additional Precautions

NET WEIGHT FIVE POUNDS

\$2.69

slug-killer

kills snails, slugs, and cutworms
one pound baits up to 1,500 sq. ft.

ACTIVE INGREDIENTS:	
Calcium Arsenate	5.00%
Metaldhyde	3.25%
INERT INGREDIENTS	91.75%
Total	100.00%
Arsenic expressed as metallic	
Water Soluble Arsenic (As)	1.88%
	1.88%

DANGER: Keep Out of Reach of Children and Domestic Animals

See Right Panel for Antidote and Additional Precautions

NET WEIGHT FIVE POUNDS

STOP! READ CAREFULLY

DANGER: May be fatal if swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of

IMPORTANT NOTE: Keep Slug-Killer away from dogs, pets, and children. Use and store under conditions where there will not be a hazard. Bait may be attractive to dogs. Keep dogs off treated area. Use gloves if Slug-Killer is distributed by hand.

CONTAINER DISPOSAL: Do not reuse container. Destroy it by perforating or crushing. Bury or discard in a safe place.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

605-973