CAUTION

Harmful if swallowed. Avoid breathing of dust or spray mist. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs.

This product will kill fish. Keep out of lakes, streams and ponds.

Do not reuse empty container. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.

NOTICE: Seller warrants that this product conforms to the ingredient statement of the label.



ALFA-SPRAY

ACTIVE INGREDIENTS:	•
Methoxychlor* Technical	23.787%
Malathion (O. O-dimethyl dithiopho	sphate
of diethyl mercaptosuccinate)	23.807%
Xylene	43.510%
INERT INGREDIENTS	
TOTAL	100.000%
*23,787 methoxychlor, technical is e	quivalent to
20.814% 2,2-bis (p-methoxyphenyl)-1, 1	, 1-trichloros-
thane and 2.973% other isomers	and reaction

READ ENTIRE LABEL, USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTION.

Keep Out of Reach of Children See Side Panels for Additional Cautions NET CONTENTS ONE GALLON

products.

GORDON CORPORATION

DIRECTIONS FOR USE

AERIAL APPLICATION: Use recommended amount of ALFA-SPRAY for each specific crop with a minimum of 3 gallon of water per acre consistent with crop growth and good coverage.

ALFALFA: For control of alfalfa weevil, spittlebug, lecthopper and aphis.

FIRST CUTTING: Apply 2-3 quarts per acre as soon as weevil damage appears in the field. Early application is desired to control the weevil when they are small. Do not apply within 7 days of harvest. Do not apply after full bloom.

STUBBLE SPRAYING: In areas where weevil appears at cutting time. Remove the first cutting and apply 2-3 quarts per acre to the stubble to control veevil larvae and aphis

भागामा स्थापक जास्यापक करू

Track.

Our Name Is Your Guarantee

Quality

机械化

ACCEPTED

SEP4 1969
UNDER LAGENTICIDE PRINCIPE AND ROLENTICIDE AND ROLENTICIDE ACT
FOR ECONOMIC POISON REGISTER.
ED UNDER NO. 2217-470

221-(F0