

Gordon's Weeder Granules

controls weeds & grasses around shrubs, flower beds and vegetables.

ACTIVE INGREDIENT
 Trifluralin 2,6-dinitro-
 N,N-dipropylacetamide
 INERT INGREDIENTS

1.47%
 98.53%
 TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for additional precautionary information.

NET WEIGHT ONE POUND

EPA REG. NO. 2217-480
 EPA EST. NO. 2217-KS-1

M'd by **G** pbi/gordon CORPORATION
 KANSAS CITY, KANSAS 64116

BEST DOCUMENT AVAILABLE

ACCEPTED

OCT 27 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act
 registered at
 EPA Reg. No. 2217-480

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

- Hazards to Humans & Domestic Animals -

CAUTION: Harmful if swallowed or absorbed through the skin. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. In case of contact with eyes, flush with plenty of water and get medical attention. If on skin, wash with soap and water.


- Environmental Hazards -

Do not apply directly to water. Use with care when applying in areas adjacent to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL: Do not store where children have access. Do not reuse empty container. Wrap and put in trash collection.



2217-480



USES

GORDON'S WEEDER GRANULES provides long-lasting control of many annual grasses and broad leaf weeds. It kills the weed seeds as they germinate in the soil. Therefore, to be effective, the product must be applied to the soil and mixed into the top 1 to 3 inches of soil. Since Weeder Granules mostly attack germinating seeds, any existing weeds should be removed.
Application to Ornamental Trees, Shrubs & Flowers: In addition to roses and other flowering shrubs, Weeder Granules can be applied under the following established flowers:

Established Flowers			
Achillea	Centaura	Gladiolus	Periwinkle
Ageratum	Chrysanthemum	Hollyhock	Petunia
Arctost	Clematis	Isis	Phlox
Asier	Delila	Lobelia	Portulaca
Balsam	Dianthus	Lupinus	Rudbeckia
Calendula	Dimorphotheca	Mirigold	Sakia
California Poppy	Forget-Me-Not	Morning Glory	Scabiosa
Campanula	Four O'Clock	Nasturtium	Shasta Daisy
Carnation	Gallardia	Nicotiana	Snapdragon
			Stocks
			Snow on the Mountain
			Sunflower
			Sweet Alyssum
			Sweet Pea
			Sweet William
			Zinnia

NOTE: Do not use on ground cover as injury may occur. Also, Gladiolus corms less than 1 inch in diameter may be injured.
Directions for Mixing into Soil: Apply Weeder Granules to established trees, ornamentals, roses and flowers that are 2 to 3 inches or more in height. Apply uniformly at the recommended rate. The rate of application varies with soil type. Use the table below to determine the proper rate for your type of soil. Mix into the soil surface with hand rake being careful not to disturb the roots of established plants. Sprinkle cultivation during the growing season will not reduce effectiveness when Weeder Granules have been adequately mixed into the soil. When it is not possible to mix Weeder Granules into the soil surface, they should be spread on the surface at the rate of 1 pound per 160 sq. ft. and water thoroughly. When applied at this rate, do not mix into the soil. Remove all existing grasses and weeds prior to application since Weeder Granules control weeds from seed only.

Application Chart—Ornamentals

Soil Type	Rate of Application
Light Sandy	1 lb. to 1,280 sq. ft.
Medium Loam	1 lb. to 960 sq. ft.
Heavy Silt or Clay	1 lb. to 640 sq. ft.

Application Chart—Vegetables

Apply Weeder Granules according to the following chart:

Vegetable Crop	Before Planting	After Planting
Vegetables from Seed:	Apply anytime up to 10 weeks before planting these vegetables from seed. Mix into top 3" soil at these rates depending on soil type: Light Sand—1 lb. 1,280 sq. ft. (See A) Medium Loam—1 lb. 960 sq. ft. Heavy Silt or Clay—1 lb. 640 sq. ft. (See B)	NOT RECOMMENDED
Transplants: Broccoli Brussels Sprouts Cabbage Cauliflower Pepper Tomatoes	Apply and mix into soil before transplanting. Do not apply after transplanting.	
Lima Beans from Seed (See C)	Apply anytime up to 10 weeks before planting these vegetables from seed. Mix into light and medium soils at the rate of 1 lb. 1,280 sq. ft. in heavy soils apply at rate of 1 lb. 960 sq. ft.	NOT RECOMMENDED
Seedlings: Cucumbers and Cantaloupe	NOT RECOMMENDED	Mix carefully into soil at rates given for vegetables from seed when plants have 3 or 4 true leaves. Leaves should not be wet.
High Potatoes	NOT RECOMMENDED	Mix into soil at vegetable rates soon after planting.

A: One standard measuring cup contains 1/2 lb. Weeder Granules.
 B: Weeder Granules is not recommended on much soils with very high organic matter content.
 C: Note reduced rate on lima and snap beans.
NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application, are subject to change, Seller makes no claim other than those stated on this label.

WEEDS CONTROLLED

ANNUAL GRASSES:	ANNUAL BROADLEAF WEEDS:
Barnyard grass	Carrot seed
Brachiaria	Chickweed
Bromegrass	Goosefoot
Oat	Knotweed
Crabgrass	Lamb's quarter
Foxtail	Pigweed
	Purslane
	Puncturevine
	String Nettle

BEST DOCUMENT AVAILABLE