



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 29, 2025

Eric D. Smith
Director, Regulatory Affairs
Gordan's Weeder Granules
PBI/Gordon Corporation
P.O. Box 860350
Shawnee, KS 66286

Subject: Label Amendment - Registration Review Mitigation for Trifluralin
Product Name: Gordon's Weeder Granules
EPA Registration Number: 2217-480
Case Number: 474979
Application Dates: March 27, 2020

Dear Eric D. Smith:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Trifluralin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must

submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Tracy Jackson by phone at 202-566-2268, or via email at jackson.tracy@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read 'M. Perch', with a long, sweeping horizontal line extending to the right.

Maryam K. Muhammad-Perch, Team Lead
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

ACCEPTED

09/29/2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 2217-480

TRIFLURALIN

GROUP

3

HERBICIDE

GORDON'S WEEDER GRANULES

ACTIVE INGREDIENT:

Trifluralin: alpha, alpha, alpha-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine..... 1.47%

INERT INGREDIENTS..... 98.53%

TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for Precautionary Statements and First Aid

NET WEIGHT: THREE POUNDS

669/ AP090805

EPA REG. NO. 2217-480

EPA EST. NO. 2217-KS-X

MANUFACTURED BY:

Manufactured For
 **pbi/gordon**
corporation
An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101

Telephone: 1-800-821-7925



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE):

All mixers, loaders, applicators and other handlers must wear:

- ◆ Long-sleeved shirt and long pants,
- ◆ Socks plus shoes,
- ◆ Goggles or face shield, and
- ◆ Chemical resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- ◆ Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- ◆ Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

First Aid	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for treatment advice.
For additional information in case of emergency call toll-free, 1-877-800-5556. Have the product container or label with you when calling a poison control center or doctor or going for treatment.	

ENVIRONMENTAL HAZARDS:

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides 40 CFR Part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Noncrop weed control is not within the scope of the Worker Protection Standard.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container only. Separate pesticides during storage to prevent cross-contamination of other pesticides, fertilizer, food, and feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Paper and plastic bags: Completely empty bag by shaking and tapping sides and bottom to loosen any clinging particles into application equipment. Then dispose of bags in a sanitary landfill, or by incineration, or if allowed by State and local authorities by burning. If burned stay out of smoke.

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GRASS AND BROADLEAF WEEDS CONTROLLED	
Annual Grasses	Annual Broadleaf Weeds
Annual bluegrass	Carpetweed
Annual ryegrass	Chickweed
Barnyardgrass (Watergrass)	Florida pusley (Florida purslane)
Bromegrass	Henbit
Cheat	Knotweed
Crabgrass	Lambsquarters (Goosefoot)
Foxtails	Pigweed
Johnsongrass (from seed)	Puncturevine
Junglerice	Purslane
Sandbur	Stinging Nettle
Signalgrass	
Sprangletop	
Stinkgrass	
Wildcane (shattercane)	
Wild oat (partial control)	
Woolly cupgrass	

- GENERAL INFORMATION -

- ◆ Gordon's Weeder Granules is a pre-emergence herbicide for control of many annual grasses and broadleaf weeds.
- ◆ Gordon's Weeder Granules controls weeds by killing germinating seeds and does not control established weeds.
- ◆ Gordon's Weeder Granules provides effective weed control when applied and incorporated into the soil surface.
- ◆ Gordon's Weeder Granules is recommended for use on a wide variety of ornamentals (trees, shrubs, and flowers), fruits, nuts, and vegetables. See below for specific uses and instructions.

- GENERAL PRECAUTIONS AND RESTRICTIONS -

- ◆ Do not apply this product through any type of irrigation system.
- ◆ Gordon's Weeder Granules must be applied prior to the emergence (pre-emergence) of annual broadleaf and grass weeds. To assure prolonged control (seasonal) of weeds shown below, soil incorporation is essential.
- ◆ Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

WEED RESISTANCE MANAGEMENT

For resistance management, Gordon's Weeder Granules contains a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- ◆ Rotate the use of this product or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- ◆ Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use, and that considers mechanical control methods, cultural

(e.g., timing to favor the turf and not the weeds), biological (weed-competitive varieties) and other management practices.

- ◆ Scout area prior to application to identify the weed species present and their growth stage to determine if the intended application will be effective.
- ◆ Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: 1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; 2) a spreading patch of non-controlled plants of a particular weed species; 3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- ◆ If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- ◆ Contact your local extension specialist or pest control advisor for additional pesticide resistance-management and/or integrated weed-management recommendations for specific types of turf and weed biotypes.
- ◆ For further information or to report suspected resistance, call 877-800-5556.

- APPLICATION INSTRUCTIONS -

Apply Gordon's Weeder Granules to a smooth soil surface to assure uniform distribution. Excessive groundcover such as crop residues or existing weeds can interfere with the incorporation of this product into the soil and should be removed prior to application. Apply Gordon's Weeder Granules prior to the germination of weed seeds or to clean-cultivated, weed-free areas.

Apply Gordon's Weeder Granules in a "ready-to-use" shaker container or with hand held or hand propelled rotary or drop-type granular applicator. Prior to use, calibrate the equipment according to the manufacturer's directions and check frequently during application to be sure equipment is working properly and distributing the granules uniformly.

For small areas, use a "ready-to-use" container or a shaker-type applicator (such as a coffee can with holes punched in the bottom) and distribute the granules uniformly. Do not apply more than the recommended amount.

Gordon's Weeder Granules must be soil-incorporated within 24 hours of application either by mechanical incorporation or at least 0.5 acre inch of water (rainfall, irrigation). If a second incorporation is needed it should be delayed 3 to 5 days after the first incorporation and be completed prior to planting. Application rates for various uses and incorporation methods differ. See below for specific use instructions.

Under normal growing conditions and if applied according to directions, Gordon's Weeder Granules will not harm the treated crop. Crop injury or soil residue may result from over-application. Erratic weed control or crop injury may result from uneven application or improper soil incorporation of Gordon's Weeder Granules. Seedling diseases, cold weather, deep planting, excessive moisture, high salt concentration or drought may weaken crop seedlings and increase the possibility of damage from Gordon's Weeder Granules. Under these conditions, delayed crop development or reduced yields may result. Gordon's Weeder Granules is not recommended for use on muck soils.

Seeded crops (vegetables or flowers) other than those listed on this label for preplant soil incorporation of Gordon's Weeder Granules should not be planted within 5 months following the application.

Use Instructions for Ornamental Trees, Groundcovers, Shrubs, and Flowers:

Apply Gordon's Weeder Granules prior to planting new nursery stock liners, ornamental trees, ornamental groundcovers, woody shrubs, and gladiolus (corms greater than 1 inch diameter). Gordon's Weeder Granules may also be applied to these and other listed ornamentals (see below) after they are established.

Apply Gordon's Weeder Granules to a weed free, smooth, soil surface on ornamental beds or around established plants. Do not apply to established plantings when the foliage is wet.

The broadcast application rate of Gordon's Weeder Granules for these crops is 1.0 ounce per 100 square feet. Optimum weed control performance will be obtained when followed by overhead irrigation (sprinkler or rainfall) within a few hours after application.

ORNAMENTAL TREES

Common Name	Scientific Name
Almond	<i>Prunus dulcis</i>
Apple, Crabapple	<i>Malus spp.</i>
Apricot	<i>Prunus armeniaca</i>
Ash, White	<i>Fraxinus americana</i>
Bald cypress	<i>Taxodium distichum</i>
Birch, European White	<i>Betula pendula</i>
Black gum	<i>Nyssa sylvatica</i>
Cherry	<i>Prunus spp.</i>
Chestnut, Chinese	<i>Castanea mollissima</i>
Cottonwood	<i>Populus deltoides</i>
Dogwood, Flowering	<i>Cornus florida</i>
Dogwood, Kousa	<i>Cornus kousa</i>
Douglas fir	<i>Pseudotsuga menziesii</i>
Fir, Balsam	<i>Abies balsamea</i>
Hemlock, Canada	<i>Tsuga canadensis</i>
Honey locust	<i>Gleditsia triacanthos</i>
Larch, Japanese	<i>Larix kaempferi</i>
Locust, Black	<i>Robinia pseudoacacia</i>
Maple, Norway	<i>Acer platanoides</i>
Maple, Red	<i>Acer rubrum</i>
Maple, Silver	<i>Acer saccharinum</i>
Maple, Sugar	<i>Acer saccharum</i>
Oak, Pin	<i>Quercus palustris</i>
Oak, Red	<i>Quercus rubra</i>
Oak, Scarlet	<i>Quercus coccinea</i>
Peach	<i>Prunus persica</i>
Pine, Austrian	<i>Pinus nigra</i>
Pine, Eastern White	<i>Pinus strobes</i>
Pine, Japanese Black	<i>Pinus thunbergiana</i>
Pine, Loblolly	<i>Pinus taeda</i>
Pine, Red	<i>Pinus resinosa</i>
Pine, Scotch	<i>Pinus sylvestris</i>
Planetree, London	<i>Platanus acerifolia</i>
Plum	<i>Prunus spp.</i>
Redbud, Eastern	<i>Cercis canadensis</i>
Spruce, Colorado	<i>Picea pungens</i>
Spruce, Norway	<i>Picea abies</i>
Spruce, White	<i>Picea glauca</i>
Sweet gum	<i>Liquidambar styraciflua</i>
Sycamore	<i>Platanus occidentalis</i>
Tulip tree	<i>Liriodendron tulipifera</i>
Walnut, Black	<i>Juglans nigra</i>

ORNAMENTAL GROUNDCOVER PLANTING

Common Name	Scientific Name
Aarons beard	<i>Hypericum calycinum</i>
Bellflower, Adriatic	<i>Campanula elatines</i>
Bellflower, Poscharsky	<i>Campanula poscharskyana</i>
Ceanothus	<i>Ceanothus spp.</i>
Coreopsis	<i>Coreopsis spp.</i>
Cotoneaster	<i>Cotoneaster spp.</i>
Coyote Brush	<i>Baccharis pilularis</i>
Crown Vetch	<i>Coronilla varia</i>
Daisy, Trailing African	<i>Osteospermum fruticosum</i>
Fern, Asparagus	<i>Asparagus densiflorus</i>
Gazania	<i>Gazania spp.</i>
Germander	<i>Teucrium chamaedrys</i>
Ice Plant, Largeleaf	<i>Carpobrotus edulis</i>
Ivy, Algerian	<i>Hedera canariensis</i>
Ivy, English	<i>Hedera helix</i>
Lily-of-the-Nile	<i>Agapanthus spp.</i>
Lilyturf, Bigblue	<i>Liriope muscari</i>
Marigold	<i>Tagetes spp.</i>
Myoporum	<i>Myoporum laetum</i>
Plumbago, Dwarf	<i>Ceratostigma plumbaginoides</i>
Rosemary	<i>Rosmarinus officinalis</i>
Rupturewort	<i>Hemiaria glabra</i>
Snow-in-Summer	<i>Cerastium tomentosum</i>
Speedwell	<i>Veronica spp.</i>
St. Johnswort	<i>Hypericum coris</i>
Stonecrop (Sedum)	<i>Sedum spp.</i>
Strawberry, Beach	<i>Fragaria chiloensis</i>
Thrift	<i>America maritima</i>
Verbena	<i>Verbena spp.</i>
Wirevine, Creeping	<i>Muehlenbeckia axillaris</i>
Yarrow, Woolly	<i>Achillea tomentosa</i>
Zoysiagrass	<i>Zoysia tenuifolia</i>

ORNAMENTAL WOODY SHRUBS

Common Name	Scientific Name
Andromeda, Japanese	<i>Pieris japonica</i>
Arborvitae, American	<i>Thuja occidentalis</i>
Azalea	<i>Rhododendron spp.</i>
Barberry, Japanese	<i>Berberis thunbergii</i>
Barberry, Mento	<i>Berberis mentorensis</i>
Boxwood, Common	<i>Buxus sempervirens</i>
Boxwood, Harlands	<i>Buxus harlandi</i>
Boxwood, Littleleaf	<i>Buxus microphylla</i>
Camellia, Japanese	<i>Camellia japonica</i>
Camellia, Sasanqua	<i>Camellia sasanqua</i>
Cherry laurel, American	<i>Prunus caroliniana</i>
Cinquefoil	<i>Potentilla spp.</i>
Cleyera, Japanese	<i>Cleyera japonica</i>
Cotoneaster, Cranberry	<i>Cotoneaster apiculata</i>
Cotoneaster, Zabel	<i>Cotoneaster zabelii</i>
Deutzia	<i>Deutzia spp.</i>
Elaeagnus, Silverberry	<i>Elaeagnus pungens</i>
Euonymus, Spreading	<i>Euonymus kiautschovica</i>
Euonymus, Winged	<i>Euonymus alatus</i>
Euonymus, Wintercreeper	<i>Euonymus fortunei</i>
Firethorn	<i>Pyracantha spp.</i>
Forsythia	<i>Forsythia spp.</i>
Guava, Pineapple	<i>Feijoa sellowiana</i>
Holly	<i>Ilex spp.</i>
Honeysuckle	<i>Lonicera spp.</i>
Indian hawthorn	<i>Raphiolepis indica</i>
Juniper	<i>Juniperus spp.</i>
Laurel, Mountain	<i>Kamia latifolia</i>
Lilac, Common	<i>Syringa vulgaris</i>
Mockorange	<i>Philadelphus spp.</i>
Pittosporum, Japanese	<i>Pittosporum tobira</i>
Privet	<i>Ligustrum spp.</i>
Red cedar, Eastern	<i>Juniperus virginiana</i>
Rhododendron	<i>Rhododendron spp.</i>
Spiraea, Vanhoutte	<i>Spiraea vanhouttei</i>
Viburnum	<i>Viburnum spp.</i>
Weigela	<i>Weigela spp.</i>
Willow	<i>Salix spp.</i>
Yew, Anglojap	<i>Taxus media</i>
Yew, Japanese	<i>Taxus cuspidata</i>
Yew, Southern	<i>Podocarpus macrophyllus</i>

ROSES AND OTHER ESTABLISHED FLOWERS

African Daisy	Four O'Clock	Poppy, California
Aster (perennial)	Gaillardia	Portulaca
Balsam	Gladiolus	Rose
Blackeyed Susan	Golden Glow	Salvia
Calendula	Impatiens	Shasta Daisy
Carnation	Ixora	Snapdragon
Centaurea, Velvet	Lobelia	Snow on the Mountain
Chrysanthemum	Lupine	Stock
Coreopsis	Marigold	Sunflower
Cornflower	Marigold, Cape	Sweet Alyssum
Cosmos	Morningglory	Sweet Pea
Dahlia	Nasturtium	Sweet Sultan
Dianthus	Nicotiana	Sweet William
Dusty Miller	Petunia	Vinca
Floss Flower	Phlox	Yarrow
Forget-Me-Not	Pincushion Flower	Zinnia

Instructions for Fruit, Vegetable, Nut, and Vineyard Crops:

For control of weeds in the listed crops in this section use the following application rate table.

Application Rates	
Soil Texture	Broadcast Application Rates, (ozs./1,000 sq.ft.)
Sand, loamy sand, sandy loam	12.5 fl. oz. (1 lb./1,280 sq.ft.)
Loam, silt, silt loam	16.66 fl. oz. (1 lb./ 960 sq.ft.)
Clay, clay loam, sandy clay	25 fl. oz. (1 lb./ 640 sq.ft.)

Apply and incorporate the indicated amount of Gordon's Weeder Granules to a smooth, weed free, soil surface. Optimum weed control performance will be obtained if Gordon's Weeder Granules is mechanically incorporated, such as with a rake or rototiller, 2 to 3 inches deep within a few hours after application.

Gordon's Weeder Granules is recommended for application to the following crops at the indicated stage of growth.

A. BEFORE PLANTING
Beans (Castor, Dry, Guar, Lima, Mungbean and Snap)
Carrot
Celery, direct seeded or transplant
Chicory/Endive
Cole Crops (Broccoli, Brussels Sprout, Cabbage and Cauliflower), transplant
Fruit and Nut Crops (Almond, Citrus, Grape, Nectarine, Peach, Pecan, Plum, Prune, Walnut)
Greens (Collard, Kale and Mustard)
Okra
Peas (Dry and English)
Pepper, transplant
Radish
Sunflower
Tomato, transplant

B. AT PLANTING
Celery, direct seeded or transplant
C. AFTER PLANTING
Asparagus, established Apply and incorporate prior to spear emergence in the spring.
Celery, direct seeded & transplant
Cucurbit (Cantaloupe, Cucumber & Watermelon) Apply and incorporate between the rows and beneath plants in the 3 to 4 true leaf stage.
Fruit and Nut Crops (Almond, Citrus, Grape, Nectarine, Peach, Pecan, Plum, Prune, Walnut), established Apply 25 to 50 ounces per 1,000 square feet on all soil types. Do not apply to grapes within 60 days of harvest.
Onion (Grown for dry bulb only) Avoid covering exposed bulbs with treated soil or injury may occur.
Potato Apply after planting, prior to crop emergence, or after plants have emerged. Do not cover foliage with treated soil if applied after emergence.
Tomato, direct seeded Apply at thinning after plants are 2 to 3 inches tall to soil between rows and beneath plants and incorporate.

LIMITED WARRANTY AND DISCLAIMER.

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

Advertising claims that may be presented on the container label.

- Stop(s) weeds before they start.
- Stops weeds before they start for a beautiful weed-free garden.
- Stops weeds before they start by preventing weed seeds from germinating.
- For use around flowers, vegetables, trees and shrubs.
- For use on flowers, vegetables and more.
- Easy-grip handle and shaker makes application quick and convenient.
- Easy to accurately apply granules exactly where you want.
- Easy to apply granules.
- Ready-To-Use.
- No mixing, no spills, no mess.
- Pre-emergent herbicide for control of most annual grasses and broadleaf weeds.
- Recommended for use with flowers, roses, shrubs, trees, vegetables, fruits and nuts.
- Use on flowers, roses, shrubs, trees, vegetables, fruits and nuts.
- Control weeds around flowers, roses, shrubs, trees, vegetables, fruits and nuts.
- Easy to apply granular formula only takes a few minutes.
- Don't pull weeds, prevent them!
- Peel open for [See next page for] [See attached booklet for] First Aid and additional Precautionary Statements.
- *Optional graphics that may be presented on the container labeling::*



Sprinkle Weed
Preventer Granules



Activate with ½ -1" of
water or rain, or uniformly
incorporate in soil.



Germinating weed seeds
are shut down before
you see them.

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00480.20201020.amend-proposed-clean
2. **Reason for Issue:** Registration Review comments – Trifluralin