

GORDON'S
prescription for care of lawns
trimecTM
dandelion
 and
broadleaf
 weed killer

ACCEPTED

8-15-69

327-455

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall and again in the spring if needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot spray" weed treatment, however, may be made at any time with a small pressure sprayer spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolens down in the bare spots. Poor weed control will result if spray is applied during drought or just before rain is expected. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE: TO SPRAY ENTIRE LAWN
 1 Pint covers 400 to 600 sq. ft. of lawn.
 Dilute with water as indicated below.

For Pressure Sprayers	Lawn Hose Sprayers	Area to Spray
2 tblsp. in 1/2 gal.	2 tblsp. for 2 gals.	250 sq. ft.
4 oz. (8 tblsp.) in 1 1/2 gal.	8 tblsp. for 6 gals.	1000 sq. ft.
8 oz. in 3 gals.	16 tblsp. for 12 gals.	2000 sq. ft.

CAUTION: Keep Out of Reach of Children
 see back panel for additional cautions
 CONTENTS 1 PINT.

MANUFACTURED BY GORDON CORPORATION KANSAS CITY, KANSAS

19



GORDON'S
 trimec
 dandelion
 and
 broadleaf
 weed killer

ACCEPTED

8-15-69

3-7-68

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall and again in the spring if needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. Spot spray weed treatment, however, may be made at any time with a small pressure sprayer spraying only the weeds at ground level. Wait 3 weeks before pulling new Grass seed or stolons down in the bare spots. Poor weed control will result if spray is applied during drought or just before rain is expected. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE: TO SPRAY ENTIRE LAWN

1 Pint (16 oz.)		to 4,000 sq. ft. of lawn area
Dilute with water as indicated below.		
For Pressure Sprayers	Lawn Hose Sprayers	Area to Spray
2 tblsp. in 1/2 gal.	2 tblsp. for 2 gals.	250 sq. ft.
4 oz. (8 tblsp.) in 1 1/2 gal.	8 tblsp. for 6 gals.	1000 sq. ft.
8 oz. in 3 gals.	16 tblsp. for 12 gals.	2000 sq. ft.

CAUTION: Keep Out of Reach of Children
 see back panel for additional cautions

CONTENTS 1 PINT

MANUFACTURED BY GORDON CORPORATION KANSAS CITY, MISSOURI

NON-EMULSIFIABLE

SORGHUM --Post-Emergence Only: Use 1/3 pint per acre. Sorghums are most tolerant to 2,4-D when 4 to 12 inches high. Spray a check plot first. Do not apply during flowering or early dough stage.

FIELD CORN: To control certain annual or perennials such as field bindweed, or cocklebur, use 1/3 to 1/2 pint per acre, applying at the most susceptible stage of weed growth. Apply post-emergence when 10 to 18 inches high. Some injury to corn is to be expected at any time and may be increased by high temperatures at time of treatment, a storm or cultivation immediately following treatment, or during a stage (which may last for a week or more) just after the leaves unfold, and is increasingly more common following overall sprays as the corn gets taller.

Do not use on light sandy soil. Use drop nozzles when corn is over 10 inches high.

PASTURES, GRASS CROPS, NON-CROP LAND (waste areas, fence rows, drainage ditchbanks, rights-of-way and roadsides): For control of many broadleaved annual weeds such as mustard, cocklebur, sunflower, and certain perennial weeds such as bindweed apply 1 1/3 pints per acre.

Do not allow dairy animals to graze treated areas within seven days after treatment.

Do not spray seedling grasses or when grass is from boot to milk stage.

WHEAT, OATS, BARLEY AND RYE: For the control of susceptible species of weeds such as frenchweed, marshelder, mustard, sunflower, ragweed, cocklebur, hedge bindweed, and similar weeds, use 1/3 to 1/2 pint per acre, and apply at full tiller stage. In general applications

2
WEED
A LOW

Contains 6
ACTIVE INGREDIENTS.
*Mixed octyl ester of
INERT INGREDIENTS

*Equivalent to 62.43 2,4-D
gallon.

NON-

(WILL)


CAUTION: Keep
See Si
CONT

Manufactured
Kansas Cit
customers

to cereal
tillered
jointing
control of
bindweed,
or similar
from 1/2
at late t
stage (the
inches hig
harvest ma

Do not for
fields for
with 2,4-D
Do not use
feed.

ACCEPT
F.
UNDER THE FEDERAL
FUNGICIDE AND ROUEN
FOR ECONOMIC POISON
UNDER NO 2-17



GORDON'S
prescription for care of lawns
trimecTM
dandelion
 and
broadleaf
 weed killer

ACCEPTED

8-15-69

UNDER THE U.S. PATENT OFFICE
 PATENT NO. 2,217,483

Chickweed Henbit Manna Poison Ivy Speedwell Yarrow

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall and again in the spring if needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot spray" weed treatment, however, may be made at any time with a small pressure sprayer spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolens down in the bare spots. Poor weed control will result if spray is applied during drought or just before rain is expected. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE: TO SPRAY ENTIRE LAWN

1 Pint covers up to 4,000 sq. ft. of Lawn Area.
 Dilute with water as indicated below.

For Pressure Sprayers	Lawn Hose Sprayers**	Area to Spray
2 tblsp. in 1/2 gal.	2 tblsp. for 2 gals.	250 sq. ft.
4 oz. (8 tblsp.) in 1 1/2 gal.	8 tblsp. for 6 gals.	1000 sq. ft.
8 oz. in 3 gals.	16 tblsp. for 12 gals.	2000 sq. ft.
1 pint in 6 gals.	1 pint for 24 gals.	4000 sq. ft.

CAUTION: Keep Out of Reach of Children
 see back panel for additional cautions

CONTENTS 1 PINT

MANUFACTURED BY **GORDON CORPORATION** KANSAS CITY, KANSAS