

-TAM-

# TRIMEC 2XF TURF HERBICIDE

FORMERLY KILLEX

**CAUTION: Keep out of Reach of Children.**

SEE SIDE PANEL FOR ADDITIONAL CAUTION  
STATEMENTS

U.S. PATENT NO. 3,284,186

<b>ACTIVE INGREDIENTS:</b>	
*Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	20.95%
**Dimethylamine salt of 2,4-dichlorophenoxy acetic acid	31.85%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	1.97%
<b>INERT INGREDIENTS</b>	<b>45.23%</b>
<b>TOTAL</b>	<b>100.00%</b>
*Equivalent to 2-(2-methyl-4-chlorophenoxy) propionic acid	17.32%
**Equivalent to 2,4-dichlorophenoxy acetic acid	25.46%
***Equivalent to 3,6-dichloro-o-anisic acid	1.64%

**PRIVATE BRANDS, INC.**


300 SOUTH 3RD ST., KANSAS CITY, KANSAS 66118

**NET CONTENTS: 55 GALLONS**

873-569

U.S.D.A. Reg. No. 2217-488

**MANUFACTURING CONCENTRATE**  
For use in formulating a Dry Turf Herbicide.  
Not for resale.



**CAUTION**—Harmful if swallowed. Do not get in eyes, on skin or clothing. In case of contact, wash with soap and water.

Do not contaminate or store near feed or foodstuffs.

Avoid inhalation of spray mist.

**DON'T OVER DOSE!** Follow directions carefully. Do not let spray drift to sensitive plants such as vegetables, legumes, grapes, flowers, cotton or other 2,4-D susceptible plants. Use coarse spray when there is less chance for "wind drift." Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate water to be used for domestic or irrigation purposes. Local Spray conditions will vary. Check local recommendations first. Use separate spray equipment for insecticides and fungicides. Do not reuse empty container. Wash thoroughly with water and detergent. Discard in a safe place. Do not make aerial applications near cotton or sensitive plants.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

T.M. P.B.I., K.C.K.

## DIRECTIONS

Use to "spot-treat" weeds in the lawn. For best results apply when weeds are actively growing. Do not water within 24 hours after spraying.

Shake well. Spot Weeder is designed to permit easy treatment of lawn weeds without bending or stooping. Invert can, aim at weed and press valve. Only a short burst is needed. Overspraying only wastes the spray.

Effects begin to show after a few days and weeds gradually die. Up to 4 weeks may be required for complete kill. Repeat in 3-4 weeks if necessary.

For lawns with many weeds, requiring an overall treatment, spray with TRIMEC Turf Herbicide.

**CAUTION**—Harmful or fatal if swallowed. If swallowed, do not induce vomiting. Call a physician immediately. Vapor harmful. Avoid breathing of vapor or spray mist. Use with adequate ventilation. Avoid contact with skin, eyes, or clothing. In case of contact, flush with plenty of water, and, in case of contact with eyes, get medical attention. Wash after each use.

Avoid spraying susceptible plants (flowers, shrubs, etc.). Avoid accidental drift or application to nearby garden plants, flowers, or shrubs. Do not apply within drip line of desirable plants. Bentgrass, Carpetgrass, and St. Augustine grass may be injured if contacted. Do not contaminate water used for irrigational or domestic purposes. Do not store near seeds, fertilizers, insecticides or fungicides.

**WARNING**—Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130 F may cause bursting. Never throw container into fire or incinerator.

**NOTICE**—Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA Reg. No. 2217-486 AA

670 374

## KILLS

DANDELIONS

PLANTAINS

CHICKWEED

CLOVER

BEDSTRAW

BLACK MEDIC  
(Yellow Clover)

ENGLISH DAISY

CURLED DOCK

HEAL-ALL

GROUND IVY  
(Creeping Charlie)

KNOTWEED

RAGWEED

SHEPHERD'S PURSE

BUCKHORN

CANADA THISTLE

CUDWEED

DOLLARWEED

FLEABANE

FILAREE

HENBIT

LAMBSQUARTERS

MATCHWEED

FLORIDA PUSLEY

MUSTARD

PENNYWORT

PEPPERWEED

PIGWEEED

PRICKLY LETTUCE

POISON IVY

PURSLANE

SHEEP SORREL

OXALIS

SPURGE

YARROW