


2(of 3)

2217-495

DIRECTIONS FOR USE - HOUSEFLIES

2217-499



Residual wall spray: For the control of houseflies, including resistant strains, in dairy barns, hog pens, calf pens, poultry houses, and other farm buildings, apply a 1% residual spray to the ceilings, walls, stanchions, etc. Prepare the spray by mixing 1 pint of Cygon 2-E in 3 gallons of water. Thoroughly wet all fly resting areas to the point of run-off. One gallon of spray will cover 500 to 1,000 square feet of surface.

Repeat applications should be made when necessary.

Remove all animals from buildings when applying residual wall sprays.

CYGON 2-E IS REGISTERED FOR USE AS A RESIDUAL-TYPE INSECTICIDE IN AND AROUND FEDERALLY INSPECTED MEAT-PACKING PLANTS EXCEPT WHERE MEAT FOOD PRODUCTS ARE PROCESSED OR HANDLED

Spot Sprays: For localized housefly control, apply a spray containing  $\frac{1}{2}$  pint of Cygon 2-E in 5 quarts of water with a knapsack or similar type sprayer to areas frequented by flies, such as doorways, around windows, etc.

Repeat applications should be made when necessary. Good sanitation is a necessary part of any effective fly control program.

Manure sprays: For the control of housefly manure, mix  $\frac{1}{2}$  pint Cygon 2-E in 5 quarts of water and apply as a coarse spray or with a sprinkling can to fly-breeding areas, such as poultry droppings in caged-layer houses, garbage dumps and manure piles.

Repeat application as additional manure or garbage is added.

General outside use: For the control of houseflies around homes and recreation areas, garbage cans, animal quarters, food-processing plants, warehouses, loading docks and refuse areas, thoroughly spray exposed surfaces, such as walls, fences, garbage and refuse containers with 1 pint of Cygon 2-E in 3 gallons of water.

Repeat applications should be made when necessary.

DO NOT CONTAMINATE FEED AND FOODSTUFFS, DRINKING FOUNTAINS, MILK AND FEED TROUGHES. DO NOT USE IN MILK-PROCESSING ROOMS, INCLUDING MILK HOUSES AND MILK STORAGE ROOMS. DO NOT USE IN HOMES.

000000

3(of 3)

2217-495

ACCEPTED

2217-495

Jun 13 1969

BY THE FEDERAL INSPECTOR  
FUNGICIDE AND PESTICIDE  
FOR THE U.S. DEPARTMENT OF AGRICULTURE  
ED IN THE NO. \_\_\_\_\_ SUBJECT  
TO ATTACH DOCUMENTS.

**CAUTION**

**VAPOR IRRITANT - HARMFUL IF SWALLOWED**

**COMBUSTIBLE**

Concentrated material causes eye irritation - Avoid breathing vapor or spray mist - Use only with adequate ventilation - Avoid contact with skin and eyes - Wash thoroughly after handling - Do not contaminate food or feed products - Do not use, pour, spill or store near heat or open flame.

In case of contact with eyes, flush eyes with plenty of water for at least 15 minutes; get medical attention.

NOTE TO PHYSICIANS: This product, upon repeated or prolonged use, may cause cholinesterase inhibition. Atropine is antidotal.

This product is toxic to wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift or spray from areas treated.

**DO NOT STORE BELOW FREEZING TEMPERATURES**

Do not use this product for any uses other than those specified on this label.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

The use of the within contained material for systemically protecting living plants is covered by Patent No. 3,310,242. The selling price includes a royalty for the right of any purchaser of said material to practice the methods claimed in said patent. A nontransferable license under said patent may be obtained from American Cyanamid Company on current royalty terms whatever the source of material used.

U.S. Patent 2,494,203

0000

0000

1 (of 3)

CYGON™ 2-  
Fly Spray  
SYSTEMIC INSECT

Active Ingredient:

Dimethoate (O,O-dimethyl S-(M-meth  
phosphorothioate) .....

Inert Ingredients .....

(1 gallon contains 2 pound

CONTROLS HOUSE

(14 pt.)... → CAUTION: } Keep Out of Reach of Children  
(10 pt.)..... → { See Side Panels for Antidote

NET CONTENTS:

USDA Reg. No. 2217-X

Manufactured by

Private Brands, Inc., Kansas  
for various customers as sub


\* trademark of American Cyanamid Company  
U.S. Patent 2,424,233

ACCEPTED

2(of 3)

DIRECTIONS FOR USE - HOUSEFLIES

227-499



Residual Wall Sprays: For the control of houseflies, including resistant strains, in dairy barns, hog pens, calf barns, poultry houses, and other farm buildings, apply a 1% residual spray to the ceilings, walls, stanchions, etc. Prepare the spray by mixing 1 pint of Cygon 2-E in 3 gallons of water. Thoroughly wet all fly resting areas to the point of run-off. One gallon of spray will cover 500 to 1,000 square feet of surface.

Repeat applications should be made when necessary.

Remove all animals from buildings when applying residual wall sprays.

CYGON 2-E IS NOT TO BE USED AS A RESIDUAL-TYPE INSECTICIDE IN AND AROUND FEDERALLY INSPECTED MEAT-PACKING PLANTS EXCEPT WHERE MEAT FOOD PRODUCTS ARE PROCESSED OR HANDLED

Spot Sprays: For localized housefly control, apply a spray containing 1/2 pint of Cygon 2-E in 5 quarts of water with a knapsack or similar type sprayer to areas frequented by flies, such as doorways, around windows, etc.

Repeat applications should be made when necessary. Good sanitation is a necessary part of any effective fly control program.

Manure Sprays: For the control of housefly manure, mix 1/2 pint Cygon 2-E in 5 quarts of water and apply as a coarse spray or with a sprinkling can to fly-breeding areas, such as poultry droppings in caged-layer houses, garbage dumps and manure piles.


Repeat application as additional manure or garbage is added.

General Sprays: For the control of houseflies around homes and recreation areas, garbage cans, animal quarters, food-processing plants, warehouses, loading docks and refuse areas, thoroughly spray exposed surfaces, such as walls, fences, garbage and refuse containers with 1 pint of Cygon 2-E in 3 gallons of water.

Repeat applications should be made when necessary.

DO NOT USE IN MILK MILKING AREAS, MILK STORAGE ROOMS, MILK PROCESSING PLANTS, MILK CONDENSING PLANTS, MILK CANNERIES, MILK STORAGE TANKS, MILK TRUCKS, MILK PROCESSING PLANTS, INCLUDING MILK HOUSES AND MILK STORAGE ROOMS. DO NOT USE IN HOMES.

3(of 3)



CAUTION!

VAPOR HARMFUL - HARMFUL IF SWALLOWED

COMBUSTIBLE

Concentrated material causes eye irritation - Avoid breathing vapor or spray mist - Use only with adequate ventilation - Avoid contact with skin and eyes - Wash thoroughly after handling - Do not contaminate food or feed products - Do not use, pour, spill or store near heat or open flame.

In case of contact with eyes, flush eyes with plenty of water for at least 15 minutes; get medical attention.

HEALTH HAZARDS: This product, upon repeated or prolonged use, may cause cholinesterase inhibition. Atropine is antidotal.

This product is toxic to wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift or spray from areas treated.

DO NOT STORE BELOW FREEZING TEMPERATURES

Do not use this product for any uses other than those specified on this label.

NOTES: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, seller makes no claims other than those stated on this label.

The use of the within contained material for systemically protecting living plants is covered by Patent No. 3,410,242. The selling price includes a royalty for the right of any purchaser of said material to practice the methods claimed in said patent. A nontransferable license under said patent may be obtained from American Cyanamid Company on current royalty terms whatever the source of material used.

U.S. Patent 2,494,205