



LIVESTOCK SPRAY CONCENTRATE

Contains .85 Lbs. Vapona, 1.25 Lbs. Ciodrin per Gal.

2217-499
ACCEPTED
 Feb 4 1971
 UNDER THE FEDERAL INSECTICIDE ACT
 FOR ECONOMIC PEST CONTROL
 ED UNDER NO. _____
 TO ATTACH _____

ACTIVE INGREDIENTS:

*2,2-Dichlorovinyl dimethyl phosphate	9.35%
*Related Compounds	.70%
Dimethyl Phosphate of Alpha-methyl-benzyl 3-Hydroxy-cis-crotonate	14.78%
Heavy Aromatic Naphtha	64.67%
INERT INGREDIENTS	10.50%
TOTAL	100.00%



DANGER: Keep Out of Reach of Children
 See Side Panels for Additional Cautions
 See Antidote Statement and Other Warnings on Side Panel
 *Equivalent to .85 lbs. of Vapona per gallon
CONTENTS 1 GALLON

964-1270

MANUFACTURED BY
GORDON CORPORATION
 KANSAS CITY, KANSAS

DIRECTIONS FOR USE

LIVESTOCK SPRAY CONCENTRATION is formulated to enable the user to prepare dilutions of varying concentrations. Refer to the directions below for proper dilution rates. Always thoroughly cover all parts of the animal. Do not apply the spray in windy weather.

ANIMALS TO WHICH APPLICABLE

DAIRY CATTLE, BEEF CATTLE—(Face, Horn, Hoof, Hooves, Stable, Mastitis, Gnats, Lice)
 2 qt. in 10 gal. water (1% Vapona and .75% Ciodrin solution). Apply daily using a good hand or automatic mist sprayer. Do not spray the animal's hair in wet weather. Thoroughly cover all parts of the animal, including the udder. Do not exceed 2 ounces daily per animal. Do not spray 1/2 hr. for younger animals. Do not contaminate feed, food, or drinking water.

HOUSEhold ANIMALS—(Face, Horn, Hoof, Hooves, Stable, Mastitis, Gnats, Lice)
 2 qt. in 10 gal. water (1% Vapona and .75% Ciodrin solution). Apply daily using a good hand or automatic mist sprayer. Do not spray the animal's hair in wet weather. Thoroughly cover all parts of the animal, including the udder. Do not exceed 2 ounces daily per animal. Do not spray 1/2 hr. for younger animals. Do not contaminate feed, food, or drinking water.

GOATS, HORSE BARN, STABLE, HORSE, HOOF, HOVES, STABLE, Mastitis, Gnats, Lice
 2 qt. in 10 gal. water (1% Vapona and .75% Ciodrin solution). Apply daily using a good hand or automatic mist sprayer. Do not spray the animal's hair in wet weather. Thoroughly cover all parts of the animal, including the udder. Do not exceed 2 ounces daily per animal. Do not spray 1/2 hr. for younger animals. Do not contaminate feed, food, or drinking water.

BEEF & DAIRY CATTLE
 1/2 pt. in 5 gal. water (1% Vapona and .75% Ciodrin solution). Apply as a coarse, wet spray. Spray premises such as inside and outside of barn walls, eaves, fences, window sills, alleyways and other cracks or openings. Repeat as necessary. Use approximately 1 qt. per 100 sq. ft. Improved house and farm fly control may be achieved by adding 1 lb. of sugar per 10 gallons of spray solution. Do not contaminate feed, food, or drinking water.

BARN, DAIRY BARN, LOADING SHEDS, HQ HOUSES, HORSE BARN, STABLE, PUMP HOUSES, UTILITY BUILDINGS, MILK HOUSES—(Face, Horn, Hooves, Stable, Gnats, Mosquitoes)
 2 qt. in 10 gal. of water (1% Vapona and .75% Ciodrin solution). Apply as a coarse, wet spray. Spray premises such as inside and outside of barn walls, eaves, fences, window sills, alleyways and other cracks or openings. Repeat as necessary. Use approximately 1 qt. per 100 sq. ft. Improved house and farm fly control may be achieved by adding 1 lb. of sugar per 10 gallons of spray solution. Do not contaminate feed, food, or drinking water.

964-1270

WARNINGS

- Poisonous if swallowed, inhaled or absorbed through skin or eyes.
- Do not get on skin, in eyes, or on clothing.
- Do not breathe fumes.
- Avoid prolonged exposure to fumes.
- Wear natural rubber gloves, protective clothing and goggles when handling the concentrate to prepare emulsions.
- Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for DDVP (Vapona insecticide).
- Keep all unprotected persons out of operational area.
- In case of contact, wash immediately with soap and water.
- Wash hands, arms and face thoroughly with soap and water before eating or smoking.
- Wash all contaminated clothing with soap and hot water before reuse.
- Keep out of reach of children.
- Store at temperature above 32° F.

ADDITIONAL

- Do not contaminate milk or streams. Do not contaminate feed, water or feedstuffs. Keep out of reach of children. Transport the product immediately to the nearest physician.
- **INTERNAL:** If material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving salt water (1 tablespoon of salt to a glass of water). Never give anything by mouth to an unconscious person.
- **EXTERNAL:** If spilled on the skin, immediately remove person from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If material gets into the eyes, wash immediately with running water for at least 10 minutes, get medical attention.
- **NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 2217-499

964-1270


GORDON'S NET CONTENTS ONE GALLON

®

DIRECTIONS FOR USE

LIVESTOCK SPRAY CONCENTRATION is formulated to enable the user to prepare dilutions of varying concentrations. Follow specific directions below for proper dilution rates. Apply to thoroughly cover all parts of the animals. Do not apply the spray in poultry houses.

ANIMALS OR USE, INSECTS

DAIRY CATTLE, BEEF CATTLE—Flies (Face, Horn, House, Stable), Mosquitoes, Gnats, Lice:

2 qts. in 10 gals. water (.5% Vapona and .75% Ciodrin solution). Apply daily using a good hand or automatic mist sprayer so that the animal's hair is wet. Spray to thoroughly cover all parts of the animal, including the legs. Do not exceed 2 ounces daily per adult animal. Reduce dosage 1/2 for younger animals. Do not contaminate food, feed or drinking water.

HORSES (INCLUDING RACE HORSES)—Flies (Face, Horn, House, Stable), Mosquitoes, Gnats, Lice:

2 qts. in 10 gals. of water (.5% Vapona and .75% Ciodrin solution). Apply daily using a good hand or automatic mist sprayer so that the animal's hair is wet. Spray to thoroughly cover all parts of the animal, including the legs. Do not exceed 2 ounces daily per adult animal. Reduce dosage 1/2 for smaller or younger animals. Do not contaminate food, feed or drinking water.

GOATS, HOGS, SHEEP—Flies (Face, Horn, House, Stable), Mosquitoes, Gnats, Lice:

2 qts. in 10 gals. water (.5% Vapona and .75% Ciodrin solution). Apply daily using a good hand or automatic mist sprayer so that the animal's hair is wet. Spray to thoroughly cover all parts of the animal, including the legs. Do not exceed 1 ounce daily per adult animal. Reduce dosage 1/2 for younger animals. Do not contaminate food, feed, or drinking water.

BEEF & DAIRY CATTLE—Face Fly:

1/2 pt. in 5 pts. water (.5% Vapona and .75% Ciodrin solution). Using a calibrated sprayer, apply 5 ml. (1 tsp.) of spray to the face and a total of 5-10 ml. (1-2 tsp.) to the back and sides of the animal 3 times per week. Improved control may be achieved by the addition of 1 lbs. of sugar per pint of spray solution. Do not contaminate food, feed, or drinking water.

BARN, DAIRY BARN, LOADING SHEDS, HOG HOUSES, HORSE BARN, STABLES, PUMP HOUSES, UTILITY BUILDINGS, MILK HOUSES—Flies (Face, Horn, House, Stable), Gnats, Mosquitoes:

2 qts. in 10 gals. of water (.5% Vapona and .75% Ciodrin solution). Apply as a coarse, wet spray. Spray premises such as inside and outside of barn walls, eaves, fences, window sills, alleyways and other resting or breeding places. Repeat as necessary. Use approximately 1 quart per 1000 sq. ft. Improved house and face fly control may be achieved by adding 1 lb. of sugar per 10 gallons of spray solution. Do not contaminate food, feed, or drinking water.

964-1270


LIVESTOCK SPRAY CONCENTRATE

Contains .85 Lbs. Vapona, 1.25 Lbs. Ciodrin per Gal.

ACTIVE INGREDIENTS:

*2,2-Dichlorovinyl dimethyl phosphate	9.35%
*Related Compounds	.70%
Dimethyl Phosphate of Alpha-methylbenzyl 3-Hydroxy-cis-crotonate	14.78%
Heavy Aromatic Naphtha	64.67%
INERT INGREDIENTS	10.50%

TOTAL 100.00%



— POISON —

DANGER: Keep Out of Reach of Children

See Side Panels for Additional Cautions
See Antidote Statement and Other Warnings on Side Panel

*Equivalent to .85 lbs. of Vapona per gallon

CONTENTS 1 GALLON

964-1270

WARNING

Poisonous if swallowed, inhaled or absorbed through skin or eyes.

Do not get on skin, in eyes, or on clothing.

Do not breathe fumes.

Avoid prolonged exposure to fumes.

Wear natural rubber gloves, protective clothing and goggles when handling the concentrate to prepare emulsions.

Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for DDVP (Vapona Insecticide).

Keep all unprotected persons out of operational area.

In case of contact, wash immediately with soap and water.

Wash hands, arms and face thoroughly with soap and water before eating or smoking.

Wash all contaminated clothing with soap and hot water before reuse.

Keep out of reach of children.

Store at temperatures above 32° F.

ANTIDOTE

Do not contaminate lakes or streams. Do not contaminate feed, water or foodstuffs. Keep out of reach of children. Transport the patient immediately to the nearest physician.

INTERNAL: If material has been swallowed, induce vomiting immediately. This may be done by introducing a finger into the throat or by giving salt water (1 tablespoon of salt to a glass of water). Never give anything by mouth to an unconscious person.

EXTERNAL: If spilled on the skin, immediately remove person from the vicinity of the insecticide, remove all contaminated clothing and wash skin with soap and running water. If material gets into the eyes, wash immediately with running water for at least 10 minutes, get medical attention.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 2217-499

964-1270



MANUFACTURED BY
GORDON CORPORATION
KANSAS CITY, KANSAS