

Use herbicides safely.
Do not overdose.
Read DIRECTIONS FOR USE
carefully.

Net Wt. 35 Lbs.

TREATS
Up To 5000 Sq. Ft.



ACCEPTED
July 17-1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND REFRIGERANT ACT
REGISTERED SUBJECT
EPA REG. NO. 2217-594

ST. AUGUSTINE GRASS LAWN WEEDER & FEEDER

For Use On St. Augustine Grass
And Other Grasses For
Control Of Growing Broadleaf Weeds

CAUTION: Keep Out Of Reach Of Children.

See Back Panel for Additional Cautions

WEEDS AS IT FEEDS

ACTIVE INGREDIENTS:	
*2,4-dichlorophenoxyacetic acid	0.0820%
**2-(2-methyl-4-chlorophenoxy) propionic acid	0.2952%
***Dicamba (3,6-dichloro-o-anisic acid)	0.0295%
INERT INGREDIENTS	99.5933%
TOTAL	100.0000%

Used at 35 lbs. on 5,000 sq. ft. equivalent to:
*2,4-D ACID 0.25 lbs./acre or 1.64 lbs./ton
**MCPP ACID 0.90 lbs./acre or 5.904 lbs./ton
***Dicamba ACID 0.09 lbs./acre or 0.59 lbs./ton

U.S. Patent No. 3,284,186

E.P.A. Reg. No. 2217-594

MANUFACTURED BY
GORDON CORPORATION
KANSAS CITY, KANSAS

NET WEIGHT 35 POUNDS COVERS UP TO 5000 SQUARE FEET

GORDON'S

ST. AUGUSTINE GRASS LAWN WEEDER & FEEDER

For Control of Growing Broadleaf Weeds In
St. Augustine and Other Grasses

16-4-8

GUARANTEED ANALYSIS

TOTAL NITROGEN, not less than	16.00%
NITRATE NITROGEN, not less than	None
AMMONIACAL NITROGEN, not less than	2.66%
WATER SOLUBLE ORGANIC NITROGEN, not less than	8.67%
WATER INSOLUBLE NITROGEN, not less than	4.67%
AVAILABLE PHOSPHORIC ACID, not less than	4.00%
INSOLUBLE PHOSPHORIC ACID, not less than	0.50%
WATER SOLUBLE POTASH, not less than	8.00%
TOTAL AVAILABLE PRIMARY PLANT FOOD, not less than	28.00%
CHLORINE, not more than	1.00%

DIRECTIONS FOR USE

GORDON'S ST AUGUSTINE GRASS LAWN WEEDER AND FEEDER destroys growing broadleaf weeds as it feeds St. Augustine, Bermuda, Centipede, Zoysia and other turf grasses.

CONTROLS THESE WEEDS: Florida Pusley, Black Medic, Chickweed, Clover, Cocklebur, Dandelion, Dog, Horbit, Knotweed, Lambquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poisson Ivy, Purslane, Ragweed, Spurge, Wild Lettuce, and most other broadleaf weeds.

TIME TO APPLY: Apply any time broadleaf weeds are present except avoid treating St. Augustine grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or watering on weed foliage at time of application aids control. Do not water for 24 to 48 hours after application.

HOW MUCH TO APPLY: Use 35 lbs. on 5,000 sq. ft. for St. Augustine Grass and Bentgrass. Over-application may cause temporary discoloration. On other grasses such as Bluegrass, Bermuda, Zoysia, Fescue, and Bahia use 35 lbs. on 3,500 sq. ft. of lawn.

SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

Make of Spreader	SPREADER SETTINGS	
	Bentgrass & St. Augustine Grass	Other Grasses
K-Mart	3 1/4	4 1/4
Lawn Beauty	7 1/2	8 1/4
Scotts	4 1/4	5
Sears	5 1/4	6 1/2
Wards	6	6 1/2

CAUTION: Keep out of reach of children. Keep away from feed and food products. Do not take internally. Avoid inhalation or contact with eyes. Do not use on dichondra lawns. Do not re-use empty bag. Destroy when empty. Do not apply in areas underlaid by roots of desirable trees and shrubs.

Avoid drift to susceptible desirable plants. Do not use on newly seeded lawns.

Store in a dry place, away from plants, seeds, fertilizers, insecticides, or fungicides.

515-672