

**GORDON'S**



# crabgrass killer

AMOUNT OF CRABGRASS KILLER		Crabgrass with Seed Heads		WATER IN SPRAYER TO BE TREATED	
Young Crabgrass or 90° Weather	Fl. oz.	Tbs.	Fl. oz.	Gallons	Sq. Feet
2	1	3	1 1/2	1	250
4	2	6	3	2	500
6	3	9	4 1/2	3	750

**HOSE-END SPRAYER:**

Use a hose-end sprayer designed for treating lawns. Measure the recommended amount of chemical into bottle on sprayer and add water into bottle to desired level.

AMOUNT OF CRABGRASS KILLER		FILL BOTTLE TO	AREA TO TREAT
Young Crabgrass	Older Crabgrass		

Recommended rates of application in the presence of Kentucky, Merion Blue and Zoysia Grasses. Under ideal weather and turf conditions, such as cool days and moist turf, it is also selective in the presence of sensitive grasses, such as Fescues and Bents. Discoloration may occur in Fescues and Bent grass areas, hence only the "Young Crabgrass" dosage is recommended when these grasses predominate. This will require a number of treatments if spraying is delayed until crabgrass is mature in these areas. Will injure St. Augustine, Bahia, Carpet and Centipede grasses.

**CAUTION:** Avoid contact with skin or breathing of spray mist. Avoid contamination of food or foodstuffs. Keep away from children and domestic animals. Wash hands after using. Harmful if swallowed. Keep livestock off treated areas.

MANUFACTURED BY  
**GORDON CORPORATION**  
KANSAS CITY, KANSAS

UNION  
REGISTERED  
TRADE  
MARK  
UNDER NO. 2

**DIRECTIONS FOR USE**  
**WHERE TO USE:** New Lawn Weed Killer may be used on Bluegrass, Bermuda, or Zoysia lawns, vacant lots, play grounds, and other similar areas where hard-to-kill weeds are present. This product is especially effective on all types of hard-to-kill weeds such as HENBIT, CHICKWEED, KNOTWEED, SPOTTED SPURGE, OXALIS, BLACK MEDIC, and SHEEP SORREL. In addition to these, TRIMEC controls most other broadleafed lawn weeds, including Dandelion and Plantain, and a partial list is given on the side panel.

Do not allow spray to get on vegetable or ornamental plants. Injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot-Spray" weeds only. Clover present will be damaged if sprayed.

**WHEN TO USE:** Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results mow before treatment. Do not water lawn within 24 hours after treatment.

**CAUTION:** Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpet.



**NEW LAWN WEED KILLER**

Contains \*TRIMEC® Herbicide

1 Pint Covers 2700 Sq. Ft.

**ALSO KILLS: DANDELION, KNOTWOOD, HENBIT, PLANTAINS, SPURGE, POISON IVY, and most other broadleaf weeds.**

Contains: Dimethylamine salts of 2,4-D, MCPP and Dicamba  
**KEEP FROM FREEZING**

**ACTIVE INGREDIENTS:**

• Dimethylamine salt of 2,4-dichlorophenoxypropionic acid	5.900
• Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	8.990
• Dimethylamine salt of Dicamba 3,6-dichloro-o-anisic acid	0.553
<b>INERT INGREDIENTS</b>	84.557
<b>Total</b>	<b>100.000</b>

2,4-dichlorophenoxypropionic acid equivalent 0.42% pounds per gallon or 4.88% Acid.  
 2,4-dichlorophenoxyacetic acid equivalent 0.65 pounds per gallon or 7.465% Acid.  
 3,6-dichloro-o-anisic acid equivalent 0.04 pounds per gallon or 0.459% Acid.

**CAUTION: Keep Out of Reach of Children**  
 See Side Panels for Additional Precautions  
 Registered trademark

MANUFACTURED BY  
**PBI - GORDON CORPORATION**  
 KANSAS CITY, KANSAS

<b>CONTROLS</b>	<b>HENBIT</b>	<b>POISON OAK</b>
BEDSTRAW	KNOTWEED	PURSLANE
BLACK MEDIC	LAMBSQUARTER	RAGWEED
BUCKHORN	LESPEDEZA	SHEPHERDS PURSE
CHICKORY	MALLOW	SPEEDWELL
CHICKWEED	MORNING GLORY	SPURGE
CLOVER	PEPPERGRASS	WILD CARROT
DANDELION	PIGWEEED	WILD LETTUCE
DOCK	PLANTAINS	WILD ONION
GROUND IVY	POISON IVY	YARROW
HEAL-ALL		AND MANY OTHERS

**HOW MUCH TO USE—ENTIRE LAWN**  
**HOSE-END SPRAYER:** Use a hose-end sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2) and spray on area indicated. For example—to spray 1,700 sq. feet, measure 10 ozs. (20 tbsp.) in jar and add water to the 10 gallon level and mix. Turn on hose and spray as evenly as possible over 1,700 sq. feet.

Amt. New Lawn Weed Killer to use: (Tbsp.)	Col. 1 (Fl. Oz.)	Col. 2 Fill Bottle To: (Gallons)	Col. 3 Area to Treat: (Sq. Feet)
2	1	1	170
4	2	2	335
10	5	5	850
20	10	10	1700
32	16	16	2700

**PRESSURE SPRAYERS:** Measure indicated amount of chemical into tank and dilute with water according to the following chart. Use a spray pattern which gives small spray droplets without any fine mist and cover the area indicated.

*WILD GARLIC*

*Our Name Is Your*

*Guarantee*

*of*

*Quality*

ACCEPTED WITH COMMENTS