

G
GORDON'S NET CONTENTS ONE GALLON

DIRECTIONS FOR USE

ORNAMENTAL LAWNS AND TURF (Other than bent grass): Apply TRIMEC FAIRWAY HERBICIDE at the rate of 3¼ to 4 pints per acre in sufficient water for adequate coverage. Maximum kill of weeds will be obtained from spring or early fall application when weeds are actively growing. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Grass seed can be safely sown 3 to 4 weeks after application at recommended rate. Do not use on dichondra, carpetgrass, and St. Augustine grass.

BENT GRASS TURF: In closely mowed bent grass (Putting and bowling greens) apply TRIMEC FAIRWAY HERBICIDE at a maximum rate of 1 fluid ounce per 1,500 square feet preferably in May, or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE: Care should be taken to avoid overdosing bent grasses or injury may result. Large volumes of spray water (i.e., 1 fl. oz. in 5 gal. water per 1,500 sq. ft.) will aid in obtaining uniform coverage. If hand-type sprayers are used it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local overapplication and subsequent turf discoloration or injury.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 2217-517 881-170

trimec
FAIRWAY
BROADLEAF HERBICIDE

Contains New Patented TRIMEC™ Weed Killer.
 U. S. Patent No. 3,284,186

Kills: DANDELIONS KNOTWEED HENBIT
 CHICKWEED PLANTAINS SPURGE
 and Many Other Broadleaf Weeds

Contains: 2,4-D, MCPP and Dicamba
 ONE GALLON COVERS 2-2½ ACRES.

KEEP FROM FREEZING

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 21.85%
 Dimethylamine salt of 2,2-dimethyl-4-chlorophenoxypropionic acid 20.95%
 Dimethylamine salt of Dicamba (3,6-dichloro-2,4,5-triaminobenzoic acid) 1.97%

NET INGREDIENTS:

Total 45.23%
 Total 100.00%

2,4-Dichlorophenoxyacetic acid equivalent 7.66 pounds per gallon
 2,2-Dimethyl-4-chlorophenoxypropionic acid equivalent 7.26 pounds per gallon
 3,6-Dichloro-2,4,5-triaminobenzoic acid equivalent 1.00 pounds per gallon

CAUTION: Keep Out of Reach of Children
 See Back Panel for Additional Cautions

CONTENTS 1 GALLON

WEEDS CONTROLLED: TRIMEC FAIRWAY HERBICIDE is especially effective against weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

Do not allow spray to get on vegetable or ornamental plants since injury may result. Some injury can be expected to turf when used on Centipede or St. Augustine grass lawns. "Spot Spray" weeds only. Clover present will be damaged if sprayed.

Bedstraw	Knotweed	Sheep Sorrel
Black Medic	Lambsquarter	Shepherd's Purse
Buckhorn	Lespedeza	Speedwell
Burdock	Mallow	Spurge
Chicory	Morning Glory	Wild Carrot
Chickweed	Peppergrass	Wild Garlic
Clover	Pigweed	Wild Lettuce
Dandelion	Plantains	Wild Onion
Dock	Poison Ivy	Yarrow
Ground Ivy	Poison Oak	
Heal-all	Purslane	
Henbit	Ragweed	

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solutions near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

881-170



MANUFACTURED BY
GORDON CORPORATION
 KANSAS CITY, KANSAS