
46125 120 20017-519



METHOD OF APPLICATION

Note: The use of this product containing Seed Lube Plus will also lubricate and protect the internal parts of the planter.

USE ONLY ON SEED THAT HAS PREVIOUSLY BEEN TREATED WITH A FUNGICIDE.

Planter-Box Treatment:

Corn

Seed Corn Beetles: Mix 41/2 ounces per bushel of seed as a dry mixture in the planter box as a seed treatment prior to planting.

Corn, Peas, Soybeans and Beans (Lima, Snap, White and Kidney)

Seed Corn Maggots: Mix $1\frac{1}{2}$ -3 oz. per bushel of seed as a dry mixture in the planter box as a seed treatment prior to planting.

Planter-Box Mixing Instructions: Stir seed slightly using a mixing stick. Additional mixing will result from the operation of the planter. DO NOT MIX WITH BARE HANDS.

E.P.A. Reg. No. 2217-519-AA

1. P.A. Est. No 2217-KS-1

511-175



DIAZINON SEED TREATER WITH SEED LUBE PLUS





WARNING: DO NOT USE TREATED SEED FOR FOOD, FEED **OR OIL PURPOSES.**

Diazinon treated seed cannot be used for feed for livestock, or for the preparation of products for human consumption.

WARNING: May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of children. Keep out of reach of domestic animals.

This product is toxic to fish, birds, and other wildlife. Treated seed exposed on soil surface may be hazardous to birds and wildlife. Dispose of excess treated seeds by burial. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy it by burying or burning. Avoid smoke or fumes.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary. Seller makes no claims other than those stated on this label.

511-175

METHOD OF APPLICATION

Note The use of this product containing Seed Lube Plus will Hso lubricate and protect the internal parts of the planter

E NEEN SEEL THAT HAS PRESIDENTLY FEED TREATED CONTRACTOR OF THE

Planter Box Treatment:

Corn

which Core Beetles: Mix 44_2 outlices per bushel of seed as . Try mixture in the planter box as a seed treatment prior to tri g.

Corn, Peas Soybeans and Beans (Lima, Snap, White and Kidney)

send Corn Muggold, Mix 11, 3 oz per bushel of spectus a 11. parture in the planter box as a seed treatment price to plantering

ter box Mixing Instructions Stir Leed Might Parts is rig stick. Additional mixing will result that the the proster DO NOT MIX WITH BARE HAN.

•



DIAZINON SEED TREATER WITH SEED LUBE PLUS

mainating # O.O-diethyl Q-12-1800 phospherothics INGREDIE制法

Diazinone CIBA GEIGY Undergant E

C THUR TAN FROM







WARNING: DO NOT USE TREATED SEED FOR FOOD, FEED OR UIL PURPOSES.

Diaziron treated seed cannot be used for the stor livestock is for the preparation of products for human consumption.

WARNING: May be fatal if swallowed. May be at corted the out Chin. 10) not breathe dust. Aboid contact with skin leges and on th ing. Veash thoroughly after handling. Acoud contain nation of form and find Keep out it mark if history. King out if more it come to animals

The product is toxic to fish birds and ther wildlife. Treated seed reading that see surface may be trazar than the total and we fife The forcess treated seeds by teach the standard water to clearung of equipments in displace to we tee Applicate production by an an other to the state

Despect requires a reference and see presented at the beginning of the reference of the STOCK C future

NOTE: Solid workers that this preciable offerns to the agree end statements in the later. Since conditions of use such is weather compatibility with other demicals and our ditude of application inquipment websitary. Second concern the second second