



Use higher rates for taller alfalfa and clover.

BARLEY, OATS, RYE, WHEAT (Grain forms)	1½-2 Pts.	7 days
English Grain Aphid, Greenbugs, Cereal Leaf Beetle		
Apply as needed and use in sufficient water for good coverage.		
Armyworms	2 Pts.	7 days
PEPPERMINT, SPEARMINT —Aphids, Spider Mites, Leafhoppers	1½ Pts.	7 days
Use in sufficient water for good coverage.		
RICE (Grain form)—Rice Leaf Miner	2½ Pts.	7 days
Apply soon after first rice plants appear on surface of the water.		
Rice Stink Bug	1-1½ Pts.	7 days
Apply by airplane during early milk or dough stage and repeat as necessary. Mix with approximately 3 to 10 gallons per acre, consistent with crop growth and good coverage.		
SUGAR BEETS —Aphids, Spider Mites	1½-2 Pts.—	
Apply as needed and repeat if necessary. Do not use within 7 days of harvest if tops are to be used for feed.		
TOBACCO —Aphids (Field)	1-1½ Pts.	7 days***
Plant Beds, 2 pts. per 100 gals. of water but not within 3 days of transplanting. Apply at the rate of 5 gals. of finished spray per 100 sq. ft.		
ASPARAGUS —Thrips	1½-2 Pts.	1 day
Apply on foliage as needed.		
BEANS —Mexican Bean Beetles, Leafhoppers, Spider Mites, Japanese Beetle	1½-2 Pts.	1 day***
BEETS —Aphids	1½-2 Pts.	7 days***
BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE, BROCCOLI, KALE, KOHLRABI, TURNIP, MUSTARD GREENS, CARROTS	1½-2 Pts.	7 days***

MALATHION

CONTAINS 5 LBS. MALATHION PER GALLON

ACTIVE INGREDIENTS:

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	55.2%
Heavy Aromatic Naphtha	35.5%

INERT INGREDIENTS	9.3%
Total	100.0%

CAUTION: Keep Out of Reach of Children
See Side Panels For Additional Cautions

CAUTION

HARMFUL IF SWALLOWED. AVOID PROLONGED BREATHING VAPOR OR SPRAY MIST. AVOID PROLONGED OR REPEATED CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF

2027-1971
 2217-520



MALATHION

CONTAINS 5 LBS. MALATHION PER GALLON

ACTIVE INGREDIENTS:

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate) 55.2%
 Heavy Aromatic Naphtha..... 35.5%

INERT INGREDIENTS 9.3%
 Total..... 100.0%

CAUTION: Keep Out of Reach of Children
 See Side Panels For Additional Cautions.

CAUTION

IF SWALLOWED, AVOID PROLONGED BREATHING VAPOR OR SPRAY MIST. AVOID PROLONGED

2217-520
 025-6172

CROPS AND INSECTS	Rate per 100 Gals. of Water	INTERVAL (Last Spray to Harvest)
PECANS —Spider Mites, Aphids	1-2 Pts.	0 days***
GRAPEFRUIT, LEMONS, LIMES, ORANGES, TANGERINES, TANGELUS, KUMQUATS California Red Scale, Yellow Scale, Purple Scale, Black Scale (single brooded), Soft Scale, Citricola Scale	1-2 Pts.	7 days***
Florida Red & Purple Scales (light and medium infestations)	2 Pts.	7 days***
Thrips	2½ Pts. per 200 gals per acre	7 days***
Do not apply when trees are in bloom.		
BLUEBERRIES —Japanese Beetle	1½ Pts per acre	1 day
Apply when first berries turn blue and repeat at 10 day intervals as long as necessary.		
BLACKBERRIES, BOYSENBERRIES, LEWBERRIES, LOGANBERRIES, RASPBERRIES —Mites, Thrips, Leafhoppers, Japanese Beetle	1½ Pts.	1 day***
Aphids, Rosa Scale	3 Pts. per acre	1 day***
CRANBERRIES —Leafhoppers, Blackheaded Fireworm, Spittlebug Nymphs, Cranberry Fruitworm	1½ Pts.	3 days***
GRAPES —Leafhoppers, Spider Mites, Mealybugs	1½ Pts.	3 days***
Injury may occur on Ribier, Italia, Cardinal & Almeria varieties if applied after the cluster stage. ***Apply as needed and repeat as necessary.		



50

Use higher rates for taller alfalfa and clover.

BARLEY, OATS, RYE, WHEAT (Grain forms) 1 1/2-2 Pts. 7 days

English Grain Aphid, Greenbugs, Cereal Leaf Beetle
Apply as needed and use in sufficient water for good coverage

Armyworms 2 Pts. 7 days

PEPPERMINT, SPEARMINT--Aphids, Spider 1 1/2 Pts. 7 days

Mites, Leafhoppers
Use in sufficient water for good coverage.

RICE (Grain form) 2 1/2 Pts. 7 days

Apply soon after first rice plants appear on surface of the water.

Rice Stink Bug 1-1 1/2 Pts. 7 days

Apply by airplane during early milk or dough stage and repeat as necessary. Mix with approximately 5 to 10 gallons per acre. Use tent with stop growth of insects.

SUGAR BEETS--Aphids, Spider Mites 1 1/2-2 Pts.

Apply as needed. Repeat if necessary. Do not use within 7 days of harvest if tops are to be used for feed.

TOBACCO Aphids, Flea 1-1 1/2 Pts. 7 days**

Plant Beds, 2 pts. per 100 gals. of water but not within 3 days of transplanting. Apply at 1/2 rate for 1/2 acre. 1 1/2 times 3 spray per acre.

ASPARAGUS Thrips 1 1/2 Pts. 1 day

Apply on foliage as needed.

BEANS Mexican Bean Beetle, Leafhoppers, Spider Mites, Japanese Beetle 1 1/2 Pts. 1 day**

BEEFS Aphids 1 1/2 Pts. 7 days**

BRUSSELS SPROUTS, CALLIFLOWER, CABBAGE, BROCCOLI, KALE, KOHLRABI, TURNIP, MUSTARD GREENS, CARROTS 1 1/2-2 Pts. 7 days**

Aphids, Imported Cabbage Worm, Cabbage

MALATHION

CONTAINS 5 LBS. MALATHION PER GALLON

ACTIVE INGREDIENTS:
 Malathion (0,0-dimethyl dithiophosphate)
 diethyl mercaptosuccinate
 Heavy Aromatic Napthalene

INERT INGREDIENTS:
 Total

CAUTION: Keep Out of Reach of Children
 See side of package for directions.

CAUTION
 HARMFUL IF SWALLOWED. AVOID PROLONGED BREATHING OF FOG OR SPRAY MIST. AVOID PROLONGED OR REPEATED CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF FOOD AND FEEDSTUFFS. KEEP AWAY FROM CHILDREN AND DOMESTIC ANIMALS. STORE ABOVE 32° F.



MALATHION

CONTAINS 5 LBS. MALATHION PER GALLON

ACTIVE INGREDIENTS:

Malathion (Oxidimethyl diethyl phosphorothioate) diethyl mercaptosuccinate	55.2%
Heavy Aromatic Naphtha	35.5%
INERT INGREDIENTS	9.3%
Total	100.0%

CAUTION: Keep Out of Reach of Children
 See side panels for directions and cautions

CAUTION

IF SWALLOWED, AVOID PROLONGED BREATHING VAPOR OR SPRAY MIST. AVOID PROLONGED CONTACT WITH SKIN. WASH THOROUGHLY AFTER USE. AVOID CONTAMINATION OF

Wings, Nymphs

PELANS —Spider Mites, Aphids	1-2 Pts	0 days**
GRAPEFRUIT, LEMONS, LIMES, ORANGES, TANGERINES, TANGELUS, KUMQUATS California Red scale, Yellow Scale, Purple scale, Black scale (Single Brooded, 2nd brood), Citricola scale	1-2 Pts	7 days***
Florida Red & Purple scales (light and medium infestations),	2 Pts	7 days***
Nymphs	2 1/2 Pts per 250 gals per acre	7 days**

Do not apply when trees are in bloom

BLUEBERRIES —Japanese Beetle	1 1/2 Pts per acre	1 day
Apply when first berries turn blue and repeat at 10 day intervals as long as necessary		

BLACKBERRIES, BOYSENBERRIES, CRAWBERRIES, LOGANBERRIES, RASPBERRIES —Mites, Thrips, Leafhopper, Japanese Beetle	1 1/2 Pts	1 day**
Scale, Rose scale	1 Pts per acre	1 day***

CRAWBERRIES —Leafhoppers, Black-headed fireworm, Spittlebug, Thrips, Strawberry rootworm	1 1/2 Pts	3 days**
---	-----------	----------

GRAPES —Leafhoppers, Spider Mites, and young	2 1/2 Pts	3 days**
---	-----------	----------

Injury may occur in fruit if applied during bloom
 Cautions apply to all crops and insects
 **Apply when insects are present

CROPS AND INSECTS	Rate per 100 Gals of Water	INTERVAL (last spray to harvest)
ORNAMENTALS —Oystershell Scale, Leafhopper, European Spruce Sawfly, Magnolia Scale	2 1/2 Pts. or less	0 days

CABBAGE, BROCCOLI, KALE, KOHLRABI, TURNIP, MUSTARD GREENS, CARROTS
 Aphids, Imported Cabbage Worm, Cabbage Looper, Leafhoppers
 1 1/2-2 Pts 7 days***

LETTUCE, SPINACH—Aphid Mites
 Cabbage Looper, Leafhoppers
 Leaf Lettuce—do not apply within 14 days of harvest
 2 Pts 7 days***

ONIONS, GARLIC—Thrips
 Onion Maggot
 1/2 Pt 7 days***

PEAS—Pea Aphid, Pea Weevil
 Apply in early bloom period before eggs are laid
 Repeat as necessary
 1 1/2-2 Pts 7 days***

POTATOES—Aphid, Colorado Potato Beetle
 RABBIES, PRUNING KNIVES
 MOSBRAD, H. BUTANERS—BORERS
 TOMATOES—Spider Mites, Aphids,
 Tomato Russet, Flea, Striped
 STRAWBERRIES—Aphid, Flea Beetle
 Lygus Bugs, Spittlebug
 Potato Leafhopper, Strawberry
 Whiteflies
 Strawberry Root Weevil
 Apply in early bloom period before eggs are laid
 1 1/2-2 Pts 7 days***

APPLES, PEARS, CHERRIES
 LIME & SULFUR
 Cover trees completely. Malathion
 injury to Monarch and Gortala
 is possible in wetting

HARMFUL IF SWALLOWED. AVOID PROLONGED BREATHING VAPOR OR SPRAY MIST. AVOID PROLONGED OR REPEATED CONTACT WITH SKIN. WASH THOROUGHLY AFTER USING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. KEEP AWAY FROM CHILDREN AND DOMESTIC ANIMALS. STORE ABOVE 32° F.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Apply this product only as specified on label.
 Empty containers and contents and dispose of wastes by burning in non-crop lands away from water. Containers should be disposed of by punching holes in them and burying with wastes or by incineration (not of waste).
 Malathion is highly toxic to bees exposed to direct treatment or residues on crops. Protective measures may be obtained from your cooperative Agricultural Extension Service.
 Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in growing areas while food is expected.

ANTIDOTE

ANTIDOTE IF SWALLOWED: Give a tablespoon of salt in glass of warm water and repeat until vomit ceases. Have victim lie down and keep quiet.
CALL A PHYSICIAN IMMEDIATELY!
NOTE TO PHYSICIANS: Malathion upon repeated or prolonged use may cause cholinesterase inhibition. Atropine is antidotal.

NOTE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application are not stated, Seller makes no claims other than those stated on this label.

NET CONTENTS

GALS

MANUFACTURED BY

PBI - GORDON CORPORATION

ST. LOUIS, MISSOURI