

DROON'S NET CONTENTS CHE GALLON

INSTITUTIONAL AREA SPRAY

ACTIVE INGREDIENTS:

Petroleum Distillate	98.92%
*Technical Piperonyl Butoxide	.90%
Pyrethrins	.18%
Total	100.00%

*Equivalent to 0.72% (butyl carbityl) (6-propyl piperonyl) ether and 0.18% related compounds.

CAUTION: Keep Out of Reach of Children
See Side Panels for Additional Cautions

NET CONTENTS CAL (S)

988-371

MANUFACTURED BY

GORDON CORPORATION

KANSAS GITTY, KANSAS

DIRECTIONS FOR USE

STORED PRODUCTS IN ELEVATORS, GRANARIES, WAREHOUSES AND MILLS:

For control of the exposed stages of Field Crickets, Granary, Weevils, Confused Flour Beetles, Grain Moths, Cadelles, Rust Red Flour Beetles, Saw Toth Grain Beetles, Spider Beetles, Grain Mites, Rice Weevils and Cheese Skipper Flies: After thorough Cleanup and prior to filling with grain apply at the rate of I gallon for each 1800 square feet of area. Wet surfaces of the storage sarea thoroughly. Hit insects directly with the spray whenever possible. Repeat application as often as necessary. Do not apply directly to grain or oil-bearing seeds.

CAUTION

CAUTION: Harmful if swallowed. Keep out of reach of children and domestic animals. Do not spray into or near open flame. Avoid spraying directly on feed, foodstuffs and food equipment. Keep container tightly sealed when not in use

Before using this product, exposed food products and packaging materials must be removed from the room or protected. After using this compound, equipment and utensils must be thoroughly rinsed with potable water before operations are resumed.

NOTICE—Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 2217-523

988-371

DIRECTIONS FOR USE

Institutional Area Spray is a "Ready to Use" insecticide to be used in appropriate spray equipment (fog applicators, sprayers, vaporizers or other dispensers) according to the following dispetiment. lowing directions.

When using in a confined area, vacate treated area and ventilate when treatment is completed.

FOOD PROCESSING PLANTS AND INSTITUTIONS:

Food should be removed by covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When ising the product as a space spray in these areas apply only when plant is not in operation.

Roaches, Waterbugs, Anti, Silverfish and Spiders: Apply as a scoarse spray to all cracks and crevices where these timesets mormally hide. Spraying of dark places such as behind pipes, beneath sinks and along funways infested with roaches, waterbugs, ants, silverfish and spiders recommended. Hit these insects with spray whenever possible. Repeat application as often as required.

Bedbugs: Spray bed frames, springs, and all sides of mattresses, especially seams and tufts. Spray behind baseboards, loose wall paper and into all cracks around the room where this in-sect may hide and breed. Repeat application when necessary.

Clothes Moths and Carpet Beetles: Use as a fine spray. Apply lightly to clothing or interior surfaces of storage container such as closet or trunks. When spraying clothing for clothes moth control and applications are all moth control pay particular attention to all seams and folds of garments. Repeat treatment every 30 days unless articles are stored in moth-tight containers. For Carpet beetles in food processing plants treat same as roaches.

Flies, Mosquitoes and Gnats: Use as a mist spray. Close room and discharge spray throughout room. Use 1 ounce of spray for each 1000 cubic feet of space. Leave room closed for at least 10 minutes. Sweep up fallen insects. Repeat application when necessary. Avoid direct contact with food or food utensils.

988-371

ACCEPTED

MAR 2 4 1971

FUNGICIDE AND RODE TICIDE ACT
FOR E.U... 15 POISON TEGESTER.
TO ATTACHED COMMENTS.

.1