



disease. Elm bark beetle can be controlled with sprays of Methoxychlor Tree Spray but more than spraying is required. The first and most important phase of Dutch Elm disease control is destruction of all dead or diseased wood in which elm bark beetles might develop.

Elm Bark Beetle Control:

To control Elm Bark beetles, make a dormant spray application in the spring before any leaves appear. Do not spray unless temperature is 40° F or warmer.

Hydraulic Sprayers:

Mix 8 gallons Methoxychlor Tree Spray with 92 gallons of water and apply 20 to 30 gallons to each 50 foot elm tree. Be sure all bark on trunk, limbs and twigs is thoroughly wet

ACCEPTED

10-11-74

METHOXYCHLOR

TREE SPRAY

For leafhopper control use 1 -- 2 quarts of Methoxychlor Tree Spray in 100 gallons of spray (2 to 4 tablespoons per gallon). Apply full coverage spray. Repeat spray as needed.

For Elm leaf beetle control or for the control of cankerworms, use 2 to 3 quarts per 100 gallons of spray (4 to 6 tablespoons per gallon). Apply full coverage spray.

For best results add a spreader-sticker at the rate of 2 to 4 ounces per 100 gallons. Repeat spray as needed.

WARNING: This product will tend to

METHOXYCHLOR

TREE SPRAY

Use on Flowers:

Mix 5 pints per 10 gallons of water and apply 2 to 3 gallons per 50-foot tree wetting all bark surfaces thoroughly.

ATTENTION: Harmful if swallowed. Avoid inhalation and contact with eyes. Wash with soap and water.

Keep product in tight container. Keep out of gutters, streams, and ponds. Do not spray where children play. Do not spray when weather conditions favor drift. Do not spray. Do not contaminate water, soil, or equipment. Do not use if label is missing. Apply. Do not use if label is missing. Do not use if label is missing.

Do not use if label is missing. Do not use if label is missing. Do not use if label is missing.

Do not use if label is missing. Do not use if label is missing. Do not use if label is missing.

ACTIVE INGREDIENTS

Metnoxychlor, Technical	24.0%
Aromatic Petroleum Hydrocarbon	72.5%
INERT INGREDIENTS	3.5%
	100.0%

Consists of 1,1,1-(p-methoxyphenyl)-1,1,1-trichloroethane and its isomers and related compounds.

EFFECTIVE FOR USE ON ORNAMENTAL SHADE TREES AND SHRUBS

WHEN USED AS DIRECTED

CAUTION: Keep Out of Reach of Children
See Side Panel for Additional Cautions

NET CONTENTS ONE GALLON

above 40° F.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

Rinse containers and equipment, and dispose of wastes in non-crop lands away from water supplies. Containers should be disposed of by puncturing holes in them and burying with wastes. **DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.**

NOTICE: Seller warrants that the product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claim other than those stated on the label.