

DIRECTIONS FOR USE

Bentgrass Lawns and Turf — Apply Bentgrass Herbicide at the rate of 3 to 4 pints per acre in sufficient water for uniform coverage. If hardy weeds persist, make a second application after three weeks. Maximum kill of weeds will be obtained from spring or early fall application when weeds are actively growing. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. If application is made when soil is quite warm, use reduced rates to reduce potential injury. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Seed can be safely sown 3 to 4 weeks after application at recommended rate. Do not use on dichondra.

If dandelions persist as a special problem, individual spot treatment with a single nozzle hand sprayer is recommended. Do not overdose Bentgrass. In closely mowed Bentgrass (putting and bowling greens) apply Bentgrass Herbicide at a maximum rate of 1 fluid ounce per 1,000 square feet preferably in May, or through September. Any yellowing will disappear after about one week.

NOTE: Care should be taken to avoid overdosing Bentgrass or injury may result. Large volumes of spray water will aid in obtaining uniform coverage. If hand type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, Seller makes no claims other than those stated on this label.


840-571

GORDON'S HERBICIDES ON

BENTGRASS BROADLEAF HERBICIDE

Contains **TRIMEC** Herbicide

Kills: DANDELIONS KNOTWEED HENBIT
CHICKWEED PLANTAINS SPURGE
and Many Other Broadleaf Weeds



Contains: 2,4-D, MCPP and Diquat
ONE GALLON COVERS UP TO 2.9 ACRES
KEEP FROM FREEZING

CAUTION: Keep Out of Reach of Children

CONTENTS 1 GALLON

840-571

WEEDS CONTROLLED: Bentgrass Herbicide is especially effective against weeds such as SHEEP SORREL, KNOT WEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

Do not allow spray to get on vegetable or ornamental plants since injury may result. Some injury can be expected to turf when used on Centipede or St. Augustine grass lawns. "Spot Spray" weeds only. Clover present will be damaged if sprayed.

Bedstraw	Knotweed	Sheep Sorrel
Black Medic	Lambsquarter	Shepherd's Purse
Eurockhorn	Lespedeza	Speedwell
Eurodock	Mallow	Spurge
Chicory	Morning Glory	Wild Carrot
Chickweed	Peppergrass	Wild Garlic
Clover	Pigweed	Wild Lettuce
Candelion	Plantains	Wild Onion
Cock	Poison Ivy	Yarrow
Ground Ivy	Poison Oak	
Heal-all	Purslane	
Henbit	Ragweed	

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. If contact with eyes occurs flush with plenty of water.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solutions near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable clovers are present. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

When container is empty, wash thoroughly and destroy. Never re-use. Destroy by perforating or crushing, and burying in a safe place. Do not burn.

U.S.D.A. Reg. No. 2217-529 840-571

MANUFACTURED BY

PBI - GORDON CORPORATION

KANSAS CITY, KANSAS