



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals:

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly after handling.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

—KEEP FROM FREEZING—

Prohibitions: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides.

Pesticide Disposal: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation & Recovery Act.

Container Disposal: Triple rinse (or equivalent) and offer for recycling, or dispose of in a sanitary landfill, or by other approved State and local procedures.

OTHER USE PRECAUTIONS:

1. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
2. Do not pour spray solution near these plants.
3. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable clovers are present.
4. Use only lawn-type sprayers.
5. Coarse sprays are less likely to wind-drift.
6. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species.
7. Do not spray while grass is emerging from dormancy.
8. Do not apply to newly seeded grasses until well established.
9. Do not spray when air temperature exceeds 85° F.
10. See 1 can be safely sown 3 to 4 weeks after application.
11. Do not contaminate domestic or irrigation waters.
12. Some injury can be expected to turf when used on Centipede or St. Augustine grass lawns. "Spot spray" weeds only on these types of grasses. Clover present will be damaged if sprayed.
13. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
14. Failure to observe the above precautions may result in injury.

GORDON'S

PROFESSIONAL TURF PRODUCTS



TRIMEC[®]

BROADLEAF HERBICIDE

BENTGRASS FORMULA

Kills: DANDELIONS KNOTWEED HENBIT
CHICKWEED PLANTAINS SPURGE

and Many Other Broadleaf Weeds

ONE GALLON COVERS UP TO 2.9 ACRES

KEEP FROM FREEZING

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid.....	19.84%
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	6.12%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	2.53%
INERT INGREDIENTS	71.51%

Total.....100.00%

THIS PRODUCT CONTAINS:

- *1.46 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 16.40%.
- **0.45 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 5.08%.
- ***0.19 lb. 3,6-dichloro-o-anisic acid per gallon or 2.10%.

***Isomer Specific by AOAC Method 8.D01-5.

KEEP OUT OF REACH OF CHILDREN

CAUTION


Statement of Practical Treatment

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See side panels for additional precautionary statements.

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation. U.S. Patent No. 3,284,188.

NET CONTENTS ONE U.S. GALLON



Application: Apply Bentgrass Formula at the rate of 8 to 4 pints per acre in sufficient water for uniform coverage. If hardy weeds persist, make a second application after three weeks. Maximum kill of weeds will be obtained from spring or early fall application when weeds are actively growing. Avoid application on excessively dry or hot periods unless irrigation is available. If application is made when soil is quite warm, use reduced rates to reduce potential injury. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Seed can be safely sown 3 to 4 weeks after application at recommended rate. Do not use on dichondra.

If dandelions persist as a special problem, individual spot treatment with a single nozzle hand sprayer is recommended. Do not overdo Bentgrass. In closely mowed Bentgrass (putting and bowling greens) apply Bentgrass Formula at a maximum rate of 1 fluid ounce per 1,000 square feet preferably in May, or through September. Any yellowing will disappear after about one week.

For hard-to-kill summer weeds, use 6 pints per acre. Do not apply this rate to bentgrass, centipede or St. Augustine grasses.

NOTE: Care should be taken to avoid overdoing Bentgrass or injury may result. Large volumes of spray water will aid in obtaining uniform coverage. If hand type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

WEEDS CONTROLLED: Bentgrass Formula is especially effective against weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

Bentgrass	Knotweed	Regweed
Black Medick	Lambsquarter	Sheep Sorrel
Buckhorn	Loosestrife	Shepherd's Purse
Burdock	Mallow	Speedwell
Chicory	Morning Glory	Spurge
Chickweed	Peppercorn	Wild Carrot
Clover	Pigweed	Wild Garlic
Dandelion	Plantain	Wild Lettuce
Deck	Polygon	Wild Onion
Ground Ivy	Polygon	Yarrow
Heart-ail	Purslane	
Henbit		

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, Seller makes no claims other than those stated on this label.

MADE BY **pbi/gordon**
corporation
KANSAS CITY, KANSAS 64108