



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals:

Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly after handling.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

—KEEP FROM FREEZING—

Prohibitions: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not store near other pesticides.

Pesticide Disposal: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation & Recovery Act.

Container Disposal: Triple rinse (or equivalent) and offer for recycling, or dispose of in a sanitary landfill, or by other approved State and local procedures.

OTHER USE PRECAUTIONS:

1. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
2. Do not pour spray solution near these plants.
3. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable clovers are present.
4. Use only lawn-type sprayers.
5. Coarse sprays are less likely to wind-drift.
6. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species.
7. Do not spray while grass is emerging from dormancy.
8. Do not apply to newly seeded grasses until well established.
9. Do not spray when air temperature exceeds 85° F.
10. Seed can be safely sown 3 to 4 weeks after application.
11. Do not contaminate domestic or irrigation waters.
12. Some injury can be expected to turf when used on Centipede or St. Augustine grass lawns. "Spot spray" weeds only on these types of grasses. Clover present will be damaged if sprayed.
13. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides.
14. Failure to observe the above precautions may result in injury.

BEST AVAILABLE COPY
GORDON'S
PROFESSIONAL TURF PRODUCTS

TRIMEC®

BROADLEAF HERBICIDE

BENTGRASS FORMULA

Kills: DANDELIONS KNOTWEED HENBIT
CHICKWEED PLANTAINS SPURGE

and Many Other Broadleaf Weeds

ONE GALLON COVERS UP TO 2.9 ACRES

KEEP FROM FREEZING

ACTIVE INGREDIENTS:

2217-529
*Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid..... 19.84%
*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid..... 6.12%
*Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)..... 2.53%
INERT INGREDIENTS..... 71.51%

Total..... 100.00%

THIS PRODUCT CONTAINS:

*1.46 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid per gallon or 16.40%.

**0.45 lb. 2,4-Dichlorophenoxyacetic acid per gallon or 5.08%.

***0.19 lb. 3,6-dichloro-o-anisic acid per gallon or 2.10%.

**Isomer Specific by AOAC Method 6.D01-5.

KEEP OUT OF REACH OF CHILDREN
CAUTION


Statement of Practical Treatment

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See side panels for additional precautionary statements.

TRIMEC® is a registered trademark of PBI/Gordon Corporation U.S. Patent No. 3,284,186

NET CONTENTS ONE U.S. GALLON



Bentgrass Formula Apply Bentgrass Formula at the rate of 6 to 4 pints per acre in sufficient water for uniform coverage. If hardy weeds persist, make a second application after three weeks. Maximum kill of weeds will be obtained from spring or early fall application when weeds are actively growing. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. If application is made when soil is quite warm, use reduced rates to reduce potential injury. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Seed can be safely sown 3 to 4 weeks after application at recommended rate. Do not use on dichondra.

If dandelions persist as a special problem, individual spot treatment with a single nozzle hand sprayer is recommended. Do not overdo Bentgrass. In closely mowed Bentgrass (putting and bowling greens) apply Bentgrass Formula at a maximum rate of 1 fluid ounce per 1,000 square feet preferably in May, or through September. Any yellowing will disappear after about one week.

For hard-to-kill summer weeds, use 6 pints per acre. Do not apply this rate to bentgrass, centipede or St. Augustine grasses.

NOTE: Care should be taken to avoid overdosing Bentgrass or injury may result. Large volumes of spray water will aid in obtaining uniform coverage. If hand type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent turf discoloration or injury.

WEEDS CONTROLLED: Bentgrass Formula is especially effective against weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS and HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

Bentgrass	Knotweed	Ragweed
Black Medic	Lambsquarters	Sheep Sorrel
Buckhorn	Loosestrife	Shepherd's
Burdock	Mallow	Purse
Chicory	Morning Glory	Speedwell
Chickweed	Peppercorn	Spurge
Clover	Pigweed	Wild Carrot
Dandelion	Plantain	Wild Garlic
Deck	Poison Ivy	Wild Lettuce
Ground Ivy	Poison Oak	Wild Onion
Heart-ail	Parsley	Yarrow
Henbit		

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, Seller makes no claims other than those stated on this label.


MADE BY **G** pbi / gordon
CORPORATION
KANSAS CITY, KANSAS 64116

The Certification of Compliance for the 2,4-D Exposure Reduction Program


"I, being an authorized representative of PBI/Gordon Corporation, certify that all containers of **Trimec® Broadleaf Herbicide Bentgrass Formula (EPA No. ~~2217-529~~)** produced by June 15, 1994 for products registered for use only on residential or turf sites, excluding sod farms will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all container of said product sold or distributed by this company by January 1, 1995 for products registered for use only on residential or turf sites, excluding sod farms will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products."



James A. Armbruster, Ph.D.
Mgr. Regulatory Services



Date



TRIMEC® BROADLEAF HERBICIDE

BENTGRASS FORMULA

2.5 GALLONS COVERS UP TO 7.25 ACRES

CONTROLS: Dandelions, Chickweed, Knotweed, Plantains, Henbit, Spurge,
and many other broadleaf weeds.

- KEEP FROM FREEZING -

ACTIVE INGREDIENTS

* Dimethylamine salt of 2-(2-Methyl-4-chlorophenoxy) propionic acid	19.84%
** Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	6.12%
*** Dimethylamine salt of Dicamba (3,6-Dichloro-o-anisic acid)	2.53%
INERT INGREDIENTS	<u>71.51%</u>
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- *1.46 lbs. 2-(2-Methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 16.40%.
 - **0.45 lbs. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 5.08%
 - ***0.19 lbs. 3,6-Dichloro-o-anisic acid equivalent per gallon or 2.10%
- Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

KEEP OUT OF REACH OF CHILDREN

CAUTION


Statement of Practical Treatment

In case of contact, wash skin with plenty of soap and water;
For eyes, flush with water for 15 minutes and get medical attention.
See side panel for additional Precautionary Statements.

NET CONTENTS 2 1/2 U.S. GALLONS (9.46 LITERS)

840/1262 APXXXXXX
EPA REG. NO. 2217-529
EPA EST. N O. 2217-KS-1

Mfd. by PBI/GORDON CORPORATION
KANSAS CITY, MISSOURI 64101



3 7 5
STOP!

READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS
AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wear long pants, long-sleeved shirt, shoes and gloves when mixing, loading or applying this product.

When mixing, loading, or applying this product, wear long-sleeved shirt, long pants, socks, shoes and chemical resistant gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Containers over 1 gallon and less than 5 gallons: Persons engaged in open purging of this product must also wear coveralls or a chemical resistant apron.

Containers of 5 gallons or more: A mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Environmental Hazards: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.


Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

FOR USE ON RESIDENTIAL AND OTHER TURF SITES EXCLUDING SOD FARMS.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application.
Do not enter treatment areas until spray has dried or dust has settled.



4 2 5

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

USE PRECAUTIONS: Do not apply this product through any type of irrigation system. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solution near these plants. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable clovers are present. Use only lawn-type sprayers. Coarse sprays are less likely to wind-drift. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within the dripline of tree and other ornamental species. Do not spray while grass is emerging from dormancy. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 85° F. Seed can be sown 3 to 4 weeks after herbicide application. Do not contaminate domestic or irrigation waters. Some injury can be expected to turf when used on centipedegrass or St. Augustinegrass lawns; "spot spray" weeds only on these types of grasses. Clover present will be damaged if sprayed. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury.

Bentgrass Lawns and Turf—Apply Bentgrass Formula at the rate of 3 to 4 pints of product per acre in sufficient water for uniform coverage. If hardy weeds persist, make a second application after three weeks. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. If application is made when soil is quite warm, use reduced rates to reduce potential injury. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until the second or third mowing. Seed can be sown 3 to 4 weeks after herbicide application. Do not use on dichondra.

If dandelions persist as a special problem, individual spot treatment with a single nozzle hand sprayer is recommended. Do not overdose bentgrass. In closely mowed bentgrass (putting and bowling greens) apply Bentgrass Formula at a maximum rate of 1 fluid ounce per 1,000 square feet preferably in May, or through September. Any yellowing will disappear after about one week.

For hard-to-kill summer weeds, use 6 pints of product per acre. Do not apply this rate to bentgrass, centipedegrass or St. Augustinegrass.

The maximum application rate to turf is 0.4 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast application per treatment site is 2 per year.

NOTE: Care should be taken to avoid overdosing bentgrass or injury may result. Large volumes of spray water will aid in obtaining uniform coverage. If hand type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom. Sweeping movements with a multiple nozzle boom will result in heavy over-application and subsequent turf discoloration or injury.

5 of 5
WEEDS CONTROLLED: Bentgrass Formula is especially effective against weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. Broadleaf weeds are listed below.

Bedstraw
Black Medic
Buckhorn
Burdock
Chicory
Chickweed
Clover
Dandelion
Dock
Ground Ivy
Healall

Henbit
Knotweed
Lambsquarter
Lespedeza
Mallow
Morningglory
Peppergrass
Pigweed
Plantains
Poison Ivy
Poison Oak

Purslane
Ragweed
Sheep Sorrel
Shepherdspurse
Speedwell
Spurge
Wild Carrot
Wild Garlic
Wild Lettuce
Wild Onion
Yarrow

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

