

**DIRECTIONS FOR USE**

**ORNAMENTAL LAWNS AND TURF (Other than Bentgrass):** Apply COMMERCIAL HERBICIDE at the rate of 2 to 3 pints per acre in sufficient water for adequate coverage. Use the higher rate in heavy weed growth or on older weeds or during periods of poor growth. Maximum kill of weeds will be obtained from spring or early fall application when weeds are actively growing. Spray when average temperatures are 50° F. or higher. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Grass seed can be safely sown 3 to 4 weeks after application at recommended rate. Do not use on dichondra, carpetgrass, and St. Augustine grass. Do not spray under drip line of ornamentals.

**BENTGRASS TURF:** In closely mowed Bentgrass (putting and bowling greens) apply COMMERCIAL HERBICIDE at a maximum rate of 1/2 fluid ounce per 1,000 square feet preferably in May, or mid-August through September. Slight turf yellowing will disappear after about one week.

**NOTE:** Care should be taken to avoid overdosing Bentgrass or injury may result. Large volumes of spray water (i.e., 1 gallon per 250 or 300 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local overapplication and subsequent turf discoloration or injury.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary. Seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 2217-530 839-1270



# COMMERCIAL BROADLEAF HERBICIDE

Contains \*TRIMEC® Herbicide

**Kills:** DANDELIONS KNOTWEED HENBIT  
CHICKWEED PLANTAINS SPURGE  
and Many Other Broadleaf Weeds

Contains: 2,4-D, MCPP and Dicamba

ONE GALLON COVERS UP TO 4 ACRES

KEEP FROM FREEZING

**ACTIVE INGREDIENTS:**

- \*Dimethylamine salt of 2,4-dichloro-phenoxyacetic acid ..... 32.07%
- \*\*Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid ..... 16.12%
- \*\*\*Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) ..... 4.04%
- Dimethylamine Salts of Related Compounds ..... 0.05%
- INERT INGREDIENTS: ..... 47.72%
- Total ..... 100.00%

- \*26.7% 2,4-dichlorophenoxyacetic acid equivalent—2.62 pounds per gallon
- \*\*13.4% 2-(2-methyl 4-chlorophenoxy) propionic acid equivalent—1.31 pounds per gallon
- \*\*\*3.36% 3,6-dichloro-o-anisic acid equivalent—0.33 pounds per gallon

**CAUTION: Keep Out of Reach of Children**

See Back Panel for Additional Cautions

\*Trimec® is a registered trademark of P.B.I.—Gordon Corporation.  
U.S. Patent No. 3,284,186

839-1270

CONTENTS 1 GALLON

MANUFACTURED BY



**GORDON CORPORATION**

KANSAS CITY, KANSAS

**WEEDS CONTROLLED:** COMMERCIAL HERBICIDE is especially effective against weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

Do not allow spray to get on vegetable or ornamental plants since injury may result. Some injury can be expected to turf when used on Centipede or St. Augustine grass lawns. "Spot Spray" weeds only. Clover present will be damaged if sprayed.

Bedstraw	Knotweed	Sheep Sorrel
Black Medic	Lambsquarter	Shepherd's Purse
Buckhorn	Lespedeza	
Burdock	Mallow	Speedwell
Chicory	Morning Glory	Spurge
Chickweed	Peppergrass	Thistle
Clover	Pigweed	Wild Carrot
Dandelion	Plantains	Wild Garlic
Dock	Poison Ivy	Wild Lettuce
Ground Ivy	Poison Oak	Wild Onion
Heal-all	Furslane	Yarrow
Henbit	Ragweed	

**CAUTION:** Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solutions near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.



839-1270