

DIRECTIONS FOR USE

ORNAMENTAL LAWNS AND TURF (Other than Bentgrass): Apply TRIMEC COMMERCIAL HERBICIDE at the rate of 2 to 3 pints per acre in sufficient water for adequate coverage. Use the higher rate in heavy weed growth or on older weeds or during periods of poor growth. Maximum kill of weeds will be obtained from spring or early fall application when weeds are actively growing. Spray when average temperatures are 50° F. or higher. Avoid applying during long, excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Grass seed can be safely sown 3 to 4 weeks after application at recommended rate. Do not use on dichondra, carpetgrass, and St. Augustine grass. Do not spray under drip line of ornamentals.

BENTGRASS TURF: In closely mowed Bentgrass (putting and bowling greens) apply TRIMEC COMMERCIAL HERBICIDE at a maximum rate of 1/2 fluid ounce per 1,000 square feet preferably in May, or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE: Care should be taken to avoid overdosing Bentgrass or injury may result. Large volumes of spray water (i.e., 1 gallon per 250 or 300 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local overapplication and subsequent turf discoloration or injury.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary. Seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 2217-530 839-1070



trimec®
COMMERCIAL
BROADLEAF HERBICIDE

Contains New Patented TRIMEC Weed Killer.
 U. S. Patent No. 3,284,186

Kills: DANDELIONS KNOTWEED HENBIT
 CHICKWEED PLANTAINS SPURGE
 and Many Other Broadleaf Weeds

Contains: 2,4-D, MCPP and Dicamba
ONE GALLON COVERS UP TO 1 ACRES
KEEP FROM FREEZING

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	32.07%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	16.12%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	4.04%
Dimethylamine Salts of Related Compounds....	0.05%
INERT INGREDIENTS:	47.72%
Total	100.00%
*26.7% 2,4-dichlorophenoxyacetic acid equivalent— 2.62 pounds per gallon	
**13.4% 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent—1.31 pounds per gallon	
***3.36% 3,6-dichloro-o-anisic acid equivalent— 0.33 pounds per gallon	

CAUTION: Keep Out of Reach of Children
 See Back Panel for Additional Cautions
 T.M. Registered PBI, K.C., K. 839-1070

CONTENTS 1 GALLON

MANUFACTURED BY

PBI - GORDON CORPORATION
 KANSAS CITY, KANSAS

ACCEPTED
 Nov. 16, 1970
 UNDER THE FEDERAL INSTITUTE
 PUNISHING AND RESENTING ACT
 FOR ECONOMIC POISON REGISTRATION
 UNDER NO. 3317-530

WEEDS CONTROLLED BY THIS COMMERCIAL HERBICIDE:
 against weeds such as:
 KNOTWEED
 WEED, OXALIS
 to these, it kills
 weeds, some of which
 Do not allow this herbicide on
 ornamental plants.
 Some injury may be
 used on Centaurea
 lawns. "Spot" applications
 present will

Bedstraw
 Black Medic
 Buckhorn
 Burdock
 Chicory
 Chickweed
 Clover
 Dandelion
 Dock
 Ground Ivy
 Heal-all
 Henbit

CAUTION: Avoid contact with skin or clothing. Avoid contact with eyes. Avoid breathing spray. Avoid drift on flowers, ornamentals, and other desirable plants. Do not pour spray on desirable plants. Do not store in areas where children can reach. Do not spray on turf when mowed. When used on lawn, spray areas are less likely to be injured and shrubs to St. Augustine.



trimec®

COMMERCIAL BROADLEAF HERBICIDE

Contains New Patented TRIMEC Weed Killer.
U. S. Patent No. 3,284,186

Kills: DANDELIONS KNOTWEED HENBIT
CHICKWEED PLANTAINS SPURGE
and Many Other Broadleaf Weeds

Contains: 2,4-D, MCPP and Dicamba
ONE GALLON COVERS UP TO 1 ACRES
KEEP FROM FREEZING

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-dichloro-phenoxycetic acid	32.07%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	16.12%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	4.04%
Dimethylamine Salts of Related Compounds....	0.05%
INERT INGREDIENTS:	47.72%
Total	100.00%

- *26.7% 2,4-dichlorophenoxyacetic acid equivalent—2.62 pounds per gallon
- **13.4% 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent—1.31 pounds per gallon
- ***3.36% 3,6-dichloro-o-anisic acid equivalent—0.33 pounds per gallon

CAUTION: Keep Out of Reach of Children

See Back Panel for Additional Cautions
T.M. Registered PBI, K.C., K. 839-1070
CONTENTS 1 GALLON

MANUFACTURED BY

PBI - GORDON CORPORATION

KANSAS CITY, KANSAS

WEEDS CONTROLLED: TRIMEC COMMERCIAL HERBICIDE is especially effective against weeds such as SHEEP SORREL, KNOTWEED, SPOTTED SPURGE, CHICKWEED, OXALIS AND HENBIT. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below. Do not allow spray to get on vegetable or ornamental plants since injury may result. Some injury can be expected to turf when used on Centipede or St. Augustine grass lawns. "Spot Spray" weeds only. Clover present will be damaged if sprayed.

Bedstraw	Knotweed	Sheep Sorrel
Black Medic	Lambsquarter	Shepherd's Purse
Buckhorn	Lespedeza	Speedwell
Burdock	Mallow	Spurge
Chicory	Morning Glory	Thistle
Chickweed	Peppergrass	Wild Carrot
Clover	Pigweed	Wild Garlic
Dandelion	Plantains	Wild Lettuce
Dock	Poison Ivy	Wild Onion
Ground Ivy	Poison Oak	Yarrow
Heal-all	Purslane	
Henbit	Ragweed	

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solutions near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.



339-1070

UNDER THE FEDERAL INDEBTEDNESS ACT
 FUNDS AND RESOURCES NOT
 FOR ECONOMIC REVENUE INVESTMENT
 UNDER NO. 237-530