



RES-Q*

SEED DISINFECTANT POWDER**

MANUFACTURED BY
PCB - GORDON CORPORATION
KANSAS CITY, KANSAS

DIRECTIONS

Mix GORDON RES-Q SEED DISINFECTANT DIRECTLY with the seed in the drill or planter box or in a regular dust treater at the rate of:

WHEAT	BARLEY, OATS, RYE	FLAX
1 oz. per bushel	2 oz. per bushel	4 oz. per bushel

Follow these steps to ensure thorough mixing with the seed:

When starting with an empty drill or planter box enough seed to cover the bottom of the box is to be treated in a separate container and placed in the bottom of the empty drillbox. This seed can be treated in an empty carton or drum which can later be destroyed. This treatment of seed in a separate container is not necessary when, at refilling, enough treated seed remains to cover the bottom of the drill or planter box.

1. Fill the planter box to one-half capacity. (Do not fill the planter box over one-half full).
2. Sprinkle seed disinfectant over the seed surface at the rate of 1 oz. per bushel of wheat, 2 oz. per bushel of barley, rye and oats and 4 oz. per bushel of flax.
3. Mix well by stirring with a stick or paddle until the seed is evenly coated with the powder. Do not mix with bare hands.
4. Add the remaining half to fill the seed box and sprinkle the seed disinfectant over the seed surface at the above recommended rates, per bushel, and mix as directed in step 3 above.

NOTE: Not effective against Loose Smut of Wheat and True Loose Smut of Barley.

U.S.D.A. Reg. No. 2217-534

513-271



RES-Q*

SEED DISINFECTANT POWDER**

PBI

MANUFACTURED BY
GORDON CORPORATION

KANSAS CITY, KANSAS

CAUTION: Caution should be observed in using. Do not mix seed with the bare hands.

1. Avoid inhaling plus wear a dry dust filter and fume mask suitable for protection from Maneb, Captan and Hexachlorocyclohexene when handling this seed disinfectant.
2. Avoid contact with eyes, skin and clothing. In case of contact with skin, wash with soap and water. For eyes, flush with water for 15 minutes and get medical attention. May cause irritation of eyes, nose and throat.
3. Wash hands before eating or smoking.
4. Do not allow to become wet during storage. If Maneb becomes wet during storage, its fungicidal effectiveness may be reduced and it may also produce vapors which may be flammable.
5. Keep container tightly closed except when removing contents and store in a dry, well ventilated area away from food.
6. Treat only the amount of seed to be sown to avoid the possibility of storing treated seed and the associated hazards of food contamination.
7. Do not use treated or left over seed for food or feed purposes.
8. This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, disposal of wastes.

NOTICE: Seller warrants that this product conforms to the present statement on the label. Since conditions of use, storage, weather, compatibility with other chemicals and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

PBI

REGISTERED
May 25, 1971
221534

513-271



RES-Q*

SEED DISINFECTANT POWDER**

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

DIRECTIONS

Mix GORDON RES-Q SEED DISINFECTANT DIRECTLY with the seed in the drill or planter box or in a regular dust treater at the rate of:

WHEAT	BARLEY, OATS, RYE	FLAX
1 oz. per bushel	2 oz. per bushel	4 oz. per bushel

Follow these steps to ensure thorough mixing with the seed:

When starting with an empty drill or planter box enough seed to cover the bottom of the box is to be treated in a separate container and placed in the bottom of the empty drillbox. This seed can be treated in an empty carton or drum which can later be destroyed. This treatment of seed in a separate container is not necessary when, at refilling, enough treated seed remains to cover the bottom of the drill or planter box.

1. Fill the planter box to one-half capacity. (Do not fill the planter box over one-half full).
2. Sprinkle seed disinfectant over the seed surface at the rate of 1 oz. per bushel of wheat, 2 oz. per bushel of barley, rye and oats and 4 oz. per bushel of flax.
3. Mix well by stirring with a stick or paddle until the seed is evenly coated with the powder. Do not mix with bare hands.
4. Add the remaining half to fill the seed box and sprinkle the seed disinfectant over the seed surface at the above recommended rates, per bushel, and mix as directed in step 3 above.

NOTE: Not effective against Loose Smut of Wheat and True Loose Smut of Barley.

U.S.D.A. Reg. No. 2217-534

513-271



RES-Q*

SEED DISINFECTANT POWDER**

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

CAUTION: Caution should be observed in using. Do not mix in seed with the bare hands.

1. Avoid inhaling plus wear a dry dust filter and fume mask suitable for protection from Maneb, Captan and Hexachlorobenzene when handling this seed disinfectant.
2. Avoid contact with eyes, skin and clothing. In case of contact with skin, wash with soap and water. For eyes, flush with water for 15 minutes and get medical attention. May cause irritation of eyes, nose and throat.
3. Wash hands before eating or smoking.
4. Do not allow to become wet during storage. If Maneb becomes wet during storage, its fungicidal effectiveness may be reduced and it may also produce vapors which may be flammable.
5. Keep container tightly closed except when removing contents and store in a dry, well ventilated area away from food stuffs.
6. Treat only the amount of seed to be sown to avoid the problem of storing treated seed and the associated hazards of feed and food contamination.
7. Do not use treated or left over seed for food or feed purposes.
8. This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.



513-2

513-271