

# RES-Q\*

SEED DISINFECTANT POWDER\*\*

Contains Hexachlorobenzene, Captan and Maneb

Controls Stinking Smut or Bunt of Wheat, False Loose and Covered Smut of Barley, Oat Smuts and Many Seed or Soil-Borne Seedling Blights (Seed Borne Root Rots) of Wheat, Barley, Oats, Rye and Flax.

GORDON RES-Q Seed Disinfectant is for use directly in the drill or planter box at time of seeding, or may be used in regular dust treaters.

ACTIVE INGREDIENTS:

Hexachlorobenzene .....	20%
***Captan .....	20%
Maneb .....	15%
INERT INGREDIENTS .....	45%
TOTAL .....	100%

ACCEPTED

MAR 31 1983

U.S. Federal Insecticide,  
Fungicide and Rodenticide Act,  
the pesticide

2217-634

KEEP OUT OF REACH OF CHILDREN ←

CAUTION ←

See side panels for additional  
precautionary statements.

12 pt. bold.

18 pt. bold.

NET WEIGHT THREE POUNDS

\*Trademark Pending.

\*\*Patent Pending.

\*\*\*N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide.

STOP! READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND FOLLOW  
DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

-Hazards to Humans & Domestic Animals-

CAUTION: May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. When using, wear protective clothing such as long sleeves, long pants, hat, gloves, covering for face and eyes. Wear a dry dust filter and fume mask when handling. Wash thoroughly after use and launder clothing before reuse. Wash hands before eating or smoking.

Do not use treated or left over seed for food or feed purposes. Do not mix seed with bare hands.

-Statement of Practical Treatment-

In case of contact with skin, wash with soap and water. For eyes, flush with water for 15 minutes and get medical attention.

-Environmental Hazards-

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

-Physical or Chemical Hazards-

Do not allow this product to become wet during storage as such may produce vapors which may be flammable.

USE PRECAUTIONS:

Treat only the amount of seed to be sown to avoid the problem of storing treated seed and the associated hazards of feed and food contamination. Dispose of excess treated seeds by burial.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL ← *12 pt. bold.*

Storage:

Do not allow to become wet during storage. If Maneb becomes wet during storage, its fungicidal effectiveness may be reduced and such may also produce vapors which may be flammable. Keep container tightly closed except when removing contents. Store in a dry, well-ventilated area away from foodstuffs.

Pesticide Disposal:

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved State procedures under Subtitle C of the Resource Conservation and Recovery Act.

Container Disposal:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

513/383

EPA REG. NO. 2217-534  
EPA EST. NO. 2217-KS-1

Mfd. By  
PBI/GORDON CORPORATION  
Kansas City, Kansas 66118

Limited Warranty. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

Exclusion of Other Warranties and Remedies. Except where such disclaimers and exclusions are specifically prohibited by applicable law, THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESS OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty, BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

Mix GORDON RES-Q SEED DISINFECTANT DIRECTLY with the seed in the drill or planter box or in a regular dust treater at the rate of:

WHEAT	BARLEY, OATS, RYE	FLAX
1 oz. per bushel	2 oz. per bushel	4 oz. per bushel

Follow these steps to ensure thorough mixing with the seed:

When starting with an empty drill or planter box enough seed to cover the bottom of the box is to be treated in a separate container and placed in the bottom of the empty drillbox. This seed can be treated in an empty carton or drum which can later be destroyed. This treatment of seed in a separate container is not necessary when, at refilling, enough treated seed remains to cover the bottom of the drill or planter box.

1. Fill the planter box to one-half capacity. (Do not fill the planter box over one-half full).
2. Sprinkle seed disinfectant over the seed surface at the rate of 1 oz. per bushel of wheat, 2 oz. per bushel of barley, rye and oats and 4 oz. per bushel of flax.
3. Mix well by stirring with a stick or paddle until the seed is evenly coated with the powder. Do not mix with bare hands.
4. Add the remaining half to fill the seed box and sprinkle the seed disinfectant over the seed surface at the above recommended rates, per bushel, and mix as directed in step 3 above

NOTE: Not effective against Loose Smut of Wheat and True Loose Smut of Barley.