ACCEPTED 1-5-71



BL 10,000 LAWN WEED KILLER

Contains *TRIMEC® Herbicide

KEEP FROM FREEZING One Quart Covers 10880 Sq. Ft. Contains: 2,4-D, MCPP and Dicamba

Kills: DANDELION, KNOTWEED, CLOVER, HENBIT, PLAN-TAINS, SPURGE, POISON IVY and most other broadleaf weeds.

ACTIVE INGREDIENTS: "Dimethylamine salt of 2,4-dichlerophenezyacetic acid 12.78%
"Dimethylamine salt of 2,6-dichlerophenezyacetic acid 5.82%
"Dimethylamine salt of 2,6-dichlero-anisic acid 1.51%
IMERT INGREDIENTS 29.83% Tetal 100.00%

Equivalent to:

10.5295 2.4-D Acid or 0.94 lbs. acid per gallon

14.7559 MCPP Acid or 0.42 lbs. acid per gallon

15.1 1056 Dicamba Acid or 0.027 lbs. acid per gallon

CAUTION: Keep Out of Reach of Children See Back Panel for Additional Cautions

CONTENTS 1 QUART

879-1170

U.S.D.A. Reg. No. 2217-536 *Trimec® is a registered trademark of P.B.I.—Gordon Corporation. U.S. Patent No. 3,284,185



UNDER THE FEDERAL INSECTICIDED ONTROLS
FUNDICIDE AND ROUTATICHE ACT
FOR ECONOMIC POISON REGISTER.
BEDSTRAW
UNDER NO. 230 SEACH MEDIC BUCKHORN CHICORY CHICKWEED CLOVER DANDELIGN DOCK GROUND IVY

HEAL-ALL HENBIT KNOTWEED LAMBSQUARTER

LESPEDEZA MALLOW **MORNING GLORY PEPPERGRASS** PIGWEED **PLANTAINS**

POISON IVY POISON CAK **PURSLANE**

RAGWEED SHEPHERD'S PURSE

SPEEDWELL SPURGE WILD CARROT WILD LETTUCE WILD ONION

and many others

YARROW

DIRECTIONS FOR USE

WHERE TO USE: BL 10,000 Lawn Weed Killer may be used on Bluegrass, Bermuda, or Zoysig lawns, vacant lots, playgrounds and other similar areas where hard-to-kill weeds are present. This product is especially effective on all types of hard-to-kill weeds such as HENBIT, CHICKWEED, KNOTWEED, SPOTTED SPURGE, OXALIS, BLACK MEDIC, and SHEEP SORREL. in addition to these, it controls most other broadlest lawn weeds, including Dandellon and Plantain, and a partial list is given on the side panel. Do not allow spray to get on vegetables or ornamental plants since injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot-Spray" weeds only. Clover present will be damaged if sprayed.

when To Use: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot-Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results mow before treating. Do not water lawn within 24 hours after treatment.

40W MUCH TO USE-ENTIRE LAWN

HOSE-END SPRAYER: Use a hose-end sprayer designed for lawns, Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated. For example—to spray 1,700 sq. teet, measure 5 ozs. (10 thsp.) in jar and add water to the 10 gallons level and mix Turn on hose and spray as evenly as possible over 1,700 sq. feet.

ed. teer			
Col. 1 Amt. BL 10,000 to Use: (Tbsp.) (Fl. Oz.)		Col. 2 Filf Bottle To: (Galfons)	Gol. 3 Area to Treat: (Sq. Ft.)

PRESSURE SPRAYERS: Measure indicated amount of chemical into tank and dilete with water according to the following chart. Use a spray pattern which gives small spray droplets without any fine mist and cover the area indicated:

Amt. BL 10,000 to Use: (Tbsg.) (F1, Oz.)		Amt. of Water in Sprayer: (Gallons)	Area to be Sprayed (Sq. Ft.)
(Tbsp.)	(81. 02.)	(Galiona)	(3q. rt.)
2	1	1/2	335
4	2	1	670
8	4	2	1300
12	6	3	2000

NOTE: Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

NOW MUCH TO USE: SPOT SPRAY
To spray small areas or individual weeds, do not use hose-end sprayers. Mix 1 or. (2 thsp.) in ½ gal. water in pressure sprayer and spray as required being certain to "wet" each weed. The GORDON Aerosoi Spot Y/ceder is ideal for this use. Be sure to use GORDON'S Spreader Sticker according to the directions on that container so that each weed is thoroughly wet.

cautions are desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable bentgrass or clovers are present. When container is empty, wash thoroughly and destroy. Never re-use. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin, and clothing. In case of contact, flush with water, Avoid contamination of feed or foodstuffs.

NOTICE. Selter warrants that this product conforms to the ingredient statement on the labet. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Selter makes no claims other than those stated on this label.

879-1170