

BL 10,000 LAWN WEED KILLER

Contains *TRIMEC® Herbicide

KEEP FROM FREEZING One Quart Covers 10880 Sq. Ft. Contains: 2.4-D. MCPP and Dicamba

Kills: DANDELION, KNOTWEED, CLOVER, HENBIT, PLAN-TAINS, SPURGE, POISON IVY and most other broadleaf weeds.

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid
*Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid
**Dimethylamine salt of Dicamba (3,6-dichloro-e-anislo acid)....... 12.875% 5.840% 1.341% INERT INGREDIENTS 79.944% Total.

Equivalent to:

*10.693% 2.4-D Actd or 0.93 lbs. acid per gallon

**4.826% MCPP Acid or 0.42 lbs. acid per gallon

**1.113% Dicamba Acid or 0.097 lbs. acid per gallon

Keep Out of Reach of Children CAUTION: See Back Panel for Additional Cautions

CONTENTS 1 QUART

U.S.D.A. Reg. No. 2217-536

879-971

100.000%

*Trimec® is a registered trademark of P.B.I.—Gordon Corporation. U.S. Patent No. 3,284,186



MANUFACTURED BY

GORDON CORPORATION KANSAS CITY, KANACCEPTED

11-16-71

FUNGICIDE AND RODENTICIDE ACT FOR ECO IOMIC POISON REGISTER-ED UNDER NO.

CONTROLS

BEDSTRAW

BLACK MEDIC BUCKHORN. CHICORY CHICKWEED_ CLOVER DANDELION DOCK GROUND IVY... HEAL-ALL -__ HENBIT KNOTWEED LAMBSQUARTER LESPEDEZA MALLOW MORNING GLORY **PEPPERGRASS** PIGWEED **PLANTAINS** POISON IVY POISON OAK **PURSLANE** RAGWEED **SHEPHERD'S**: < PURSE SPEEDWELL **SPURGE** WILD CARROT WILD LETTUCE WILD ONION

and many others

YARROW

WILD GARLIE

DIRECTIONS FOR USF

WHERE TO USE: BL 10,000 Lawn Weed Killer may be used on Bluegrass, Bermuda, or Zoysia lawns, vacant lots, playgrounds and other similar areas where hard-to-kill weeds are present. This product is especially effective on all types of hard-to-kill weeds such as HENBIT, CHICKWEED, KNOTWEED, SPOTTED SPOTTED SPURGE, OXALIS, BLACK MEDIC, and SHEEP SORREL. In addition to these, it controls most other broadless lawn weeds, including Dandalion and Plantain, and a partial list is given on the side panel. Do not allow spray to get on vegetables or ornamental plants since injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot-Spray" weeds only. Clover present will be damaged if sprayed.

WMEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot-Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results mow before treating. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE-ENTIRE LAWN

HOSE-END SPRAYER: Use a hose-end sprayer HOSE-END SPRAYER: Use a hose-end sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated tevel (Column 2), and spray on area indicated. For example—to spray 1,700 sq. feet, measure 5 ozs. (10 tbsp.) in jar and add water to the 10 gallons level and mix. Turn on hose and spray as evenly as possible over 1,700 sq. feet.

Col. 1		Coi. 2	Col. 3
	. 10,000 Use: (Fl. Oz.)	Fill Bottle To: (Gallons)	Area to Treat: (Sq. Ft.)
1 2	1/2	1 2	170 340
10 15	21/2 5 71/2	5 10 15	850 1700 2550
20 32	10 16	20 32	3400 5440

PRESSURE SPRAYERS: Measure indicated amount of chemical into tank and dilute with water according to the following chart. Use a spray pattern which gives small spray droplets without any fine mist and cover the area indicated:

Amt. BL 10,000 to Use:		Amt. of Water in Sprayer: (Gallons)	Area to be Sprayed
(Tbsp.)	(F1. Oz.)	(Gallons)	(Sq. Ft.)
2	1	1/2	335
4	2	1	670
8	4	2	1300
12	6	3	2000

NOTE: Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

HOW MUCH TO USE: SPOT SPRAY

To spray small areas or individual weeds, do not use hose-end sprayers. Mix 1 oz. (2 tbsp.) in ½ gal. water in pressure sprayer and spray as required being certain to "wet" each weed. The GORDON Aerosol Spot Weeder is ideal for this use.

Be sure to use GORDON'S Spreader Sticker according to the directions on that container so that each weed is thoroughly wet.

CAUTION: Avoid drift of spray mist onto vegetables, flowers, ornamental plants, strubs, trees and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable bentgrass or clovers are present. When container is empty, wash thoroughly and destroy. Never re-use, Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin, and clothing. In case of contact, flush with water, Avoid contamination of feed or foodstuffs.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary. Seller makes no claims other than those stated on this label.

879-971