



# BL 10,000 LAWN WEED KILLER

Contains \*TRIMEC® Herbicide

KEEP FROM FREEZING

One Quart Covers 10880 Sq. Ft.

Contains: 2,4-D, MCPP and Dicamba

**Kills: DANDELION, KNOTWEED, CLOVER, HENBIT, PLANTAINS, SPURGE, POISON IVY and most other broadleaf weeds.**

#### ACTIVE INGREDIENTS:

\*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid..... 12.875%

\*\*Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid..... 5.840%

\*\*\*Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)..... 1.341%

INERT INGREDIENTS..... 79.944%

Total..... 100.000%

#### Equivalent to:

\*10.693% 2,4-D Acid or 0.93 lbs. acid per gallon

\*\*4.826% MCPP Acid or 0.42 lbs. acid per gallon

\*\*\*1.113% Dicamba Acid or 0.097 lbs. acid per gallon

**CAUTION: Keep Out of Reach of Children**  
See Back Panel for Additional Cautions

**NET CONTENTS ONE QUART**

EPA REG. NO. 2217-536

EPA EST. NO. 2217-KS-1

879-475

\*Trimec® is a registered trademark of P.B.I.—Gordon Corporation. U.S. Patent No. 3,284,186

Mfd. by  **pbi/gordon**  
**corporation**  
KANSAS CITY, KANSAS 66118

ACCEPTED

APR 10 1975

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, and the pesticide  
registered under  
EPA Reg. No. 2217-536

## CONTROLS

BEDSTRAW  
BLACK MEDIC  
BUCKHORN  
CHICORY  
CHICKWEED  
CLOVER  
DANDELION  
DOCK  
GROUND IVY  
HEAL-ALL  
HENBIT  
KNOTWEED  
LAMBSQUARTER  
LESPEDeza  
MALLOW  
MORNING GLORY  
PEPPERGRASS  
PIGWEE  
PLANTAINS  
POISON IVY  
POISON OAK  
PURSLANE  
RAGWEED  
SHEPHERD'S  
PURSE  
SPEEDWELL  
SPURGE  
WILD CARROT  
WILD LETTUCE  
WILD ONION  
YARROW

and many others

#### DIRECTIONS FOR USE

**WHERE TO USE:** BL 10,000 Lawn Weed Killer may be used on Bluegrass, Bermuda, or Zoysia lawns, vacant lots, playgrounds and other similar areas where hard-to-kill weeds are present. This product is especially effective on all types of hard-to-kill weeds such as HENBIT, CHICKWEED, KNOTWEED, SPOTTED SPURGE, OXALIS, BLACK MEDIC, and SHEEP SORREL. In addition to these, it controls most other broadleaf lawn weeds, including Dandelion and Plantain, and a partial list is given on the side panel. Do not allow spray to get on vegetables or ornamental plants since injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot-Spray" weeds only. Clover present will be damaged if sprayed.

**WHEN TO USE:** Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot-Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results mow before treating. Do not water lawn within 24 hours after treatment.

#### HOW MUCH TO USE—ENTIRE LAWN

**HOSE-END SPRAYER:** Use a hose-end sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated. For example—to spray 1,700 sq. feet, measure 5 ozs. (10 tbsp.) in jar and add water to the 10 gallons level and mix. Turn on hose and spray as evenly as possible over 1,700 sq. feet.

Col. 1	Col. 2	Col. 3
Amt. BL 10,000 to Use: (Tbsp.)	Fill Bottle To: (Gallons)	Area to Treat: (Sq. Ft.)
1	1/2	170
2	1	340
5	2 1/2	850
10	5	1700
15	7 1/2	2550
20	10	3400
32	16	5440

**PRESSURE SPRAYERS:** Measure indicated amount of chemical into tank and dilute with water according to the following chart. Use a spray pattern which gives small spray droplets without any fine mist and cover the area indicated:

Amt. BL 10,000 to Use: (Tbsp.)	(Fl. Oz.)	Amt. of Water in Sprayer: (Gallons)	Area to be Sprayed (Sq. Ft.)
2	1	1/2	335
4	2	1	670
8	4	2	1300
12	6	3	2000

**NOTE:** Whether using a hose-end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

#### HOW MUCH TO USE: SPOT SPRAY

To spray small areas or individual weeds, do not use hose-end sprayers. Mix 1 oz. (2 tbsp.) in 1/2 gal. water in pressure sprayer and spray as required being certain to "wet" each weed.

#### —FOR COMMERCIAL USE—

**HOW MUCH TO USE—ENTIRE LAWN:** Apply BL 10,000 at the rate of 1 gallon per acre (or 1 quart per 10,880 sq. ft.) in sufficient water for adequate coverage.

**HOSE END SPRAYER—**Use sprayer designed for lawns. Put recommended amount of chemical in sprayer bottle and add water to indicated level and spray on specified area.

**PRESSURE SPRAYERS—**Mix 2 fl. oz. (4 tbsp.) in each gallon of water and uniformly spray each gallon of dilute spray on about 675 sq. ft. of lawn.

**CAUTION:** Harmful if swallowed. Avoid contact with eyes, skin, and clothing. In case of contact, flush with water. Avoid contamination of feed or foodstuffs.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra nor lawns or turf where desirable bentgrass or clovers are present. Use only lawn sprayers. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.

**DO NOT REUSE CONTAINER. WHEN EMPTY, WASH THOROUGHLY AND DESTROY. DO NOT BURN.**

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

2217-536

04/10/1975