



BL 10,000 LAWN WEED KILLER

Contains *TRIMEC® Herbicide

KEEP FROM FREEZING

One Quart Covers 10880 Sq. Ft.

Contains: 2,4-D, MCPP and Dicamba

Kills: DANDELION, KNOTWEED, CLOVER, HENBIT, PLANTAINS, SPURGE, POISON IVY and most other broadleaf weeds.

ACTIVE INGREDIENTS:	
*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	12.875%
**Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	5.840%
***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	1.341%
INERT INGREDIENTS	79.944%
Total	100.000%

Equivalent to
*10.697% 2,4-D Acid or 0.07 lbs. acid per gallon
**4.826% MCPP Acid or 0.12 lbs. acid per gallon
***1.113% Dicamba Acid or 0.097 lbs. acid per gallon

CAUTION: Keep Out of Reach of Children
See Back Panel for Additional Cautions

NET CONTENTS ONE QUART

EPA REG. NO. 2217-536 829-475
EPA EST. NO. 2217-KS-1

*Trimec® is a registered trademark of PBI Gordon Corporation, U.S. Patent No. 3,284,186

Mfr. by  **pbi / GORDON**
CORPORATION
KANSAS CITY, KANSAS 66118

CONTROLS

BEDSTRAW
BLACK MEDIC
BUCKHORN
CHICORY
CHICKWEED
CLOVER
DANDELION
DOCK
GROUND IVY
HEAL-ALL
HENBIT
KNOTWEED
LAMBSQUARTER
LESPEDeza
MALLOW
MORNING GLORY
PEPPERGRASS
PIGWEED
PLANTAINS
POISON IVY
POISON OAK
PURSLANE
RAGWEED
SHEPHERD'S PURSE
SPEEDWELL
SPURGE
WILD CARROT
WILD LETTUCE
WILD ONION
YARROW

and many others

DIRECTIONS FOR USE

WHERE TO USE: BL 10,000 Lawn Weed Killer may be used on Bluegrass, Bermuda, or Zoysia lawns, vacant lots, playgrounds and other similar areas where hard-to-kill weeds are present. This product is especially effective on types of hard-to-kill weeds such as HENBIT, CHICKWEED, KNOTWEED, SPOTTED SPURGE, OXALIS, BLACK MEDIC, and SHEEP SORREL. In addition to these, it controls most other broadleaf lawn weeds, including Dandelion and Plantain, and a partial list is given on the side panel. Do not allow spray to get on vegetables or ornamental plants since injury may result. Since injury can be expected when used on Centipede or St. Augustine grass lawns, "Spot-Spray" weeds only. Clover present will be damaged if sprayed.

WHEN TO USE: Spray at any time during the growing season when weeds are actively growing. Treat Chickweed, Henbit, Wild Onion and Garlic in the fall or in the spring, as needed. On new lawns wait until the turf has reached a height of 2 inches before making an overall spray treatment. "Spot-Spray" weed treatment, however, may be made at any time with a small pressure sprayer, spraying only the weeds at ground level. Wait 3 weeks before putting new grass seed or stolons down in the bare spots. Poor weed control may result if spray is applied during drought or just before rain. For best results, mow before treating. Do not water lawn within 24 hours after treatment.

HOW MUCH TO USE—ENTIRE LAWN

HOSE END SPRAYER: Use a hose-end sprayer designed for lawns. Put recommended amount of chemical (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated. For example, to spray 1,700 sq. feet, measure 5 ozs. (10 tbsp.) in jar and add water to the 10 gallons level and mix. Turn on hose and spray as evenly as possible over 1,700 sq. feet.

Col. 1	Col. 2	Col. 3
Amt. BL 10,000 to Use	Fll. Oz.	Area to Treat, Sq. Ft.
1	1 1/2	170
2	1	340
5	2 1/2	850
10	5	1700
15	7 1/2	2550
20	10	3400
32	16	5440

PRESSURE SPRAYERS: Measure indicated amount of chemical into tank and dilute with water according to the following chart. Use a spray pattern which gives small spray droplets without any fine mist and cover the area indicated.

Amt. BL 10,000 to Use	Amt. of Water in Sprayer	Area to be Sprayed
(tbsp.)	(Gallons)	(Sq. Ft.)
2	1	335
4	2	670
8	4	1300
14	6	2000

NOTE: Whether using a hose end sprayer or pressure sprayer, avoid sprays which are too fine because they can drift and damage ornamental or garden plants.

HOW MUCH TO USE: SPOT SPRAY

To spray small areas or individual weeds, do not use hose end sprayers. Mix 1 oz. (2 tbsp.) in 1/2 gal. water in pressure sprayer and spray as required being certain to "wet" each weed.

—FOR COMMERCIAL USE—

HOW MUCH TO USE—ENTIRE LAWN: Apply BL 10,000 at the rate of 1 gallon per acre or 1 quart per 10,880 sq. ft. in sufficient water for adequate coverage.

HOSE END SPRAYER: Use sprayer designed for lawns. Put recommended amount of chemical in sprayer bottle and add water to indicated level and spray on specified area.

PRESSURE SPRAYERS: Mix 2 fl. oz. (4 tbsp.) in each gallon of water and uniformly spray each gallon of dilute spray on about 675 sq. ft. of lawn.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin, and clothing. In case of contact, flush with water. Avoid contamination of feed or foodstuffs.

Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not contaminate domestic or irrigation water. Do not pour spray solution near these plants. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Do not spray on carpetgrass, dichondra, nor lawns or turf where desirable bentgrass or clovers are present. Use only lawn sprayers. Coarse sprays are less likely to drift. Do not apply to areas underlain by roots of desirable trees and shrubs. This product may be injurious to St. Augustine grass.


DO NOT REUSE CONTAINER WHEN EMPTY. WASH THOROUGHLY AND DESTROY. DO NOT BURN.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

NO
MIXING OR
POURING

AIM AT
WEED AND
SPRAY

PROTECT
FROM
FREEZING



Ready-to-Use
SPOT-TREAT
Contains *TRIMEC® Herbicide
NO MIXING READY-TO-USE

SPECIALLY FORMULATED
FOR "HARD-TO-KILL" LAWN WEEDS

Contains amines of 2,4-D, Mecoprop and Dicamba

ACTIVE INGREDIENTS:	
**Dinethylamine salt of 2,4-dichlorophenoxy acetic acid	.620%
**Dinethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	.281%
**Dinethylamine salt of Dicamba (3,6-dichloro-o-anisic acid)	.064%
INERT INGREDIENTS	99.035%
100.000%	
*Equivalent to 2,4-dichlorophenoxy acetic acid	0.515%
*Equivalent to 2-(2-methyl-4-chlorophenoxy) propionic acid	0.232%
*Equivalent to 3,6-dichloro-o-anisic acid	0.0536%

CAUTION: Keep Out of Reach of Children
See Back Panel for Additional Cautions

KEEP FROM FREEZING


*Trimec® is a registered trademark of P.B.I.—Gordon Corporation, U.S. Patent No. 3,284,186

NET 32 FLUID OUNCES

908-1170

MANUFACTURED BY
PBI-GORDON CORPORATION
KANSAS CITY, KANSAS

KILLS
DANDELIONS
PLANTAINS
CHICKWEED
CLOVER
REDSTRAW
BLACK MEDIC
(Yellow Clover)
ENGLISH DAISY
CURLED DOCK
HEAL-ALL
GROUND IVY
(Creeping Charlie)
KNOTWEED
RAGWEED
SHEPHERD'S PURSE
BUCKWORN
CANADA-THISTLE
CUDWEED
DOLLARWEED
FLEABANE
FILAREE
HEMBIT
LAMBSQUARTERS
MATHWEED
FLORIDA-POSSLEY
MUSTARD
PENNYWORT
PEPPERWEED
PIGWEEED
PRICKLY LETTUCE
POISON IVY
PURSLANE
SHEEP-SORREL
YARROW



DIRECTIONS

Use to "Spot-Treat" weeds in the lawn. For best results apply when weeds are actively growing. Do not water within 24 hours after spraying.

Ready-to-use Spot-Treat is designed to permit easy treatment of lawn weeds without bending or stooping. Aim at weed and spray. Only a short burst is needed. Overspraying only wastes the spray. Effects begin to show after a few days and weeds gradually die. Up to 4 weeks may be required for complete kill. Repeat in 3-4 weeks if necessary.

For lawns with many weeds, requiring an overall treatment, spray with a Turf Herbicide containing TRIMEC Herbicide Concentrate.

CAUTION: Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Wash after each use.

Avoid spraying susceptible plants (flowers, shrubs, etc.). Clovers, Bent grass, Carpetgrass and St. Augustine grass may be injured if contacted. Do not apply within drip line of desirable plants. Do not store near seeds, fertilizers, insecticides or fungicides.

Avoid accidental drift or application to nearby garden plants, flowers, or shrubs. Do not contaminate water used for irrigation or domestic purposes.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 2217-537

908-1170

NO
MIXING OR
POURING



AIM AT
WEED AND
SPRAY

PROTECT
FROM
FREEZING

Ready-to-Use SPOT-TR

Contains *TRIMEC®

NO MIXING

SPECIALLY FORMULATED
FOR "HARD-TO-KILL" LAWN

Contains amines of 2,4-D, Mecoprop and

ACTIVE INGREDIENTS:

*Dimethylamine salt of 2,4-dichlorophenoxyacetic acid
**Dimethylamine salt of 2,4-dimethyl-4-chlorophenoxyacetic acid
***Dimethylamine salt of Dicamba (3,5-dichlorobenzoic acid)

INERT INGREDIENTS:

1. Solvent to 100% methanol
2. Solvent to 20% methanol, 80% isopropyl alcohol
3. Solvent to 100% methanol and

CAUTION: Keep Out of Reach
See Back Panel for Additional

KEEP FROM FREEZING


*Trimec is a registered trademark of
Gordon Chemical Corporation, U.S. Patent No. 3,125,412

NET 32 FLUID OUNCES



MANUFACTURED BY
GORDON CORPORATION

KANSAS CITY, KANSAS



DIRECTIONS

Use to "Spot Treat" weeds in the lawn. For best results apply when weeds are actively growing. Do not water within 24 hours after spraying.

Ready-to-use Spot-Treat is designed to permit easy treatment of lawn weeds without bending or stooping. Aim at weed and spray. Only a short burst is needed. Overspraying only wastes the spray. Effects begin to show after a few days and weeds gradually die. Up to 4 weeks may be required for complete kill. Repeat in 3-4 weeks if necessary.

For lawns with many weeds, requiring an overall treatment, spray with a Turf Herbicide containing TRIMEC Herbicide Concentrate.

CAUTION: Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water; for eyes get medical attention. Wash after each use.

Avoid spraying susceptible plants (flowers, shrubs, etc.). Clovers, Bent grass, Carpetgrass and St. Augustine grass may be injured if contacted. Do not apply within drip line of desirable plants. Do not store near seeds, fertilizers, insecticides or fungicides.

Avoid accidental drift or application to nearby garden plants, flowers, or shrubs. Do not contaminate water used for irrigation or domestic purposes.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 2217-337

77

908-1170