



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Washington, D.C. 20460

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

December 4, 2012

James Kunstman PBI/Gordon Corporation P.O. Box 014090 Kansas City, MO 64101

Subject:

Notification per PR Notice 98-10 (add marketing claims)

Gordon's BL 8000 Lawn Weed Killer

EPA Reg. No. 2217-539

Application Dated November 15, 2012

Dear Dr. Kunstman:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the subject product. The Registration Division (RD) has conducted a review of this request and finds that the action falls within the scope of PRN 98-10. The label submitted with the application has been date-stamped "Notification" and will be placed in our records.

If you have any questions, please contact Mindy Ondish at (703)605-0723 or at ondish.mindy@epa.gov.

Sincerely,

Mindy or the, for

Kathryn V. Montague Product Manager 23 Herbicide Branch Registration Division (7505P)

≎ EPA	Environmental F	d States		Regist		O. Approval Expires 2-28-95 OPP Identifier Number	
	Applie	cation for Pestic	ide - Sect	ion I		STATE OF A SECOND	
1. Company/Product Nur	nber 2217-539	2. EF	A Product Ma Kathryn V	nager Montague	3. 1	Proposed Classification	
4. Company/Product (Na Gordon's E	^{me)} BL 8000 Lawn Weed Ki	iller F	roduct Mana	ger—Team 2	23	X None Restricted	
PBI/Gordon Corpora Post Office Box 014 Kansas City, Missou	090	(b)(i). EPA	my product is Reg. No		ntical in cor		
Onlook in c	no to a now address	Section -	.			STATE OF THE REAL PROPERTY.	
Amendment - Explai	n below.		-	labels in respon	se to NC	TIFICATION	
	ponse to Agency letter dated		"Me Too" Application.			DEC 0 4 2012	
Notification - Explain	below.		Other - Expla	ain below.			
review. e-mail to jkunstman@pbigorde	on.com FAX: 816-421-2731	Section –	III				
1. Material This Product Wi	Il Be Packaged In:			Samuel Contract			
Child-Resistant Packaging Yes* No * Certification must	Unit Packaging Yes No If "Yes" No. pe	Water Soluble Yes No	No. per	M N PI	f Container etal astic lass		
be submitted	Unit Packaging wgt. contain	ner Package wgt.	Container		ther (Specify	y)	
3. Location of Net Contents Label Cor		e(s) Retail Container 1 pint to 1 gallon and 2.5	gallons	5. Location of L On Label On Labelin		ons	
6. Manner in Which Label is	⊠ F	ithograph Paper glued Stenciled	Other_			6 C C C C C C C C C C C C C C C C C C C	
		Section -			cccci		
1. Contact Point (Complete	tems directly below for identifica		ntacted, if neces	ssary, to process			
Name James L.	Kunstman, Ph.D.		of Regulatory	Services	Telephor	ne No. Include Area Code) 816-460-6292	
I acknowledge that any both under applicable I	ents I have made on this form ar knowingly false or misleading st	tatement may be punisha			e	6. Data Application Received (Stamped)	
2. Signature		3. Title					

5. Date

November 15, 2012

James L. Kunstman, Ph.D.

4. Typed Name

Certification with Respect to Label Integrity

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL			
EPA Registration Number	Date Submitted to EPA	Electronic File Name	
2217-539	11/15/2012	002217-00539.20121115.notif-proposed- clean 002217-00539.20121115.notif-proposed- highlighted	

I Certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

Signature

James L. Kunstman, Ph.D. Director, Regulatory Services

Pan I Zunt

November 15, 2012

Date



1217 W. 12th STREET · P.O. BOX 014090 KANSAS CITY, MISSOURI 64101-0090 816-421-4070 · 1-800-821-7925 FAX: 816-474-0462

cecce

November 15, 2012

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard (South Building)
2777 South Crystal Drive
Arlington, VA 22202-4501
Attn: Kathryn V. Montague, PM-23

Dear Ms Montague:

NOTIFICATION: Labeling notification per PRN 98-10: Notifications, Non-notifications and Minor Formulation Amendments: Sections (N)
Gordon's BL 8000 Lawn Weed Killer (EPA Reg. No. 2217-539)

I. Labeling notification:

Request for advertising/marketing claims.

We ask to add advertising claims that are consistent with the Directions for Use previously-approved. Please refer to pages 8 of the draft labeling.

One copy of the proposed labeling with annotations and one copy of the draft labeling without notations are enclosed for your review.

II. Certification statement:

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

III. Enclosures:

- 1. Application for Pesticide Amendment (EPA Form 8570-1)
- 2. One (1) copy the draft labeling with annotations.
- 3. One (1) copy the draft labeling without notations.

If you have any questions, please call me at 816-460-6292 or contact me at jkunstman@pbisordon.com.

Sincerely,

James L. Kunstman, Ph.D. Director of Regulatory Services

GORDON'S BL 8000 LAWN WEED KILLER

EPA Reg. No. 2217-539

NOTIFICATION

DEC 04 2012

Alternate Brand Names: GroundWork Concentrate Weed Killer

ACTIVE INGREDIENT:

2,4-D, dimethylamine salt	7.59%
Mecoprop-p, dimethylamine salt	1.83%
Dicamba, dimethylamine salt	0.84%
OTHER INGREDIENTS:	89.74%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

0.54 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 6.30%

0.13 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 1.51%

0.06 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 0.70%

Isomer Specific by AOAC Method.

Trimec® is a registered trademark of PBI/GORDON CORPORATION.

2,4-D,dimethylamine salt (CAS Registry No. 2008-39-1)

Mecoprop-p, dimethylamine salt (CAS Registry No. 66423-09-4)

Dicamba, dimethylamine salt (CAS Registry No. 2300-66-5)

Note: If the design, format, or small size of the container labels makes it impractical to present the entire ingredient statement on the front panel, then add a referral statement and present this substatement on the back panel: "See back panel for substatement of ingredient statement".

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

First Aid

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, the continue rinsing eye.
- · Call a poison control center or doctor for treatment advice.



1217 West 12th Street Kansas City, Missouri 64101

If swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, experimentally result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Precautions And Restrictions

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
 Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Use only lawn-type sprayers. Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Do not exceed specified dosages for any area.
- Schedule the herbicide applications when air temperatures are 60° to 90°F with good soil moisture.
 Avoid applications during the spring transition of the warm season grasses and during periods of

extremely high temperatures. The tolerance of warm season grasses decreases at air temperatures greater than 90°F and when turfgrass is drought stressed.

- The turf tolerance of centipedegrass and St. Augustinegrass can range from sensitive to tolerant. Do not apply this product to 'Floratam' St. Augustinegrass in Florida.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse sprayer over lawn area only.

2. Where To Use

Gordon's BL 8000 Lawn Weed Killer may be used on bahiagrass, bermudagrass, Kentucky bluegrass, fescue, perennial ryegrass, zoysiagrass, centipedegrass, and St. Augustinegrass lawns, vacant lots, playgrounds and other similar areas where broadleaf weeds are present. This product is especially effective on all hard-to-kill weeds such as poison ivy, poison oak, sheep sorrel and spurge. In addition, BL 8000 Lawn Weed Killer controls most other broadleaf weeds, including sensitive weeds such as chickweed, clover, dandelion, knotweed and plantain. A partial list is given on the side panel.

3. When To Use

1. Spray during the growing season when weeds are actively growing.

- 2. On new lawns, wait until the grass has reached a height of 2 inches before making an overall spray treatment. However, a spot treatment may be made at any time with a small pressure sprayer. Wait 3 weeks before seeding or planting stolons, plugs, or sprigs.
- 3. Poor weed control may result if spray is applied during drought or just before rain.

4. For best results, mow before treating.

5. Do not water lawn within 24 hours after treatment.

4. How Much To Use (Entire Lawn)

4.1 Hose End Sprayers

Use a hose end sprayer designed for lawns. Pour specified amount of product (Column 1) in bottle and add water to indicated level (Column 2), and spray on area shown in Column 3. For example, to spray 2,500 square feet measure 10 fluid ounces (20 tablespoons) in jar, add water to the 10 gallon level and mix. Turn on hose and spray as evenly as possible over 2,500 square feet.

Chart 1: Amounts of Gordon's BL 8000 Lawn Weed Killer for bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns applied with hose end sprayers.

Column 1 Amount of Lawn Weed Killer to Use		Column 2 Fill Bottle to	Column 3
(Tablespoons)	(Fluid Ounces)	(Gallons)	(Square Feet)
10	5	5	1,250
20	10	10	2,500
30	15	15	3,750
40	20	20	5,000

Note: For the grasses in Chart 1, apply the entire contents of this container (32 fluid ounces) on 8,000 square feet of turf.

Chart 2: Amounts of Gordon's BL 8000 Lawn Weed Killer for centipedegrass and St. Augustinegrass lawns applied with hose end sprayers.

Column 1 Amount of Lawn Weed Killer to Use		Column 2 Fill Bottle to	Column 3 Area to Treat
(Tablespoons)	(Fluid Ounces)	(Gallons)	(Square Feet)
10	5	5	2,500
20	10	10	5,000
30	15	15	7,500
40	20	20	10,000

Note: For St. Augustinegrass and centipedegrass apply the entire contents of this container (32 fluid ounces) on 16,000 square feet of turf.

4. 2 Pressure Sprayers

Measure indicated amount of Gordon's BL 8000 Lawn Weed Killer, pour into tank, and dilute with water according to the following chart. Use a spray pattern that gives small [coarse] spray droplets without any fine mist, and cover the area indicated.

Chart 3. For bahiagrass, bermudagrass, Kentucky bluegrass, fescues, perennial ryegrass, and zoysiagrass lawns with pressure or tank type sprayers.

Amount of Lawn Weed Killer to Use		Amount of Water in Sprayer	Area To Treat
(Tablespoons)	(Fluid Ounces)	(Gallons)	(Square Feet)
2	1	1	250
4	2	2	500
6	3	3	750

Amount of Lawn Weed Killer to Use		Amount of Water in Sprayer	Area To Treat
(Tablespoons)	(Fluid Ounces)	(Gallons)	(Square Feet)
2	1	1	500
4	2	2	1,000
6	3	3	1,500

Spot Spraying [Treatment]

To spray small areas or individual weeds, do not use a hose end sprayer. Refer to Chart 3 or Chart 4 for mixing instructions. Spray as needed and be certain to wet the weeds.

4.3 [Spray Delivery System Name]

[When Packaged Accordingly:]
Battery Operated [Powered] Sprayer

SPRAYER: USE DIRECTIONS

The [Spray Delivery System Name] is a patented innovative home lawn, landscape, and perimeter tank spray system. Spot treat [[bugs] [and] [or] [weeds]] with this easy to use system.

- 1. Unscrew the tank cap and fill the tank with water up to the one gallon indicator. [The tank holds one gallon of water.]
- 2. Remove the [sprayer] [wand] [applicator] from the tank holster and unfold the [sprayer] [wand] [applicator].
- 3. Select the [cartridge] [bottle] [insecticide] [herbicide] labeled for intended use. READ THE ENTIRE PRODUCT LABEL PRIOR TO USE.
- 4. Remove the cap from the [cartridge] [bottle] [insecticide] [herbicide]. [Each [cartridge] [bottle] contains 4 oz. of [product] [liquid]. Each 4 oz. [cartridge] [bottle] of concentrate makes one gallon of [liquid] [ready to use]spray.]

CCCCCC

5. Insert the [cartridge] [bottle] [insecticide] [herbicide] into the top of the sprayer and twist until tight.

6. Press the trigger to spot spray [bugs] [weeds] as listed on the product label.

7. When finished spot treating remove the [cartridge] [bottle] [insecticide] [herbicide]. Replace the cap on the [cartridge] [bottle] [insecticide] [herbicide]. Follow label Storage and Disposal instructions for product [insecticide] [herbicide] [cartridge] [bottle].

After spot spraying rinse the [sprayer] [wand] [applicator] according to the RINSE: USE DIRECTIONS.

RINSE: USE DIRECTIONS

- Remove the [cartridge] [bottle] [insecticide] [herbicide]. Ensure that there is water in the one gallon tank.
- 2. Press and hold the [trigger] [sprayer] for 10 seconds to flush the [sprayer] [wand] [applicator] with clean water from the tank.

CHANGE [CARTRIDGE] [BOTTLE]: USE DIRECTIONS

- 1. Remove the [cartridge] [bottle] [insecticide] [herbicide] currently loaded in the [Spray Delivery System Name] .
- 2. Follow label Storage and Disposal instructions for product [insecticide] [herbicide] [cartridge] [bottle].
- Press and hold the [trigger] [sprayer] for 10 seconds to flush the [sprayer] [wand] [applicator] with clean water from the tank.
- Select a new [cartridge] [bottle] [insecticide] [herbicide] labeled for intended use. READ THE ENTIRE PRODUCT LABEL PRIOR TO USE.
- 4. Insert the [cartridge] [bottle] [insecticide] [herbicide] into the top of the sprayer and twist until tight.

5. Press the trigger to spot spray [bugs] [weeds] as listed on the product label.

- 6. When finished spot treating remove the [cartridge] [bottle] [insecticide] [herbicide]. Replace the cap on the [cartridge] [bottle] [insecticide] [herbicide]. Follow label Storage and Disposal instructions for product [insecticide] [herbicide] [cartridge] [bottle].
- 7. After spot spraying, rinse the [sprayer] [wand] [applicator] according to the RINSE: USE DIRECTIONS. Store the [Spray Delivery System Name] OR RINSE and change [cartridge] [bottle].

NOTE: IF OUTPUT SPRAY IS DIMINISHED OR SPUTTERING, DO THE FOLLOWING:

- 1. Check that there is water remaining in the one gallon tank. If not, refill the tank with water.
- 2. Check that there is liquid remaining in the [cartridge] [bottle]. If not, change to a new [cartridge] [bottle].

[Two [cartridges] [bottles] can be stored on the back of the [Spray Delivery System Name]] .

[{Optional:}CHANGING BATTERIES:]

If sprayer does not operate or spray performance declines significantly, change the four AA alkaline batteries in the sprayer handle as follows:

Remove the [cartridge] [bottle].

- 2. Slide the battery cover open in the underside of sprayer handle.
- 3. Allow battery cover to swing open.

4. Remove old batteries.

- Replace with four AA alkaline batteries. Make sure the positive and negative ends match the direction indicated on the inside of the battery door.
- 6. Close battery door and slide the door until latch is closed.

4.4 Limitations on broadcast applications to residential turfgrass

The maximum application rate is 4.0 fl.oz. of product per 1,000 sq.ft. per application [0.74 lb 2,4-D ae, 0.18 lb MCPP-p ae and 0.08 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl.oz. of product per 1,000 sq.ft. [1.47 lb 2,4-D ae, 0.35 lb MCPP-p ae and 0.16 lb dicamba ae per acre].

4.5 Limitations on spot treatments to residential turfgrassSpot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 4.0 fl.oz. per 1,000 sq.ft. per application [0.18 lb MCPP-p acid equivalent per acre]. The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

5. Broadleaf Weeds	3		
Annual yellow sweetclover	Cockle	Mugwort	Speedwell
Aster	Cocklebur	Musk thistle	Spiny amaranth
Austrian fieldcress	Common mullein	Mustard	Spiny cocklebur
Bedstraw	Creeping Jenny (Creeping	Narrowleaf plantain	Spotted catsear
Beggarticks	Charlie)	Narrowleaf vetch	Spotted spurge
Betony, Florida	Cudweed	Nettle	Spurweed
Bindweed, field	Curly dock	Orange hawkweed	Stinging nettle
Bird vetch	Daisy, English	Oriental cocklebur	Strawberry, India mock
Bitter wintercress	Daisy fleabane	Oxalis	Tall nettle
Bittercress, hairy	Daisy, oxeye	Parsley-piert	Tall vervain
Bitterweed	Dandelion	Parsnip	Tansy ragwort
Black-eyed Susan	Dichondra	Pearlwort	Tansy mustard
Black medic	Dogbane	Pennycress	Tanweed
Black mustard	Dogfennel	Peppergrass	Thistle
Blackseed plantain	Dollarweed	Pepperweed	Trailing crownvetch
Blessed thistle	Elderberry	Pigweed	Tumble mustard
Bloodflower milkweed	False dandelion	Pineywoods bedstraw	Tumble pigweed
Blue lettuce	Falseflax	Plains coreopsis (tickseed)	Velvetleaf
Blue vervain	False sunflower	Plantain	Venice mallow
Bracted plantain	Fiddleneck	Poison ivy	Virginia buttonweed
Brassbuttons	Florida pusley	Poison oak	Virginia creeper
Bristly oxtongue	Frenchweed	Pokeweed	Virginia pepperweed
Broadleaf dock	Galinsoga	Poorjoe	Wavyleaf bullthistle
Broadleaf plantain	Goathead	Prairie sunflower	Western clematis
Broomweed	Goldenrod	Prickly lettuce	Western salsify
Buckhorn	Ground ivy	Prickly sida	White mustard
Buckhorn plantain	Gumweed	Prostrate knotweed	Wild aster
Bulbous buttercup	Hairy fleabane	Prostrate pigweed	Wild buckwheat
Bull thistle	Hawkweed	Prostrate spurge	Wild carrot
Bullnettle	Healall	Prostrate vervain	Wild four-o'clock
Burclover	Heartleaf drymary	Puncturevine	Wild garlic
Burdock	Heathaster	Purslane, common	Wild geranium
Burning nettle	Hedge bindweed	Ragweed	Wild lettuce
Burweed	Hedge mustard	Red sorrel	Wild marigold
Buttercup	Hemp	Redroot pigweed	Wild mustafd
Buttonweed	Henbit	Redstem filaree	Mild orion
Canada thistle	Hoary cress	Rough cinquefoil	Wild parsnip
Carolina geranium	Hoary plantain	Rough fleabane	Wild radish co
Carpetweed	Hoary vervain	Russian pigweed	Wild rape
Catchweed bedstraw	Horsenettle	Russian thistle	Wild strawperry
Catnip	Jimsonweed	Scarlet pimpernel	Wild sweet potato
Catsear	Knawel	Scotch thistle	Wild vetch
Chickweed, common	Knotweed	Sheep sorrel	Woodsorrel
Chickweed, mouseear	Kochia	Shepherdspurse	Woolly croton
Chicory	Lambsquarters	Slender plantain	Woolly morningglory
Cinquefoil	Lespedeza	Smallflower galinsoga	Woolly plantain
Clover, crimson	Mallow	Smooth dock	Wormseed
Clover, hop	Matchweed	Smooth pigweed	Yarrow
Clover, red	Mexicanweed	Sorrel	Yellow rocket
Clover, strawberry	Milk vetch	Sowthistle	Yellowflower pepperweed
	Morningglory	Spanishneedles	and other broadleaf weeds
Clover, sweet Clover, white	Mouseear hawkweed	Spiny sowthistle	and other broadlear weeds
Ciover, write	INIOUSEE AI HAWKWEEU	Topiny sowunsue	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES. INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTIAL. CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

Mixes with Water – Concentrate

C

 Statements which may appear on different label components depend configuration. 	ding on packaging	
See next panel for additional Precautionary Statements and First Aid Net Contents: EPA Est. No		
 2. Advertising claims that may be presented on container labeling, advand other marketing/sales promotional materials: • One quart covers up to 8,000 square feet. 	vertisements, broch	uros
Keep from freezing.	erecet.	c
 Controls dandelion, chickweed, knotweed, henbit, plantain, clover, spunlawn weeds including those shown on the side panel. Kills even the tough weedsroots and all! Controls even the tough weedsroots and all! Treats northern and southern grasses. Kills the root. One quart treats up to 16,000 square feet. (Southern grasses) Water based. One quart covers up to square feet. Won't harm lawn grass when used according to label directions. Kills [Controls] 200+ broadleaf weeds. Starts working overnight Won't harm lawn grass (when used as directed) Controls (Number) weeds! 	ge and many other b	roadleaf

- Treated areas for this container vary from 16,000 square feet (Northern Grasses) to 32,000 square feet (Southern Grasses)
- CONTROLS
 - SENSITIVE WEEDS: Bedstraw, Black Medic, Buckhorn, Chickweed, Clover, Dandelion, Knotweed, Lambsquarters, Peppergrass, Plantain, Purslane, Ragweed, Sheperdspurse, and Wild Lettuce.
 - HARD-TO-KILL WEEDS: Chicory, Creeping Charlie, Creeping Jenny, Dock, Ground Ivy, Heal-All, Lespedeza, Mallow, Morningglory, Oxalis, Pigweed, Poison Ivy, Poison Oak, Sheep Sorrel, Speedwell, Spurge, Wild Carrot, Wild Onion, Yarrow and many other broadleaf weeds.
- For 16 fl.oz. retail container: Lawn areas for this container vary from 4,000 sq.ft.(Northern grasses) to 8,000 sq.ft.(Southern grasses).
- For 32 fl.oz. retail container: Lawn areas for this container vary from 8,000 sq.ft.(Northern grasses) to 16,000 sq.ft.(Southern grasses).
- For One Gallon retail container: Lawn areas for this container vary from 32,000 sq.ft.(Northern grasses) to 64,000 sq.ft.(Southern grasses).
- · For outdoor use
- Controls dandelion, spurge, chickweed and many other hard-to control weeds.
- · Controls dandelion, spurge, chickweed and many other listed weeds
- · Won't harm lawns when used as directed.
- Contains Trimec® Herbicide
- Kills Over 200 Weeds -- [see complete weed listing inside]
- One quart treats up to 8,000 sq. ft.(Northern grasses) to 16,000 sq. ft. (Southern grasses)*
- One quart treats 8,000 to 16,000 sq.ft.*
- Makes up to XXX[*]
- . * When mixed XXX of product with YYY of water.
- Controls Weeds

DOCUMENT CONTROL INFORMATION

- 1. Unique Label Identifier: 002217-00539.20121115.notif-proposed-highlighted.doc
- 2. Reason for Issue: ad claims

