

AFTER TEXT

TRIMEC[®] HERBICIDE

A BROADLEAF WEED & BRUSH KILLER

Controls Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge, Ash, Aspen, Brambles, Kudzu, Oak, Willows and many other species of broadleaf weeds and brush; some of which are listed on this label.

— ONE GALLON COVERS 2 TO 2 1/2 ACRES —
KEEP FROM FREEZING

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	25.93%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	13.85%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid	2.76%
INERT INGREDIENTS:	57.46%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- *2.03 lbs. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 21.54%
- **1.08 lbs. 2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 11.45%
- ***0.21 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.29%

TRIMEC[®] is a registered trademark of PBI/GORDON CORPORATION.
Isomer Specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN


DANGER

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES: In case of eye contact, immediately flush eyes with plenty of water for 15 minutes. Call a physician at once.
- IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly. If irritation develops, get medical attention.
- IF SWALLOWED: Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Call a physician at once. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED: Remove victim to fresh air and apply respiration if indicated.
See additional Precautionary Statements elsewhere on this label.

NET CONTENTS U.S. GALLONS

881,883,885,886.PC8/1092 AP101492
EPA REG. NO. 2217-543
EPA EST. NO. 2217-KS-1
Mfd. by PBI/GORDON CORPORATION
KANSAS CITY, MISSOURI 64101



2 of 4

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS
AND FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

DANGER: Corrosive, causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed or inhaled. Wear long pants, long-sleeved shirt, shoes and gloves when mixing, loading or applying this product. Wear face shield or goggles when mixing, loading or applying this product. When handling this product, wear chemical resistant gloves. Avoid exposure to spray mist. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not contaminate domestic or irrigation waters. Do not apply when weather conditions favor drift away from target area.

When cleaning of equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not reenter or permit workers to reenter treated areas until sprays have dried or dusts have settled.

STORAGE & DISPOSAL

- KEEP FROM FREEZING -

STORAGE: Store in original container in a locked storage area. To prevent cross-contamination, do not store near other pesticides, fertilizers, or seeds, food or feed.

PRODUCT DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: FOR METAL CONTAINERS - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by other procedures approved by state and local authorities. **FOR PLASTIC CONTAINERS** - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

USE PRECAUTIONS/BROADLEAF WEEDS IN ORNAMENTAL LAWNS & TURF: Do not apply this product through any type of irrigation system. Avoid drift of spray mist to vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near desirable plants. Do not use on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Use only lawn-type sprayers. Avoid fine sprays; coarse sprays are less likely to wind-drift. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area; be particularly careful within the

dripline of tree and other ornamental species. Do not apply to newly seeded grasses until well established. Do not broadcast apply when air temperature exceeds 85 °F; some injury may be expected with spot treatments when air temperatures exceed 85 °F. Seed can be safely sown 3 to 4 weeks after application. Some injury can be expected to turf when used on centipedegrass or St. Augustinegrass lawns. "Spot-spray" weeds on these types of grasses to avoid possible damage that could occur from an broadcast treatment. After using this product, clean sprayer with soap or detergent and rinse thoroughly before using other pesticides in it. Care should be taken not to make applications where runoff could contaminate bodies of water or spray drift could carry the chemical to food crops or grazing lands where cattle, sheep, goats, swine or poultry would be exposed.

ORNAMENTAL LAWNS & TURF (cool season grasses other than closely-mowed bentgrasses): Maximum kill of weeds will be obtained from spring or early fall applications when weeds are actively growing. Avoid spraying during long, excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours after application. Do not apply to newly seeded turf until after the second or third mowing. Grass seed can be safely sown 3 to 4 weeks after application at recommended rates. Do not use on dichondra and carpetgrass.

General Application: Apply at the rate of 3 1/4 to 4 pints in 20 to 260 gallons of water per acre (1.2 ounces to 1.5 ounces in 1/2 to 6 gallons of water per 1,000 square feet). Use higher rates when using the higher volume of water per acre.

Controlled Droplet Applicator: Add 2 1/2 pints to the HERBI container and fill with water. Spray contents over 33,000 square feet. Avoid overlapping between spray patterns.

Bentgrass Turf: On closely-mowed bentgrass (putting greens, bowling greens, and show lawns), apply at the maximum rate of 1 ounce per 1,500 square feet in 5 gallons of water. Do not apply when grass is under stress from drought or hot temperatures. Use only application equipment that is capable of distributing uniform droplets over the entire area to be sprayed. Slight yellowing of turf will disappear after about one week.

Note: Care should be taken to avoid overdosing bentgrasses or injury may result. If hand-type sprayers are used it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom. Sweeping movements with a multiple nozzle boom will result in heavy local over-application and subsequent turf discoloration or injury.

USE PRECAUTIONS/BRUSH & BROADLEAF WEEDS ALONG ROADS, RIGHTS-OF-WAYS, DRAINAGE DITCH BANKS & OTHER SIMILAR NON-CROP AREAS: Do not apply this product through any type of irrigation system. Avoid spray drift to cotton, soybeans, tomatoes, tobacco, grapes, fruit trees, flowers or garden crops, and all other hormone herbicide-sensitive desirable plants. Do not apply near sensitive plants since small quantities of wind-drifted herbicides may cause severe injury. Do not apply in areas underlaid by roots of desirable trees and shrubs. Do not apply herbicide when wind speed is sufficient to cause drift. Do not apply herbicides when a temperature air inversion exists. An air inversion may be detected by creating a smoke column and observing a layering effect. Do not apply when temperature exceeds 85 °F and humidity is high. To aid in avoiding spray drift, use coarse sprays and low pressure. Do not use nozzles which produce fine spray droplets under high pressure. The use of thickening agents or antidrift additives and drift-reducing equipment is of value in preventing spray drift. Care should be taken not to make applications where runoff could contaminate bodies of water or spray drift could carry the chemical to food crops or grazing lands where cattle, sheep, goats, swine or poultry would be exposed.

BRUSH & BROADLEAF WEEDS ALONG ROADSIDES, RIGHTS-OF-WAYS, DRAINAGE DITCH BANKS & OTHER SIMILAR NON-CROP AREAS: To effectively control brush, all leaves, stems and suckers must be thoroughly wetted to the ground. Apply when plants come into full leaf (spring) to the time plants begin to go dormant. Best results, however, will be obtained when brush and weeds are young and actively growing. Do not cut brush or weeds until the herbicide has translocated throughout the plant causing root death.

General Application: Mix 3/4 to 1 gallon in 30 to 100 gallons of water per acre (2.25 ounces to 3 fluid ounces in 1 to 3 gallons of water per 1,000 square feet). Heavy dense stands of brush and weeds may require the high rate with the high water volume. Use only application equipment that is capable of distributing uniform droplets over the entire area to be sprayed.

BRUSH:

Ash
Aspen
Birches
Black Cherry
Brambles

Elms
Gooseberry
Honey Locust
Kudzu
Multiflora Rose

Oak
Shor.leaf Pine
Thistle
Willows

WEEDS:

Bedstraw
Black Medic
Buckhorn
Burdock
Chicory
Chickweed
Clover
Dandelion
Dock
Ground Ivy
Healall

Henbit
Knotweed
Lambsquarter
Lespedeza
Mallow
Morningglory
Peppergrass
Pigweed
Plantains
Poison Ivy
Poison Oak

Purslane
Ragweed
Sheep Sorrel
Shepherdspurse
Speedwell
Spurge
Wild Carrot
Wild Garlic
Wild Lettuce
Wild Onion
Yarrow

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.