[2217-545]

GORDON'S

spraying on rulinture, aspirant the or plastics.

Do not apply to pets or birds. Do not treat in same area with fish tanks or caged birds.

warning: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperature above 130° F may cause bursting. Never throw container into fire or incinerator.

NOTICE: Seller warrants that this product con-

indoor pantungspray

avay from you. Hold dispenser about 18" from plant surfaces and press valve, keeping fingers out of stream. Using short sweeping motion, spray all visible surfaces and infested sur-

PB - GORDON CORPORATION
KANSAS CITY, KANSAS 66118

CAUTION

Harmful if swallowed. Avoid breathing spray mist and contact with skin or eyes. Wash thoroughly after using spray. Do not contaminate booking utensils, food, or drinking water. Avoid arming on furniture sephalt tile or plastice

forms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

ACTIVE INGREDIENTS:

34 3

Pyrethrins	0.02%
Rotenone	0.13%
Other cube resins	0.26%
Technical Piperonyl Butoxide*	0.25%
Petroleum Distiliate	0.08%
INERT INGREDIENTS	25%
Total 1	

*Equivalent to 0.2% (Butylcarbity!) (6 piperonyl) ether and 0.05% related compounds

T.M. Registered, P.B.I.

675-874

E.P.A. Est. No. 11715-TN-1

DIRECTIONS FOR USE: Read all directions. Remove cap shake can, and hold con-

indooi plant spray

! DE UNDER NO. TO ATTACHED - JAMENTS.

CAUTION:

Keep Out of Reach of Children See Side Panels for Additional Caution

EPA Reg. No. 2217-545

NET WEIGHT 5 OUNCES

darters of the traction of POISON HUGIS Maves and the stalks or _ SUBJECTIFE. Do not overapply or sprayiwhen leaves are wet or directly in hot sun. Newspapers should be used to catch spray missing the plants.

> Use for control of mites, aphids, exposed thrips, white fly and mealybug on African Violets, Azalea, Boston Fern, Fuschia, and Ivy. Frequent application is needed for mealybugs as they are hard to control.