



tomato and vegetable fungicide

Active Ingredient:

Maneb (Manganese ethylene bisdithiocarbamate) 80%

Inert Ingredients*

*Includes ZnSO₄

U.S. Patent 2,974,154


CAUTION:

NET WEIGHT 10 POUNDS

E.P.A. Reg. No. 2217-548

Controls—Brown rot and shot-hole (fungus). Use 1 1/2 to 2 1/2 pounds per acre of water at red bud, early bloom and 2 1/2 to 3 1/2 weeks later.

GORDON'S



tomato and vegetable fungicide

This product is believed compatible with other commonly used pesticides and may be combined for use with other insecticides, miticides or fungicides if and as recommended for a particular use unless such product directions specifically caution against any such combination applications. Always observe any listed timing restrictions between last application and harvest to avoid the possibility of residues in excess of the tolerances established.

DIRECTIONS FOR USE VEGETABLE CROPS

Tomatoes—Early and late blight, anthracnose, Septoria leaf spot and gray leaf spot (Stemphylium). Start applications when first cluster fruit is well formed and repeat at 7 to 10 day intervals. (To avoid injury do not use on young seedlings in greenhouses). More frequent applications may be necessary starting when plants are set in the fields if weather conditions favor disease. Tomato plants treated with this product tend to develop heavy foliage, so to assure complete coverage increase spray gallonage accordingly. Where anthracnose is a serious problem on canning tomatoes apply 1½ to 2 tablespoons per gallon of water.


applications when disease threatens and repeat at 7 to 10 day intervals as needed.

FRUITS

Peaches and Nectarines—Shot-hole (fungus) and peach leaf curl. Apply dormant and popcorn spray. For brown rot blossom blight apply at pink bud and 25% to 75% of full bloom. For Scab apply at pink bud, bloom, petal fall, shuck and in cover sprays at 7 to 14 day intervals. Use 1½ to 2 tablespoons per gallon of water. Do not apply to nectarines later than 14 days before harvest. If applied to peaches within 14 days of harvest, remove residues by brushing. Do not apply to peaches later than 2 days before harvest.

Apples—Apple scab, rust, black rot, bullseye rot, fly speck, sooty blotch, bitter rot and white rot. Use this product 1½ to 2 tablespoons per gallon of water. For a protective schedule begin applications in the pre-pink sprays and repeat in subsequent pre-bloom, bloom and cover sprays in accordance with local state spray schedules.

 **GORDON CORPORATION**
SAN ANTONIO, TEXAS



2217-546