

TURF 45

LAWN FUNGICIDE

2½ LBS. covers up to 10,000 square feet.

ACTIVE INGREDIENT:

2,4-Dichloro-6-(0-chloroanilino)-s-triazin ..... 50%

INERT INGREDIENTS..... 50%

TOTAL 100%


KEEP OUT OF REACH OF CHILDREN

DANGER

See Side Panels for additional pre-cautionary statements and Statement of Practical Treatment.

NET WEIGHT 2.5 POUNDS

**ACCEPTED**  
**REREGISTERED**  
**JUL 15 1988**  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2217-551



STOP! READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND FOLLOW  
DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS:

Hazards to Humans & Domestic Animals

**DANGER:** Causes irreversible eye damage. Harmful ~~or fatal~~ if swallowed. Avoid contact with skin. Do not get in eyes or on clothing. Wear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Protective clothing should be worn during periods of exposure, such as during application or when contacting treated foliage. ANILAZINE may cause temporary allergic skin reaction in a few susceptible persons. This condition should be treated as an allergic dermatitis. There is no evidence of after effects or permanent injury.

Statement of Practical Treatment

In case of contact in eyes, flush with water for 15 minutes. Get medical attention. In case of contact with skin, wash with plenty of soap and water.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or wetlands. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or incorporate spills.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow children or pets to play on lawn for 24 hours after treatment.

STORAGE & DISPOSAL

**STORAGE:** Do not contaminate water, food, or feed by storage or disposal. Store in original container in a locked storage area inaccessible to children or pets.

**DISPOSAL:** If bag contains unused materials, securely wrap original bag in several layers of newspaper and discard in trash. If bag is empty, do not reuse. Discard bag in trash.

Turf 45 gives effective control of Rhizoctonia (brown patch), Sclerotinia (dollar spot), Copper Spot, Rust, Snow mold (Typhula), and Helminthosporium diseases, such as melting-out of Kentucky bluegrass, red leaf spot of bentgrasses, leaf blotch of Bermuda grass, Helminthosporium blight of fescues, brown blight of rye grasses, zonate eye spot, and Helminthosporium leaf spot of bluegrasses, bentgrasses, and fescues.

#### HOW TO USE:

In Garden Hose Sprayer: Put Turf 45 into sprayer jar, fill with water and stir until uniform mixture is obtained. Apply spray mixture uniformly to lawn area. For disease prevention, use 4 ounces (approximately 1 cup) Turf 45 and spray to 1,000 square feet. For spot treatment of diseased areas, use 6 ounces (approximately 1½ cups) Turf 45 and thoroughly wet these areas.

In Compressed Air Sprayers: For use in hand or knapsack-type sprayers, add 4 ounces (approximately 1 cup) of Turf 45 to 2½ gallons of water and apply to 1,000 square feet (50 ft. x 20 ft.) of lawn or other turf area. For spot treatment of heavily diseased areas, add 6 ounces (approximately 1½ cups) to 2½ gallons of water and thoroughly wet these areas.

In Field Sprayers: Schedule applications in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service for specific use information and for State regulations which may contain additional restrictions or requirements.

Recommended Applications--To control Brown Patch, Copper Spot, Dollar Spot, Helminthosporium Leaf Spot, Melting-out, and Rust, mix 4 to 6 ounces Turf 45 in 5 gallons of water and apply per 1,000 square feet. Repeat at 1 to 2 weeks intervals throughout the spring and summer months.

#### WHEN TO APPLY:

For Foliar Turfgrass Diseases: Begin application in the spring before diseases appear and repeat at 10 to 14 days intervals. During extended periods of wet weather, applications every 7 days may be required. Continue treatments throughout the spring and summer months. If necessary, water lawn before applying Turf 45.

For Control of Snow Mold (Typhula): Use 4 ounces in 5 gallons water and apply to 1,000 square feet. Apply after the first killing frost in the fall. A second application should be made when the lawn or other turfgrass area is free of snow in mid-winter. Apply a third application after the final melting of snow in late winter or early spring.

Limited Warranty. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because it is impossible to eliminate all risks even

In Compressed Air Sprayers: For use in hand or knapsack-type sprayers, add 4 ounces (approximately 1 cup) of Turf 45 to 2½ gallons of water and apply to 1,000 square feet (50 ft. x 20 ft.) of lawn or other turf area. For spot treatment of heavily diseased areas, add 6 ounces (approximately 1½ cups) to 2½ gallons of water and thoroughly wet these areas.

In Field Sprayers: Schedule applications in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service for specific use information and for State regulations which may contain additional restrictions or requirements.

Recommended Applications--To control Brown Patch, Copper Spot, Dollar Spot, Helminthosporium Leaf Spot, Melting-out, and Rust, mix 4 to 6 ounces Turf 45 in 5 gallons of water and apply per 1,000 square feet. Repeat at 1 to 2 weeks intervals throughout the spring and summer months.

#### WHEN TO APPLY:

For Foliar Turfgrass Diseases: Begin application in the spring before diseases appear and repeat at 10 to 14 days intervals. During extended periods of wet weather, applications every 7 days may be required. Continue treatments throughout the spring and summer months. If necessary, water lawn before applying Turf 45.

For Control of Snow Mold (Typhula): Use 4 ounces in 5 gallons water and apply to 1,000 square feet. Apply after the first killing frost in the fall. A second application should be made when the lawn or other turfgrass area is free of snow in mid-winter. Apply a third application after the final melting of snow in late winter or early spring.

Limited Warranty. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

Exclusion of Other Warranties and Remedies. Except where such disclaimers and excursions are specifically prohibited by applicable law. THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESS OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES EXPRESS OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty. BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.