

FLY BAIT may be used in federally inspected poultry and red meat packing plants. In edible products areas of these plants, FLY BAIT may be used only after operations have ceased for the day, and baits must be removed before operations are resumed. Apply the bait in cardboard containers which can be readily collected before resuming operations.

RESTRICTIONS

Do not apply directly to animals. Do not contaminate drinking water, milk, or milk-handling equipment. Do not treat portions of building where poultry feed or which animals can lick. Direct applications to walls, floors, or other surfaces previously treated with lime, whitewash, or other alkaline materials may be ineffective.

Keep bait and solution out of reach of children and pets.

CAUTION

Harmful if swallowed. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Do not contaminate feed or food.

Do not apply this bait in dwellings.

If poisoning occurs, get prompt medical aid. Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

U.S.D.A. Reg. No. 2217-557

608-671



FLY BAIT

MANUFACTURED BY
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

**READ LABEL DIRECTIONS BEFORE
TO CONTROL ROACHES IN STORES
BUILDINGS:** Place a small quantity
and put in protected areas such as
of reach of children and pets, and
utensils, or drinking water. Wet the

**TO CONTROL FLIES AND ROACHES
FOOD PROCESSING PLANTS, PATIO
LISHMENTS, STABLES, STOCK PEN
AREAS, OUTDOOR LATRINES, AND
FOUND:**

NOTE: Use only as a dry bait in
plants, after operations have ceased
are resumed.

NOTE: Battery poultry establishments
alleyways and walk-ups.

FLY BAIT can be used dry "as is"
described below. Good sanitation
promptly. Apply FLY BAIT daily until
needed.

As a Dry Bait: Scatter lightly in
horizontal areas. Use 4 ounces per
bait is barely visible. Apply directly
and congregate. For faster kill, apply
tery poultry establishments scatter
and walk-ways. Keep out of reach.

As a Water Solution for Fly Control:
of water, add 2 cups of corn syrup
solved. Sprinkle on floors, window
buildings, and outside of these buildings
bothersome. Do not use liquid bait in
plants.

Where Floor Area is Lacking: Where
assure control (for example, a door
boards, or strips of cardboard and
flies gather. First wet them, they
may be treated by dissolving FLY BAIT
Use only enough water to bring

ACCEPTED

READ LABEL DIRECTIONS BEFORE USE

TO CONTROL ROACHES IN STORES, RESTAURANTS, AND SIMILAR COMMERCIAL BUILDINGS: Place a small quantity of dry bait on pieces of paper, cardboard and put in protected areas such as in closets, under sinks or refrigerators, out of reach of children and pets, and where dying insects cannot fall into food, utensils, or drinking water. Wet the bait slightly for increased killing efficiency.

TO CONTROL FLIES AND ROACHES IN AND AROUND DAIRY BARN, MILK ROOMS, FOOD PROCESSING PLANTS, PATIOS, PICNIC AREAS, BATTERY POULTRY ESTABLISHMENTS, STABLES, STOCK PENS, KENNELS, ANIMAL HOSPITALS, GARBAGE AREAS, OUTDOOR LATRINES, AND OTHER PLACES WHERE THESE INSECTS ARE FOUND:

NOTE: Use only as a dry bait in edible products areas of food processing plants, after operations have ceased for the day. Remove bait before operations are resumed.

NOTE: Battery poultry establishments must be confined to use of dry bait on alleyways and walk-ups.

FLY BAIT can be used dry "as is" or dissolved in water and sprinkled as described below. Good sanitation is essential to fly control. Remove wastes promptly. Apply FLY BAIT daily until flies are controlled and then repeat as needed.

As a Dry Bait: Scatter lightly in strips on floors, window sills, and other clean horizontal areas. Use 4 ounces per 1,000 square feet, scattering lightly so that bait is barely visible. Apply directly to garbage and manure where flies breed and congregate. For faster kill, apply FLY BAIT to a moistened surface. In battery poultry establishments scatter on droppings under cages, on alleyways and walk-ways. Keep out of reach of poultry.

As a Water Solution for Fly Control: Pour 1 pound of FLY BAIT into 1 gallon of water, add 2 cups of corn syrup or 1 pound of sugar, and stir until dissolved. Sprinkle on floors, window sills, etc., of dairy barns and other farm buildings, and outside of these buildings where fly population is intense or bothersome. Do not use liquid bait in edible products areas of food processing plants.

Where Floor Area is Lacking: Where there is not enough clean floor area to assure control (for example, a dirt floor barn or a stockyard), treat sacks, boards, or strips of cardboard and place them inside or outside building where flies gather. First wet them, then apply FLY BAIT. Vertical surfaces indoors may be treated by dissolving FLY BAIT in water and painting or spraying it on. Use only enough water to bring mixture to syrup-like consistency.

608-671

ACCEPTED WITH COMMENTS