

GORDON'S TRIMEC® WEED & FEED 4 23-3-7

(Sixteen, Eighteen, or Twenty) Pounds Covers Up To 5,000 Square Feet.

Controls Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds including those listed on the back panel.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	0.655%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	0.316%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid).....	0.072%
INERT	98.957%
INGREDIENTS	
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- *10.888 lbs. or 0.544% 2, 4-Dichlorophenoxyacetic acid equivalent per ton.
 - **5.229 lbs. or 0.261% 2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
 - ***1.206 lbs. or 0.061% 3, 6-dichloro-o-anisic acid equivalent per ton.
- Isomer Specific by AOAC Methods.
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Statement of Practical Treatment

In case of contact, wash skin with plenty of soap and water.
For eyes, flush with water for 15 minutes and get medical attention.
See back panel for additional Precautionary Statements.

97 MAY 29 A9:26


RECD EPA/OPP/DPD1

NET WEIGHT (16, 18, or 20) POUNDS

587 APXXXXXX
EPA REG. NO. 2217-559
EPA EST. NO. 2217-KS-1
MANUFACTURED BY:



KANSAS CITY, MO 64101





READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

USE PRECAUTIONS:

Do not use this product where desirable clovers are present or allow the material to wind drift onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

[This product may stain concrete and stone. Where spilling occurs, sweep off immediately, then flush with water. (For use for products when iron is present in the fertilizer.)]

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 4 controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. This product may be applied to established lawns of bermudagrass, bentgrass, bahiagrass, centipedegrass, Kentucky bluegrass, fescues, perennial ryegrass, zoysiagrass, and St. Augustinegrass. Do not use on dichondra or carpetgrass, closely-mowed bentgrass, putting greens or St. Augustinegrass. Do not apply this product to Floratam® St. Augustinegrass in Florida. In addition, a 23-3-7 fertilizer provides a balanced supply of plant nutrients or plant food. ~~needed for a normal season's growth.~~

Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

The maximum application rate to turf is 1.0 pound 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Spreader Settings	
Scott.....	6
Lawn Beauty.....	5
Jackson.....	7½
Sears.....	4

The above spreader settings are approximate. (16 pound bag: Calibrate your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.) (18 pound bag: Calibrate your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet) (20 pound bag: Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet)

23-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N).....	23%
Water Insoluble Nitrogen	5%
Available Phosphate (P ₂ O ₅).....	3%
Soluble Potash (K ₂ O).....	7%

WEEDS		
Bedstraw	Knotweed	Ragweed
Black Medic	Lambsquarters	Sheep Sorrel
Buckhorn	Lespedeza	Shepherdspurse
Burdock	Mallow	Speedwell
Chickweed	Morningglory	Spurge
Chicory	Oxalis	Thistle
Clover	Peppergrass	Wild Carrot
Dandelion	Pigweed	Wild Garlic
Dock	Plantains	Wild Lettuce
Ground Ivy	Poison Ivy	Wild Onion
Healall	Poison Oak	Yarrow
Henbit	Purslane	and many more

LIMITED WARRANTY AND DISCLAIMER. The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

APPENDIX

I. Previously Approved Fertilizer Grades:

- ◆ Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅, and K₂O include the following:

8-3-16	18-3-5	24-4-12	30-0-6
10-6-4	18-5-9	25-3-3	33-3-5
13-2-3	20-4-4	25-4-4	
15-3-6	20-10-5	26-3-3	
15-5-7	20-10-10	26-3-5	
16-4-8	23-3-7	26-4-6	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

15-3-6 GUARANTEED ANALYSIS	
Total Nitrogen (N).....	15%
12.4% Ammoniacal Nitrogen	
0.3% Urea Nitrogen	
2.3% Coated slow release urea nitrogen	
Available Phosphate (P ₂ O ₅).....	3%
Soluble Potash (K ₂ O).....	6%
Calcium (Ca).....	2%
Sulfur (S).....	14%
Derived from: Ammonium Sulfate, Sulfur-Coated Urea, Ammonium Phosphate, Rock Phosphate, Muriate of Potash	

II. Advertising claims that may be presented on the retail containers:

- ◆ (For products where iron is present) - This product may stain concrete and stone. Where spilling occurs, sweep off immediately, then flush with water.
- ◆ (Brand Name) enhances the fertilizer portion of this product with Agrico Turf[®], a stabilized form of urea nitrogen that remains in the stable available ammonium form for longer periods of time. Agrico Turf[®], when watered into the soil, attaches itself to clay particles where it remains for use by the plant, longer than if unstabilized. The result is a lawn that is greener longer than if fertilized with unstabilized urea nitrogen. Added iron and other essential micronutrients promote root development, color enhancement, and seed production.

BEST AVAILABLE COPY