

2217-559

6/29/2009

10f8

Jackie



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

2217-559

Date of Issuance:

JUN 29 2009

Term of Issuance:

Name of Pesticide Product:

Gordon's Trimec
Weed & Feed 4

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

PBI/Gordon Corporation
1217 West 12th Street
Kansas City, MO 64101

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you:

1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
2. Make the following revisions to the final printed labeling:
 - a. Per the acute toxicity review, the Hazards to Humans and Domestic Animals must be revised to read:
 "CAUTION
 Harmful if swallowed. **Harmful if absorbed through skin.** Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, shoes, socks and **chemical-resistant** gloves made of any waterproof material. **Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.**"
 - b. Make the following minor changes to the Storage and Disposal section:
 1. Change the subheading "Container Disposal" to "**Pesticide Disposal and Container Handling.**"
 2. Replace "If Empty" with "**Nonrefillable Container**" and place "If Empty" after "Do not reuse this container."

Continued on Page 2

Signature of Approving Official:

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Date:

JUN 29 2009

2018

Page 2

EPA Reg. No. 2217-559

Reregistration Notice

- c. Assure that the Net Contents/Net Weight always appear on the front panel of the product label
See FIFRA 2(q)(2)(C)(iii); 40 CFR 156.10(a)(1)(iii).
3. A stamped copy of your labeling is enclosed for your records. You must submit one (1) copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this Notice or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Enclosure

Stamp 1111100
 3 of 8
ACCEPTED
 with COMMENTS
 In EPA Letter Dated:
JUN 29 2009
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.

GORDON'S TRIMEC® WEED & FEED 4

2217-559

EPA Reg. No. 2217-559

ACTIVE INGREDIENT:

- Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 0.65%
- Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid 0.16%
- Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) 0.07%

INERT INGREDIENTS 99.12%
 TOTAL 100.00%

THIS PRODUCT CONTAINS:

- 10.80 lbs 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.54%
- 2.60 lbs (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.13%
- 1.20 lbs 3,6-dichloro-o-anisic acid equivalent per ton or 0.06%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.	



Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)]

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use this product where desirable clovers are present. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

Advisory for fertilizers containing iron:

This product may stain concrete and stone. Where spilling occurs, sweep off immediately, then flush with water.

INSTRUCTIONS:

Gordon's Trimec® Weed & Feed 4 controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. This product may be applied to established lawns of bermudagrass, bentgrass, bahiagrass, centipedegrass, Kentucky bluegrass, fescues, perennial ryegrass, zoysiagrass, and St. Augustinegrass. Do not use on dichondra or carpetgrass. Do not apply this product to 'Floritam' St. Augustinegrass in Florida. In addition, a 23-3-7 fertilizer provides plant nutrients or plant food.

Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).

3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Spreader Settings	
Scotts	6
Lawn Beauty	5
Jackson	7.5
Sears	4

The above spreader settings are approximate.

16 pound bag: Calibrate your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

18 pound bag: Calibrate your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet

20 pound bags: Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (0.94 lbs 2,4-D ae, 0.23 lb MCP-P ae and 0.10 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (1.88 lbs 2,4-D ae, 0.46 lb MCP-P ae and 0.20 lb dicamba ae per acre).

23-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N)	23%
Water Insoluble Nitrogen	5%
Available Phosphate (P2O5)	3%
Soluble Potash (K2O)	7%

Labeling Requirements:

- Information regarding the contents and levels of metals in this product is available on the Internet as <http://www.aapfco.org/metals.htm>.

WEEDS			
Bedstraw	Ground ivy	Peppergrass	Speedwell
Black medic	Healall	Pigweed	Spurge
Buckhorn	Henbit	Plantain	Thistle
Burdock	Knotweed	Poison ivy	Wild carrot
Chickweed	Lambsquarters	Poison oak	Wild garlic
Chicory	Lespedeza	Purslane	Wild lettuce
Clover	Mallow	Ragweed	Wild onion
Dandelion	Morningglory	Sheep sorrel	Yarrow
Dock	Oxalis	Shepherdspurse	and many more

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

CONTAINER DISPOSAL: **If Empty** — Do not reuse this container. Place in trash or offer for recycling if available. **If Partly Filled** — Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

- Previously Approved Fertilizer Grades: Fertilizer grades with nutrient contents expressed as the percentages of N,P2O5, and K2O include the following:

8-3-16	18-3-5	24-4-12	27-3-4
10-6-4	18-5-9	25-3-3	30-0-6
13-2-3	20-4-4	25-4-4	33-3-5
15-3-6	20-10-5	26-3-3	
15-5-7	20-10-10	26-3-5	
16-4-8	23-3-7	26-4-6	

The fertilizer grade guaranteed analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations. The following guaranteed analyses presentations have previously been submitted as notifications:

15-3-6 GUARANTEED ANALYSIS	
Total Nitrogen (N)	15%
12.4% Ammoniacal Nitrogen	
0.3% Urea Nitrogen	
2.3% Coated slow release urea nitrogen	
Available Phosphate (P2O5)	3%
Soluble Potash (K2O)	6%
Calcium (Ca)	2%
Sulfur (S)	14%
Derived from: Ammonium Sulfate, Sulfur-Coated Urea, Ammonium Phosphate, Rock Phosphate, Muriate of Potash	

- An advisory for fertilizers containing iron:
This product may stain concrete and stone. Where spilling occurs, sweep off immediately, then flush with water.

- (Brand Name) enhances the fertilizer portion of this product with Agrico Turf®, a stabilized form of urea nitrogen that remains in the stable available ammonium form for longer periods of time. Agrico Turf®, when watered into the soil, attaches itself to clay particles where it remains for use by the plant, longer than if unstabilized. The result is a lawn that is greener longer than if fertilized with unstabilized urea nitrogen. Added iron and other essential micronutrients promote root development, color enhancement, and seed production.
- Contains Trimec® Herbicide.
- Excellent for lawns of bluegrass, ryegrass, fescue, St. Augustine, bentgrass, and zoysiagrass.
- Can be used on most types of turf grasses. Will not burn when used as directed.
- Contains 1% iron to prevent and correct yellowing due to iron deficiency.
- (Brand Name) effectively controls the following weeds when used as directed.
- Controls (Number) weeds! (Listing of weeds may include the following.)

WEEDS			
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly oxtongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cress	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed

8018

WEEDS			
Clover, hop	• Morningglory	Sowthistle	• Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	• Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	and other broadleaf weeds

- **Green Cycle**

- Spring

- Crabgrass Control & Lawn Food

- Summer

- Weed & Feed

- Fall

- Lawn Food

- Winter

- Fall & Winter Lawn Food

- (Sixteen, Eighteen, or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds including those listed on the back panel.

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00559.20081105.proposed.doc

2. **Reason for Issue:** Various changes pertaining to re-registration. Re-format label.