



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 23, 2017

Mr. Eric Smith
Federal Registration Manager
PBI/Gordon Corporation
P.O. Box 014090
Kansas City, MO 64101

Subject: Notification per PRN 98-10 – Minor updates/clarifications, updated weed list
Product Name: Gordon's Trimec Weed & Feed 4
EPA Registration Number: 2217-559
Application Date: February 12, 2016
Decision Number: 514326

Dear Mr. Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you have any questions, you may contact Kathryn Montague at 703-305-1243 or via email at montague.kathryn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

01/23/2017

GORDON'S TRIMEC® WEED & FEED 4

EPA Reg. No. 2217-559

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	0.65%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.16%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.07%

INERT/OTHER INGREDIENTS	99.12%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

- 10.80 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.54%
- 2.60 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.13%
- 1.20 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.06%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

~~Note to reviewer: This language has been moved from the Appendix.~~

~~Mandatory Labeling Statements which will appear on different label components depending on packaging configuration.~~

~~Referral Statement: See attached booklet back panel for complete Precautionary Statements and Directions for Use (including First Aid and Storage and Disposal). [for similar language dependent on configuration of container label]~~

Guaranteed Analysis: See appendix for example.

Net Contents: _____

EPA Est. No. _____

Company Name and Address:

PBI/Gordon Corporation
1217 West 12th Street
Kansas City, Missouri 64101



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, shoes, socks, and chemical resistant gloves made of any waterproof material. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or on clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

[Environmental Hazards (alternate per PRN 2008-1)

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use this product where desirable clovers are present. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.

- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

Gordon’s Trimec® Weed & Feed 4 controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns **as listed**. This product may be applied to established lawns of bermudagrass, bentgrass, bahiagrass, centipedegrass, Kentucky bluegrass, fescues, perennial ryegrass, zoysiagrass, and St. Augustinegrass. Do not use on dichondra or carpetgrass. Do not apply this product to ‘Floritam’ St. Augustinegrass in Florida. In addition, a **23-3-7 [XX-XX-XX]** fertilizer provides plant nutrients or plant food.

Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and growing actively, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application **[(0.94 lb 2,4-D ae, 0.23 lb MCP-p ae and 0.10 lb dicamba ae per acre per application)]**. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. **[(1.88 lb 2,4-D ae, 0.46 lb MCP-p ae and 0.20 lb dicamba ae per acre)]**.

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, the following language on equipment settings of drop spreaders, rotary spreaders, and other granular applications is optional – it may be presented on the container labels at your discretion.]

[Spreader Settings [spreader images/charts]

Spreader Settings	
Scotts	6
Lawn Beauty	5
Jackson	7.5
Sears	4

[The above spreader settings are approximate.

16 pound bag: Calibrate your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

18 pound bag: Calibrate your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet

20 pound bags: Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet

[Coverage: To determine the square footage of the application area, multiply the length of the lawn by its width. This equals your total square footage. Deduct any non-lawn areas (driveway, sidewalk, house, etc.) One bag covers XXXX sq. ft. or XX pounds per 1,000 sq. ft.]

[End optional language]

23-3-7 GUARANTEED ANALYSIS	
Total Nitrogen (N)	23%
Water Insoluble Nitrogen	5%
Available Phosphate (P2O5)	3%
Soluble Potash (K2O)	7%

WEEDS			
Bedstraw	Ground ivy	Peppergrass	Speedwell
Black medic	Healall	Pigweed	Spurge
Buckhorn	Henbit	Plantain	Thistle
Burdock	Knotweed	Poison ivy	Wild carrot
Chickweed	Lambsquarters	Poison oak	Wild garlic
Chicory	Lespedeza	Purslane	Wild lettuce
Clover	Mallow	Ragweed	Wild onion
Dandelion	Morningglory	Sheep sorrel	Yarrow
Dock	Oxalis	Shepherdspurse	and many more

WEEDS CONTROLLED:

WEEDS [253]			
Amaranth, purple	Cupid shaving Brush	Mugwort	Speedwell, common
Amaranth, slender	Cudweed	Mustard, ball	Speedwell, corn
Aster, heath	Daisy, English	Mustard, black	Speedwell, ivyleaf
Aster, white heath	Daisy, oxeye	Mustard, blue	Speedwell, Persian
Aster, purple	Dandelion, common	Mustard, hedge	Speedwell, purslane
Aster, white prairie	Dandelion, false	Mustard, Indian,	Speedwell, slender
Artichoke	Day flower	Mustard, tansy	Speedwell, snow
Austrian fieldcress	Deadnettle	Mustard, tumble	Speedwell, thymeleaf
Bedstraw, Catchweed	Dichondra	Mustard, white	Spiny amaranth
Bedstraw, smooth	Dock, broadleaf	Mustard, wormseed	Spurge, spotted
Bedstraw, Northern	Dock, curly	Nettle, stinging (burning)	Spurge, prostrate
Bedstraw, pinewoods	Dogbane	Nettle, tall	Spurweed
Beggarticks, hairy	Dogfennel	Nettle, wood	Star of Bethlehem
Betony, Florida	Dollarweed	Parsley-piert	Strawberry, India mock
Bindweed, Field	(Pennywort)	Parsnip	Tansy ragwort
Bindweed, Hedge	Dove weed	Pearlwort	Tanweed
Bitter Wintercress	Elderberry	Pennycress (frenchweed)	Thistle, blessed
Bittercress, hairy	Falseflax	Pepperweed, perennial	Thistle, bull
Bitterweed	False sunflower	Pepperweed, Virginia	Thistle, Canada
Black-eyed Susan	Fiddleneck	(Peppergrass)	Thistle, flodman (prairie)
Black Medic	Filaree, redstem	Pigweed, prostrate	Thistle, musk
Blood flower milkweed	Filaree, whitestem	Pigweed, redroot	Thistle, platte
Blue lettuce	Fleabane, daisy	Pigweed, russian	Thistle, plumeless
Brassbuttons	(annual)	Pigweed, smooth	Thistle, Russian
Bristly oxtongue	Fleabane, rough	Pigweed, tumble	Thistle, scotch
Broomweed	Galinsoga, hairy	Pineapple weed	Thistle, wavyleaf
Buckhorn	Galinsoga, smallflower	Plains coreopsis (tickseed)	Thistle, yellow (horrible)
Bullnettle	Goldenrod	Plantain, blackseed	Trailing crownvetch
Burclover	Ground ivy	Plantain, bracted	Velvetleaf
Burdock	Groundsel	Plantain, broadleaf	Vervain, blue

Burweed, lawn (spurweed)	Gumweed	Plantain, buckhorn	Vervain, hoary
Buttercup, bulbous	Hairy fleabane	Plantain, hoary	Vervain, prostrate
Buttercup, creeping	Hawkweed, meadow	Plantain, narrowleaf	Vervain, tall
Buttonweed, common (poorjoe)	Hawkweed, mouseear	Plantain, slender	Vetch, bird
Buttonweed, Virginia	Hawkweed, orange	Plantain, woolly	Vetch, hairy
Carolina geranium (wild)	Hawkweed, yellow	Poison ivy	Vetch, milk
Carpetweed	Healall	Poison oak	Vetch, narrowleaf
Catnip	Heartleaf drymary	Pokeweed	Vetch, wild
Catsear, spotted (false dandelion)	Hemp	Prairie Sunflower	Virginia creeper
Chamber bitter	Henbit	Prickly lettuce (compass plant)	Western clematis
Chickweed, common	Hoary cress	Prickly sida	Western salsify
Chickweed, mouseear	Horsenettle	Prostrate knotweed	Wild aster
Chickweed, sticky	Horseweed	Puncturevine	Wild buckweed
Chicory	Jimsonweed	Purslane, common	Wild carrot
Cinquefoil	Johnny-Jumpup violet	Pusley, brazil	Wild four-o'clock
Clover, crimson	Knawel	Pusley, Florida	Wild garlic
Clover, alyce	Knotweed, prostrate	Pusley, large flower	Wild lettuce
Clover, hop	Kochia	Ragweed, bur	Wild marigold
Clover, rabbit foot	Lambsquarters	Ragweed, common	Wild mustard (yellow rocket)
Clover, red	Lespedeza	Ragweed, lanceleaf	Wild onion
Clover, strawberry	Mallow, alkali	Ragweed, western	Wild parsnip
Clover, yellow sweet	Mallow, bristly	Redstem filaree	Wild radish
Clover, white sweet	Mallow, common	Rough cinquefoil	Wild rape
Clover, white	Mallow, dwarf	Scarlet pimpernel	Wild strawberry
Cockle	Mallow, venice	Shepard's purse	Wild sweet potato
Cocklebur	Marcela	Smartweed, ladysthumb	Wild violet
Cocklebur, oriental	Matchweed	Smartweed, pale	Woodsorrel, creeping
Cocklebur, spiny	Mexicanweed	Smartweed, Pennsylvania	Woodsorrel, violet
Common mullein	Moneywort	Smooth chaff-flower	Woodsorrel, yellow (Oxalis)
Corn Spurry	Morningglory, bigroot	Smoothdock	Woolly croton
Creeping Beggarweed	Morningglory, ivyleaf	Sorrel, red (sheep)	Wormseed
Creeping Jenny (creeping Charlie)	Morningglory, Mexican	Sowthistle, annual	Yarrow, common
	Morningglory, red	Sowthistle, spiny	Yarrow, western
	Morningglory, tall	Spanish needles	Yarrow, yellow
	Morningglory, wolly	Speedwell, bi-lobed	

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non refillable container. Do not reuse this container. **If empty:** Place in trash or offer for recycling if available. **If Partly Filled:** Call your local solid waste agency [or www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND

DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

[Note to Reviewer: The following information was moved to the front panel]

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: _____
- EPA Est. No. _____

12. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

Herbicide Benefit Claims:

- (Brand Name) effectively controls the following weeds when used as directed.
- Controls (Number) weeds! (Listing of weeds may include the following.)
- Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds including those listed on the back panel.
- Effectively controls [dandelion][chickweed][spurge] and many listed problem weeds
- Controls dandelion, chickweed, spurge and many other problem weeds, as listed.
- Controls [over] [XXX] [plus] [listed] weeds [as listed]!
- Kills even the tough weeds like Dandelion, Plantain, and Clover
- Kills dandelions and many other broadleaf weeds [as listed]
- [Brand name] contains the broad spectrum weed killer Trimec®
- Easy one-step application of fertilizer and weed control to feed your lawn while killing the weeds

WEEDS			
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catscar
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Botany, Florida	Cudweedcove	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pinoywoods bedstraw	Trailing crownvetch
Blackseed plantain	False dandelion	Plains coreopsis	Tumble mustard
Blessed thistle	Falseflax	Plantain	Tumble pigweed
Bloodflower milkweed	False sunflower	Poison ivy	Velvetleaf
Blue lettuce	Fiddleneck	Poison oak	Venice mallow
Blue vervain	Florida pusley	Pokeweed	Virginia buttonweed
Bracted plantain	Frenchweed	Poorjoe	Virginia creeper
Bristly ox tongue	Galinsoga	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Goldenrod	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Ground ivy	Prickly sida	Western clematis
Broomweed	Gumweed	Prostrate knotweed	Western salsify
Buckhorn	Hairy fleabane	Prostrate pigweed	White mustard
Buckhorn plantain	Hawkweed	Prostrate spurge	Wild aster
Bulbous buttercup	Healall	Prostrate vervain	Wild buckwheat
Bull thistle	Heartleaf drymary	Puncturevine	Wild carrot
Bullnettle	Heath aster	Purslane, common	Wild four-o'clock

WEEDS			
Burclover	Hedge bindweed	Ragweed	Wild garlic
Burdock	Hedge mustard	Red sorrel	Wild geranium
Burning nettle	Henbit	Redroot pigweed	Wild lettuce
Burweed	Hoary cross	Redstem filaree	Wild marigold
Buttercup	Hoary plantain	Rough cinquefoil	Wild mustard
Buttonweed	Hoary vervain	Rough fleabane	Wild onion
Canada thistle	Horsenettle	Russian pigweed	Wild parsnip
Carolina geranium	Jimsonweed	Russian thistle	Wild radish
Carpetweed	Knawel	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Knotweed	Scotch thistle	Wild strawberry
Catnip	Kochia	Sheep sorrel	Wild sweet potato
Catsear	Lambsquarters	Shepherdspurse	Wild vetch
Chickweed, common	Lespedeza	Slender plantain	Woodsorrel
Chickweed, mouseear	Mallow	Smallflower galinsoga	Woolly croton
Chicory	Matchweed	Smooth dock	Woolly morningglory
Cinquefoil	Mexicanweed	Smooth pigweed	Woolly plantain
Clover, crimson	Milk vetch	Sorrel	Wormseed
Clover, hop	Morningglory	Sowthistle	Yarrow
Clover, red	Mouseear hawkweed	Spanishneedles	Yellow rocket
Clover, strawberry	Mugwort	Speedwell	Yellowflower pepperweed
Clover, sweet	Musk thistle	Spiny amaranth	

Fertilizer Benefit Claims:

- (Brand Name) enhances the fertilizer portion of this product with Agrico Turf®, a stabilized form of urea nitrogen that remains in the stable available ammonium form for longer periods of time. Agrico Turf®, when watered into the soil, attaches itself to clay particles where it remains for use by the plant, longer than if unstabilized. The result is a lawn that is greener longer than if fertilized with unstabilized urea nitrogen. Added iron and other essential micronutrients promote root development, color enhancement, and seed production.
- In addition, [product name] provides a supply of plant food needed for a healthy lawn. This fertilizer analysis has been specifically formulated in [the northwest] for [northwest] lawns.
- Contains 1% iron to prevent and correct yellowing due to iron deficiency.
- Quick green that lasts up to two months
- **Promotes a quick greening lawn**
- Builds thick, green lawns from the roots up
- Fertilizer blend with nonstaining iron [only when chelated iron, iron citrate or iron succinate are the iron source]
- General Information: The numbers in the product analysis mean: 24-3-6
 Nitrogen – is absorbed by the plants in the greatest quantity. It is necessary for rapid growth and promotes deep green leaves and stems. Nitrogen will not fix itself in the soil, and if unused by the plants, will percolate past the root zone with watering and rains, and must continually be replaced.
 Phosphorus – is essential in all phases of plant growth, but is particularly associated with early maturity, the formation of seeds and fruit, increased root growth, and disease resistance. Phosphorus stays in the soil until it is used by the plant, usually requiring fewer applications than nitrogen, and its effectiveness is enhanced by adding organic matter. Additional phosphorus is recommended for early season plantings, as availability is lowest in cool weather.
 Potassium – plants take large amount of this nutrient, almost as much in some cases as nitrogen. It contributes to the overall vigor of the plant, building strong stems and leaves, helping to overcome the effect of heavy foot traffic or adverse weather. Potassium levels in many intermountain area soils are naturally high, but become depleted as plants use it, and in heavily irrigated areas such as lawns and gardens.
 Iron – usually plentiful in Intermountain soils, but many times is not available for plants to use. Proper application will reduce symptoms from “Iron Chlorosis” which are a yellowing of the leaves with dark veins. Adding iron will improve the color of most turf grasses to a deep green.


Calibrate your spreader on 275 square feet and adjust to apply at rate of 1 pound of product per 275 square feet. / Calibre su esparcidor para 275 pies cuadrados y ajústelo para aplicar a una tasa de 1 libra de producto por cada 275 pies cuadrados.	
Drop Spreader Models	Setting
<i>Modelos de esparcidor por goteo</i>	<i>Ajuste</i>
• Scotts® AccuGreen® 1000	8.0
• Scotts® AccuGreen® 2000	8.0
• Scotts® AccuGreen® 3000	8.0
• Earthway®	13.0
• Precision Red Devil™	4.0
• Vigoro®	4.0
Rotary Spreader Models	Setting
<i>Modelos de esparcidor giratorio</i>	<i>Ajuste</i>
• Scotts® SpeedyGreen® 1000	8.0
• Scotts® SpeedyGreen® 2000	8.0
• Scotts® SpeedyGreen® 3000	8.0
• Earthway®	15.0
• Precision Red Devil™	11.0
• Vigoro®	7.5

General Marketing Claims:

- [This product] Contains Trimec® Herbicide.
- With Trimec® Herbicide
- Feed Your Lawn And Control Your Weeds
- Setting may vary with age, model and condition of spreader. Check your spreader over a known area. This bag covers [xxx sq.ft.] or [yy] pound per [zzz sq.ft.] [aaa feet by bbb feet].
- UNCONDITIONAL MONEY BACK GUARANTEE If you are not completely satisfied with this product, bring the bag and your proof of purchase to the store where it was purchased for refund of purchase price of this product.
- Excellent for lawns of bluegrass, ryegrass, fescue, St. Augustine, bentgrass, and zoysiagrass.
- Can be used on most types of turf grasses. Will not burn when used as directed.
- (Sixteen, Eighteen, or Twenty) Pounds Covers Up To 5,000 Square Feet.
- Lily Miller Ultragreen feeds and protects all year long!
- Filled by weight not by volume. Product may settle after packaging.
- Setting Your Lawn Spreader
- These settings are a guide. They were determined by landscape professionals using new spreaders. The condition of your spreader, your walking speed, the roughness of the lawn, and even the humidity can affect how fast the lawn food comes out. Watch the flow rate. If fertilizer appears to be coming out either slower or faster than you are covering the lawn, adjust the setting. Close the spreader when filling, turning, or stopping.
- Satisfaction Guaranteed - If for any reason you are not satisfied with this product, please send proof of purchase to the address shown and we will gladly refund your purchase price.
- Cover up to 5,000 sq.ft.
- Treats up to 10,000 sq.ft. [for 40lb bag]
- Information regarding the raw materials composing this product can be obtained by writing to the Guarantor listed on this label and referring to the batch number found on this container.
- IFA Country Stores [Helping to Grow the Things You Love®] Since 1923
- [Product Name] builds turf and kills dandelion and many other broadleaf weeds [as listed]
- [Product Name] is a special formulation of premium quality lawn fertilizers blended with proven Trimec® Herbicide for effective controls of most broadleaf weeds
- Made in the U.S.A.
- Manufactured in the U.S.A
- Pellitized
- [Note to reviewer: This statement may be placed on containers to comply with California Prop 65: [Attention][Notice]: This product contains a chemical known to the state of California to cause cancer and birth defects or other reproductive harm.]

Referrals to other products used sequentially:

- Northwest Best(TM) THRIVELAWN PLAN



- **Green Cycle**
Spring: Crabgrass Control & Lawn Food
Summer: Weed & Feed

Fall: Lawn Food

Winter: Fall & Winter Lawn Food

- 2. IFA 4-Step Lawncare: Apply Apr, May
- IFA Premium 4 Step Program
 1. (Feb-Mar-Apr) IFA Premium Crabgrass & Spurge
 2. (Apr-May or June) IFA Premium Weed & Feed Fertilizer Blend with Iron
 3. (June-July-Aug) IFA Premium Spring & Summer
 4. (Sept-Oct-Nov) IFA Premium Fall & Winter

Fertilizer Labeling Requirements

- A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000 square feet each growing season. When applied as directed this product supplies (1.1) pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local agricultural extension agent for the proper amount of nitrogen to be applied in your area.
- Fertilizer Labeling Requirements by Association of American Plant Food Control Officials (AAPFCO): Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metal.htm>.
- Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Advisory For Fertilizers Containing Iron:

[only when chelated iron, iron citrate or iron sucrate are NOT the iron source]

Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

This product may stain concrete and stone. Where spilling occurs, sweep off immediately, then flush with water.

2. State mandated fertilizer labeling requirements:

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P ₂ O ₅)	X%
Soluble Potash (K ₂ O)	X%
Sulfur (S)	X%
x.x% free sulfur	
Iron (Fe).....	X%
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.	
*A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.	
This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P₂O₅ and K₂O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are

relevant to your labeling, please place in the Directions for Use under the heading “State Restrictions”]

- California, Oregon, Washington: “Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: [Note to reviewer: maximum number of applications in this section is dually based upon maximum ai load as well as fertilizer analysis effect on maximum nitrogen rates in different regions in Florida]:
 - In North Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In Central Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In South Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
- Florida: “Apply only to actively growing turf.”
- Florida: “Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- Idaho: “Apply only as directed” [or similar statement]
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”
- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the “Soil Erosion and Sediment Control Act,” P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots.”
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: “In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency.”
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]
- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program."

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00559.20161027.notif-proposed-highlighted

2. Reason for Issue: EPA Comments #6 = a use of other ingredients; formatting – move mandatory language to front panel, fertilizer language to appendix; add updated weeds list; clarify label language; clarify optional Non-FIFRA language for distributors; ad claims + EPA Comments #1 + EPA Comments #2 + EPA Comments #3 + EPA Comments #4 + EPA Comments #5

Internal Usage Notes
Associated Container Label: Supplemental Distributors