



Weed-'N-Feed 5

23-3-7

Sixteen Pounds Covers 5,000 Sq. Ft.

KILLS: Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds; some of which are listed on the back panel.

ACTIVE INGREDIENTS:

**Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	0.827%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.399%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.090%
INERT INGREDIENTS	98.684%
TOTAL 100.000%	

THIS PRODUCT CONTAINS:

- *13.75 lbs or 0.687% 2,4-Dichlorophenoxyacetic acid per lb
- **6.60 lbs or 0.330% 2-(2-methyl-4-chlorophenoxy) propionic acid per lb
- ***1.50 lbs or 0.075% 3,6-dichloro-o-anisic acid per lb
- *Isomer Specific by AOAC Method

BEST AVAILABLE COPY

APR 20 1987 22-17-560

KEEP OUT OF REACH OF CHILDREN
CAUTION


STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements

*TRIMEC® is a registered trademark of PBI GORDON CORPORATION.

NET WEIGHT 16 POUNDS



KANSAS CITY, KANSAS 64110



READ THE ENTIRE LABEL FIRST. OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

1/83
PM 23
22-17-560

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash.
CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

USE PRECAUTIONS

Do not use this product where desirable clovers are present or allow the material to "wind drift" onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

INSTRUCTIONS

Weed & Feed 5 contains TRIMEC® herbicide. TRIMEC controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. In addition, a 23-3-7 fertilizer provides a generous supply of plant food needed for a normal season's growth. Possible retreatment may be necessary if (a) lawn is heavily overran by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on dichondra, carpetgrass, closely-mowed bentgrass putting greens or St. Augustine lawns.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non target plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Be patient, TRIMEC works slowly and gently and is friendly to your lawn.

SPREADER SETTINGS

Scott	6
Lawn Beauty	5
Jackson	7 1/2
Sears	4

The above spreader settings are approximate. Check your spreader on 310 sq. ft. and adjust to apply at the rate of 1 lb. per 310 sq. ft.

FERTILIZER GUARANTEE

Total Nitrogen (not less than)	23%
Water Insoluble Nitrogen (not less than)	5%
Available Phosphoric Acid (not less than)	3%
Soluble Potash (not less than)	7%

WEEDS

DEQSTRAW	DANDELION	LESPEDEZA	POISON IVY	SPURGE
BLACKBERRY	DOCK	MALLOW	POISON OAK	THISTLE
BUCKLEWHEAT	GROUNDIVY	MORNING GLORY	PURSLANE	WILD CARROT
BINDWEED	FLAT TAIL	OXALIS	WIGWAG	WILD GARLIC
CHICORY	HENBIT	PEPPERCRESS	SHEEP SORE	WILD LETTUCE
CHICKWEED	KNOTWEED	PIGWEEED	SHEPHERD'S PURSE	WILD ONION
CLOVER	LAMBSQUARTER	PLANTAINS	SPEEDWELL	YARROW

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA REG. NO. 2217-560
EPA EST. No. 2217-KS-1

588 1084

1/73



Weed-'N-Feed 5

23-3-7

Eighteen Pounds Covers 5,000 Sq. Ft.

KILLS: Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds; some of which are listed on the back panel.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4 Dichlorophenoxyacetic acid	0.827%
**Dimethylamine Salt of 2 (2-methyl-4-chlorophenoxy) propionic acid	0.399%
***Dimethylamine Salt of Dicamba (3,6 dichloro-o-anisic acid)	0.090%
INERT INGREDIENTS	98.684%
TOTAL	100.000%

THIS PRODUCT CONTAINS

- *13.75 lbs or 0.687% 2,4 Dichlorophenoxyacetic acid per ton
- **6.60 lbs or 0.330% 2 (2-methyl-4-chlorophenoxy) propionic acid per ton acre
- ***1.50 lbs or 0.075% 3,6 dichloro-o-anisic acid per ton
- Isomer Specific by AOAC Method

Handwritten: Feb 23 1987

Handwritten: 287-560

BEST AVAILABLE COPY

KEEP OUT OF REACH OF CHILDREN

CAUTION


STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautionary statements.

TRIMEC is a registered trademark of PBI GORDON CORPORATION

NET WEIGHT 18 POUNDS



KANSAS CITY, KANSAS 64118



READ THE ENTIRE LABEL FIRST. OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals
CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

Environmental Hazards
 Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

USE PRECAUTIONS

Do not use this product where desirable clovers are present or allow the material to "wind drift" onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

INSTRUCTIONS

Weed & Feed 5 contains TRIMEC* herbicide. TRIMEC controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. In addition, a 23-3-7 fertilizer provides a generous supply of plant food needed for a normal season's growth. Possible retreatment may be necessary if (a) lawn is heavily overmown by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on dichondra, carpetgrass, closely-mowed bentgrass putting greens or St. Augustine lawns.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non-target plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Be patient, TRIMEC works slowly and gently and is friendly to your lawn.

SPREADER SETTINGS

Scott	6
Lawn Beauty	5
Jackson	7 1/2
Scars	4

The above spreader settings are approximate. Check your spreader on 275 sq. ft. and adjust to apply at the rate of 1 lb. per 275 sq. ft.

FERTILIZER GUARANTEE

Total Nitrogen (not less than)	23%
Water Insoluble Nitrogen (not less than)	5%
Available Phosphoric Acid (not less than)	3%
Soluble Potash (not less than)	7%

WEEDS

BIRD SHAW	DANDELION	LESPEDeza	POISON IVY	SPURGE
BLACK MEDIC	DOCK	MALLOW	POISON OAK	THISTLE
BUCKHORN	GROUND IVY	MORNING GLORY	PURSLANE	WILD CARROT
BURDOCK	HEAL ALL	OXALIS	RAGWEED	WILD GARLIC
CHICORY	HENBIT	PEPPERGRASS	SHRETT'S HORSE	WILD LETTUCE
CHICKWEED	KNOTWEED	PLINKWEED	SHRETT'S PURSE	WILD ONION
CLOVER	LAMBSQUARTER	PLANTAINS	SPELDWELL	YARROW

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA REG NO 2217-560
 EPA EST No 2217-KS-1

580-1004

2063

5



Weed-'N-Feed 5

23-3-7

Twenty Pounds Covers 5,000 Sq. Ft.

KILLS: Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds; some of which are listed on the back panel.

ACTIVE INGREDIENTS:

**Dimethylamine Salt of 2,4 Dichlorophenoxyacetic acid	0.827%
**Dimethylamine Salt of 2 (2-methyl-4-chlorophenoxy) propionic acid	0.399%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.090%
INERT INGREDIENTS	98.684%
TOTAL 100.000%	

THIS PRODUCT CONTAINS

- *13.75 lbs. or 0.687% 2,4 Dichlorophenoxyacetic acid per ton
- **6.60 lbs. or 0.330% 2 (2-methyl-4-chlorophenoxy) propionic acid per ton
- ***1.50 lbs. or 0.075% 3,6 dichloro o-anisic acid per ton
- *Isomer Specific by AOAC Method

APR 23 1980
2217-560

KEEP OUT OF REACH OF CHILDREN
CAUTION


STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water; for eyes, flush with water for 15 minutes and get medical attention.

See back panel for additional precautions and treatments

*TRIMEC is a registered trademark of PBI-GORDON CORP.

NET WEIGHT 20 POUNDS



KANSAS CITY, KANSAS 64116



READ THE ENTIRE LABEL FIRST. OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. Wash thoroughly after handling.

Environmental Hazards

Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water used for irrigation or domestic purposes. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused product, securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

USE PRECAUTIONS

Do not use this product where desirable clovers are present or allow the material to "wind drift" onto flowers, vegetables or ornamental shrubbery. Do not apply in areas underlain by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

INSTRUCTIONS

Weed & Feed 5 contains TRIMEC® herbicide. TRIMEC controls a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others commonly found in home lawns. In addition, a 23-3-7 fertilizer provides a generous supply of plant food needed for a normal season's growth. Possible retreatment may be necessary if (a) lawn is heavily overmown by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on dichondra, carpetgrass, closely-mowed bentgrass putting greens or St. Augustine lawns.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto non-target plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

Be patient, TRIMEC works slowly and gently and is friendly to your lawn.

SPREADER SETTINGS

Scott	6
Lawn Beauty	5
Jackson	7 1/2
Scars	4

The above spreader settings are approximate. Check your spreader on 250 sq. ft. and adjust to apply at the rate of 1 lb. per 250 sq. ft.

FERTILIZED GUARANTEE

Total Nitrogen (not less than)	23%
Water Insoluble Nitrogen (not less than)	5%
Available Phosphoric Acid (not less than)	3%
Soluble Potash (not less than)	7%

WEEDS

IFOSHAW	DANDELION	LESPEDEZA	POISON IVY	SPURGE
BLACK MEDIC	DOCK	MALLOW	POISON OAK	THISLE
BUCKHORN	GROUND IVY	MORNING GLORY	POISON PIE	WILD CARROT
BUNDOCK	HEAL ALL	OXALIS	HAGWEEED	WILD GARLIC
CHICORY	HENBIT	PEPPERCRESS	SHEEP SORE	WILD LETTUCE
CHICKWEED	KNOTWEED	PIGWEEED	SHEEPHOUSSE	WILD ONION
CLOVER	LAMBSQUARTER	PLANTAINS	SPELDWELL	YARROW

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

EPA REG. NO. 2217-560

EPA EST. No. 2217-KS-1

588 1084