

2217-561

19-08-2009

Tackett
17



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

James Kuntsman
PBI / Gordon Corporation
1217 West 12th Street
Kansas City, Missouri 64101

OCT 8 2009

Dear Mr. Kuntsman:

Subject: EPA Reg. 2217-561 / Weed 'N Feed 6 Label Amendment

The labeling referred to above is acceptable provided you make the following changes:

- 1) Delete "General" from "General Precautions and Restrictions".

A stamped copy of the label is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. This label supersedes all other previously accepted labels. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

Sincerely,

Joanne Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

2
17

WEED-'N-FEED 6

A 28-2-8 Lawn Fertilizer Plus Trimec®
Contains 2,4-D, MCPP-p, and Dicamba. Covers 5,000 Square Feet
Controls chickweed, dandelion, knotweed, plantain, henbit, spurge
and many other weeds as listed.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	1.040%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid ...	0.230%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid).....	0.108%

INERT INGREDIENTS:	<u>98.622%</u>
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 17.20 lbs. 2, 4-dichlorophenoxyacetic acid equivalent per ton or 0.86%
- 3.80 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.19%
- 1.80 lbs. 3, 6-dichloro-o-anisic acid equivalent per ton or 0.09%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

28-2-8 GUARANTEED ANALYSIS	
Total Nitrogen (N)	28.0%
0.8% Ammoniacal Nitrogen	
19.9% Urea Nitrogen	
4.3% Slowly Available Water Soluble Nitrogen*	
3.0% Water Insoluble Nitrogen	
Available Phosphate(P₂O₅)	2.0%
Soluble Potash (K₂O)	8.0%
Iron (Fe)	1.0%
Primary plant foods derived from: methylene urea, muriate of potash, diammonium phosphate, and urea.	
Secondary plant foods from: ferric oxy sulfate	
*4.3% slowly available water soluble nitrogen from methylenediurea and dimethylenetriurea.	

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements

NET WEIGHT: 20 POUNDS

589 AP092509
 EPA REG. NO. 2217-561
 EPA EST. NO. 2217-KS-1
 MANUFACTURED BY:



Telephone 800-821-7925

ACCEPTED
with COMMENTS
In EPA Letter Dated:
OCT 8 2009
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.

2217-561



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

Sublabel 1: Precautionary Statements and Directions for Use for outdoor domestic use pattern

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Harmful if adsorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminate clothing before reuse.

First Aid	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

OTHER USE PRECAUTIONS:

- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.

4
17

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS:

Do not use on dichondra, carpetgrass or St. Augustinegrass.

Do not use this product where desirable clovers are present. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

INSTRUCTIONS:

Weed 'N Feed 6 is a Trimec® complex, a combination of three proven weed killers -- 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 28-2-8 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second application may be made in 30 days.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (1.5 lb 2,4-D ae, 0.33 lb MCP-p ae and 0.16 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (3.0 lb 2,4-D ae, 0.66 lb MCP-p ae and 0.32 lb dicamba ae per acre per year).

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

WHAT TO EXPECT:

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application. [Refer to table for appropriate bag sizes.]

- ◆ For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.

5
17

- ◆ For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
.....	
Scotts Drop.....	5
Scotts SpeedyGreen® Rotary.....	3½
GreenView Drop (Spred II).....	4
Central Drop (#205).....	5½
Sears Drop (#19810).....	4
Ortho Drop (#5000).....	5
Precision Drop (Red Devil).....	4

- ◆ For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- ◆ For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- ◆ For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	3.0	4,000
	3.2	3,800
	3.4	-----
	3.6	-----
	4.0	-----
13	3.0	4,300
	3.2	4,100
	3.4	3,800
	3.6	3,600
	4.0	3,250
14	3.0	4,700
	3.2	4,400
	3.4	4,100
	3.6	3,900
	4.0	3,500
15	3.0	5,000
	3.2	4,700
	3.4	4,400
	3.6	4,200
	4.0	3,750
16	3.0	5,300
	3.2	5,000
	3.4	4,700
	3.6	4,400
	4.0	4,000

6
17

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
17	3.0	5,700
	3.2	5,300
	3.4	5,000
	3.6	4,700
	4.0	4,250
18	3.0	6,000
	3.2	5,600
	3.4	5,300
	3.6	5,000
	4.0	4,500
24	3.0	8,000
	3.2	7,500
	3.4	7,100
	3.6	6,700
	4.0	6,000
25	3.0	8,300
	3.2	7,800
	3.4	7,400
	3.6	6,900
	4.0	6,250
26	3.0	8,700
	3.2	8,100
	3.4	7,600
	3.6	7,200
	4.0	6,500
28	3.0	9,300
	3.2	8,800
	3.4	8,200
	3.6	7,800
	4.0	7,000
36	3.0	12,000
	3.2	11,300
	3.4	10,500
	3.6	10,000
	4.0	9,000
37	3.0	12,300
	3.2	11,600
	3.4	10,900
	3.6	10,300
	4.0	9,250
39	3.0	13,000
	3.2	12,200
	3.4	11,500
	3.6	10,800
	4.0	9,750
42	3.0	14,000
	3.2	13,100
	3.4	12,400
	3.6	11,700
	4.0	10,500

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

7
17

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. *[Graphics Optional]*

SINGLE PASS DISTRIBUTION:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage *[Graphics Optional]*. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

Advisory For Fertilizers Containing Iron:

Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

Fertilizer Labeling Requirements:

- A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000 square feet each growing season. When applied as directed this product supplies (1.1) pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local agricultural extension agent for the proper amount of nitrogen to be applied in your area.
- Fertilizer Labeling Requirements by Association of American Plant Food Control Officials (AAPFCO): Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metal.htm>.

Referrals to other products used sequentially:

- For best results, apply (brand name) products.
 - Apply using the 4-Step Program
 - **4-Step Program:**
 - STEP 1:** Menards Premium Crabgrass Preventor - Apply March - early May - Prevents crabgrass, foxtail, goosegrass, *poa annua* and other annual weed grasses. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
 - STEP 2:** Menards Premium Weed & Feed - Apply May - June - Kills dandelion, clover, plantain and over XXX other broadleaf weeds. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
 - STEP 3:** Menards Premium Lawn Food - Apply June - August - Nourishes your lawn throughout the summer months with extended feeding of Nitrogen. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
 - STEP 4:** Menards Premium Winterizer & Weed Control - Apply late August - mid October - Kills dandelion, clover, plantain and over XXX other broadleaf weeds. - Maximizes root building. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening.
- OR**
- STEP 4:** Menards Premium Starter & Winterizer - Apply late August - November - Maximizes root building. - Nutralene feeds your turfgrass for up to 2 months with long lasting deep greening. - Use with new seeding and sodding of turfgrass.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order of Injunctive Relief in Washington Toxics Coalition. et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to www.epa.gov/espp/wtcl/.

- BROADLEAF WEEDS -

Annual yellow sweetclover	Cockle	Mustard	Spiny amaranth
Aster	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Austrian fieldcress	Common mullein	Narrowleaf vetch	Spiny sowthistle
Bedstraw	Creeping jenny	Nettle	Spotted catsear
Beggarticks	Cudweed	Orange hawkweed	Spotted spurge
Betony, Florida	Curly dock	Oriental cocklebur	Spurweed
Bindweed, field	Daisy, English	Oxalis	Stinging nettle
Bird vetch	Daisy fleabane	Parsley-piert	Strawberry, India mock
Bitter wintercress	Daisy, oxeye	Parsnip	Tall nettle
Bittercress, hairy	Dandelion	Pearlwort	Tall vervain
Bitterweed	Dichondra	Pennycress	Tansy ragwort
Black-eyed Susan	Dogfennel	Pepperweed	Tansy mustard
Black medic	Dollarweed	Pigweed	Tanweed
Black mustard	False dandelion	Pineywoods bedstraw	Thistle
Blackseed plantain	Falseflax	Plains <i>coreopsis</i>	Trailing crownvetch
Blessed thistle	False sunflower	Plantain	Tumble mustard
Bloodflower milkweed	Fiddleneck	Poison ivy	Tumble pigweed
Blue lettuce	Florida pusley	Poison oak	Velvetleaf
Blue vervain	Frenchweed	Pokeweed	Venice mallow
Bracted plantain	Galinsoga	Poorjoe	Virginia buttonweed
Bristly oxtongue	Goldenrod	Prairie sunflower	Virginia creeper
Broadleaf dock	Ground ivy	Prickly lettuce	Virginia pepperweed
Broadleaf plantain	Gumweed	Prickly sida	Wavyleaf bullthistle
Broomweed	Hairy fleabane	Prostrate knotweed	Western clematis
Buckhorn	Hawkweed	Prostrate pigweed	Western salsify
Buckhorn plantain	Healall	Prostrate spurge	White mustard
Bulbous buttercup	Heartleaf drymary	Prostrate vervain	Wild aster
Bull thistle	Heathaster	Puncturevine	Wild buckwheat
Bullnettle	Hedge bindweed	Purslane, common	Wild carrot
Burclover	Hedge mustard	Ragweed	Wild four-o'clock
Burdock	Henbit	Red sorrel	Wild garlic
Burning nettle	Hoary cress	Redroot pigweed	Wild geranium
Burweed	Hoary plantain	Redstem filaree	Wild lettuce
Buttercup	Hoary vervain	Rough cinquefoil	Wild marigold
Buttonweed	Horsenettle	Rough fleabane	Wild mustard
Canada thistle	Jimsonweed	Russian pigweed	Wild onion
Carolina geranium	Knawel	Russian thistle	Wild parsnip
Carpetweed	Knotweed	Scarlet pimpernel	Wild radish
Catchweed bedstraw	Kochia	Scotch thistle	Wild rape
Catnip	Lambsquarters	Sheep sorrel	Wild strawberry
Catsear	Lespedeza	Shepherdspurse	Wild sweet potato
Chickweed, common	Mallow	Slender plantain	Wild vetch
Chickweed, mouseear	Matchweed	Smallflower galinsoga	Woodsorrel
Chicory	Mexicanweed	Smooth dock	Woolly croton
Cinquefoil	Milk vetch	Smooth pigweed	Woolly morningglory
Clover, crimson	Morningglory	Sorrel	Woolly plantain
Clover, hop	Mouseear hawkweed	Sowthistle	Wormseed
Clover, red	Mugwort	Spanishneedles	Yarrow
Clover, strawberry	Musk thistle	Speedwell	Yellow rocket
Clover, sweet			Yellowflower pepperweed
Clover, white			and other broadleaf weeds

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

9
17

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

10
17



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

Sublabel 2: Precautionary Statements and Directions for Use for occupational users

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves made of any waterproof material.

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

11
17

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

OTHER USE PRECAUTIONS:

- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WHERE TO USE:

Weed-'N-Feed 6 is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

Weed-'N-Feed 6 is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Weed-'N-Feed 6 may be applied to newly seeded cool season turfgrass. Applications are suitable only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

Weed-'N-Feed 6 may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (½ of the use rate for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area.

12
17

Do not use this product on St. Augustinegrass, dichondra, carpetgrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescue spp., perennial ryegrass, bermudagrass, and zoysiagrass:

Apply 130-174 pounds of Weed-'N-Feed 6 per acre of established turfgrass. Or the equivalent application rates are 3.0—4.0 pounds of product per 1,000 square feet of turfgrass.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application (1.5 lb 2,4-D ae, 0.33 lb MCP-P ae and 0.16 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. (3.0 lb 2,4-D ae, 0.66 lb MCP-P ae and 0.32 lb dicamba ae per acre per year).

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications is 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

APPLICATION SUGGESTIONS:

1. Spring and fall applications are suggested for the best control. However, Weed-'N-Feed 6 can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (> ½ inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<¼ inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application. [Refer to table for appropriate bag sizes.]

- ◆ For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- ◆ For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
Scotts Drop	5
Scotts SpeedyGreen® Rotary	3½
GreenView Drop (Spred II)	4
Central Drop (#205)	5½
Sears Drop (#19810)	4
Ortho Drop (#5000)	5
Precision Drop (Red Devil)	4

13
17

- ◆ For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- ◆ For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- ◆ For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, equipment settings of drop spreaders, rotary spreaders, and other granular applicators may be presented on the container labels.

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	3.0	4,000
	3.2	3,800
	3.4	-----
	3.6	-----
	4.0	-----
13	3.0	4,300
	3.2	4,100
	3.4	3,800
	3.6	3,600
	4.0	3,250
14	3.0	4,700
	3.2	4,400
	3.4	4,100
	3.6	3,900
	4.0	3,500
15	3.0	5,000
	3.2	4,700
	3.4	4,400
	3.6	4,200
	4.0	3,750
16	3.0	5,300
	3.2	5,000
	3.4	4,700
	3.6	4,400
	4.0	4,000
17	3.0	5,700
	3.2	5,300
	3.4	5,000
	3.6	4,700
	4.0	4,250
18	3.0	6,000
	3.2	5,600
	3.4	5,300
	3.6	5,000
	4.0	4,500
24	3.0	8,000
	3.2	7,500
	3.4	7,100
	3.6	6,700
	4.0	6,000
25	3.0	8,300
	3.2	7,800
	3.4	7,400
	3.6	6,900

14
17

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
	4.0	6,250
26	3.0	8,700
	3.2	8,100
	3.4	7,600
	3.6	7,200
	4.0	6,500
28	3.0	9,300
	3.2	8,800
	3.4	8,200
	3.6	7,800
	4.0	7,000
36	3.0	12,000
	3.2	11,300
	3.4	10,500
	3.6	10,000
	4.0	9,000
37	3.0	12,300
	3.2	11,600
	3.4	10,900
	3.6	10,300
	4.0	9,250
39	3.0	13,000
	3.2	12,200
	3.4	11,500
	3.6	10,800
	4.0	9,750
40	3.0	13,300
	3.2	12,500
	3.4	11,800
	3.6	11,100
	4.0	10,000
42	3.0	14,000
	3.2	13,100
	3.4	12,400
	3.6	11,700
	4.0	10,500
45	3.0	15,000
	3.2	14,100
	3.4	13,200
	3.6	12,500
	4.0	11,300
50	3.0	16,700
	3.2	15,600
	3.4	14,700
	3.6	13,900
	4.0	12,500

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

15
17

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. *[Graphics Optional]*

SINGLE PASS DISTRIBUTION:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage *[Graphics Optional]*. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

Advisory For Fertilizers Containing Iron:

Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

Fertilizer Labeling Requirements:

- A lawn will typically utilize 1 to 4 pounds of nitrogen per 1,000 square feet each growing season. When applied as directed this product supplies (1.1) pounds of nitrogen per 1,000 square feet with each application. Use this product in conjunction with an additional fertilization program to provide at least 1 pound of nitrogen per 1,000 square feet each growing season. Consult your local agricultural extension agent for the proper amount of nitrogen to be applied in your area.
- Fertilizer Labeling Requirements by Association of American Plant Food Control Officials (AAPFCO): Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metal.htm>.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order of Injunctive Relief in Washington Toxics Coalition. et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to www.epa.gov/espp/wtc/.

WEEDS CONTROLLED:

Dandelion, chickweed, plantain, and clover are controlled effectively with Weed-'N-Feed 6. A partial list of susceptible weeds is shown below.

- BROADLEAF WEEDS -			
Annual yellow sweetclover	Cockle	Mustard	Spiny amaranth
Aster	Cocklebur	Narrowleaf plantain	Spiny cocklebur
Austrian fieldcress	Common mullein	Narrowleaf vetch	Spiny sowthistle
Bedstraw	Creeping jenny	Nettle	Spotted catsear
Beggarticks	Cudweed	Orange hawkweed	Spotted spurge
Betony, Florida	Curly dock	Oriental cocklebur	Spurweed
Bindweed, field	Daisy, English	Oxalis	Stinging nettle
Bird vetch	Daisy fleabane	Parsley-piert	Strawberry, India mock
Bitter wintercress	Daisy, oxeye	Parsnip	Tall nettle
Bittercress, hairy	Dandelion	Pearlwort	Tall vervain
Bitterweed	Dichondra	Pennycress	Tansy ragwort
Black-eyed Susan	Dogfennel	Pepperweed	Tansy mustard
Black medic	Dollarweed	Pigweed	Tanweed
Black mustard	False dandelion	Pineywoods bedstraw	Thistle
Blackseed plantain	Falseflax	Plains <i>coreopsis</i>	Trailing crownvetch
Blessed thistle	False sunflower	Plantain	Tumble mustard
Bloodflower milkweed	Fiddleneck	Poison ivy	Tumble pigweed
Blue lettuce	Florida pusley	Poison oak	Velvetleaf
Blue vervain	Frenchweed	Pokeweed	Venice mallow
Bracted plantain	Galinsoga	Poorjoe	Virginia buttonweed
Bristly oxtongue	Goldenrod	Prairie sunflower	Virginia creeper

16
17

- BROADLEAF WEEDS -

Broadleaf dock	Ground ivy	Prickly lettuce	Virginia pepperweed
Broadleaf plantain	Gumweed	Prickly sida	Wavyleaf bullthistle
Broomweed	Hairy fleabane	Prostrate knotweed	Western clematis
Buckhorn	Hawkweed	Prostrate pigweed	Western salsify
Buckhorn plantain	Healall	Prostrate spurge	White mustard
Bulbous buttercup	Heartleaf drymary	Prostrate vervain	Wild aster
Bull thistle	Heathaster	Puncturevine	Wild buckwheat
Bullnettle	Hedge bindweed	Purslane, common	Wild carrot
Burclover	Hedge mustard	Ragweed	Wild four-o'clock
Burdock	Henbit	Red sorrel	Wild garlic
Burning nettle	Hoary cress	Redroot pigweed	Wild geranium
Burweed	Hoary plantain	Redstem filaree	Wild lettuce
Buttercup	Hoary vervain	Rough cinquefoil	Wild marigold
Buttonweed	Horsenettle	Rough fleabane	Wild mustard
Canada thistle	Jimsonweed	Russian pigweed	Wild onion
Carolina geranium	Knawel	Russian thistle	Wild parsnip
Carpetweed	Knotweed	Scarlet pimpernel	Wild radish
Catchweed bedstraw	Kochia	Scotch thistle	Wild rape
Catnip	Lambsquarters	Sheep sorrel	Wild strawberry
Catsear	Lespedeza	Shepherdspurse	Wild sweet potato
Chickweed, common	Mallow	Slender plantain	Wild vetch
Chickweed, mouseear	Matchweed	Smallflower galinsoga	Woodsorrel
Chicory	Mexicanweed	Smooth dock	Woolly croton
Cinquefoil	Milk vetch	Smooth pigweed	Woolly morningglory
Clover, crimson	Morningglory	Sorrel	Woolly plantain
Clover, hop	Mouseear hawkweed	Sowthistle	Wormseed
Clover, red	Mugwort	Spanishneedles	Yarrow
Clover, strawberry	Musk thistle	Speedwell	Yellow rocket
Clover, sweet			Yellowflower pepperweed
Clover, white			and other broadleaf weeds

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

17
17

APPENDIX

Advertising claims that may be presented on the container label of supplemental registrants.

- Controlled Release Nitrogen Nutralene®
- Produce Superior Results...Every Time!
- Nutralene® is a controlled-release homogenous proven technology. It releases nitrogen gradually through hydrolysis and microbial action in the soil for sustained feeding.
- Nutralene® is unlike other nitrogen sources that depend on coatings, particle size, and temperature for their nitrogen release.
- The ultimate timed release fertilizer as used by professional golf courses.
- Use when weeds are actively growing.
- Controls Over 200 Broadleaf Weeds.
- A granular fertilizer containing Trimec® herbicide which controls: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and 200 other broadleaf weeds which are listed on the (back/side) panel.
- Kills Weeds
- Fall Lawn Food
- Kills Over 200 Weeds!
- Prepares Lawn For Winter
- 100% Satisfaction Guaranteed Or Your Money Back
- Calculating Total Lawn Area [graphic] Length x Width = Total Sq.Ft. then subtract Non-Lawn Areas (House, Garden, Driveway)
- Greenyard Professional (TM)
- Greenyard Professional Logo is a trademark of Growmark