



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 5, 2017

Mr. Eric D. Smith
Federal Registration Manager
PBI/Gordon Corporation
1217 W. 12th St.
Kansas City, MO 64101

Subject: Notification per PRN 98-10 – Adding marketing claims and other minor updates
Product Name: WEED-‘N-FEED 6
EPA Registration Number: 2217-561
Application Date: 01/13/2017
Decision Number: 525416

Dear Mr. Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records. Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact Kathryn Montague at 703-305-1243 or via email at montague.kathryn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

NOTIFICATION

2217-561

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

04/05/2017

WEED-'N-FEED 6

EPA Reg. No. 2217-561

Two sublabels represent the entire master label of WEED-'N-FEED 6. Please refer to the appropriate section of the labeling as shown as follows:

Section
Language Common to All Sublabels
Sublabel 1 – Precautionary Statements and Directions for Use for outdoor domestic use pattern
Sublabel 2 – Precautionary Statements and Directions for Use for occupational users
Appendix
Document Control Information

Language Common to All Sublabels

WEED-'N-FEED 6

EPA Reg. No. 2217-561

ACTIVE INGREDIENT:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	1.040%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.230%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.108%
OTHER INGREDIENTS:	98.622%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 17.20 lb 2, 4-dichlorophenoxyacetic acid equivalent per ton or 0.86%
- 3.80 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.19%
- 1.80 lb 3, 6-dichloro-o-anisic acid equivalent per ton or 0.09%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
-x.x% ammoniacal nitrogen	
-x.x% urea nitrogen*	
Available Phosphate (P2O5)	X%
Soluble Potash (K2O)	X%
Sulfur (S)	X%
-x.x% free sulfur (S)	



Iron (Fe).....	X%
Derived from: polymer coated sulfur coated urea, urea, ammonium phosphate, potassium chloride. *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen. This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid. OR See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

Net Contents: _____
EPA Est. No. _____

Company Name and Address:

PBI/Gordon Corporation
1217 West 12th Street
Kansas City, Missouri 64101



SUBLABEL 1

Precautionary Statements and Directions for Use for outdoor domestic use pattern

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Harmful if adsorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

First Aid	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

OTHER USE PRECAUTIONS:

- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.

- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.

USE PRECAUTIONS:

- Do not use on dichondra, carpetgrass or St. Augustinegrass.
- Do not use this product where desirable clovers are present. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

INSTRUCTIONS:

Weed 'N Feed 6 is a Trimec® complex, a combination of three proven weed killers -- 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 28-2-8 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second application may be made in 30 days.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.5 lb 2,4-D ae, 0.33 lb MCP-p ae and 0.16 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [3.0 lb 2,4-D ae, 0.66 lb MCP-p ae and 0.32 lb dicamba ae per acre per year].

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

WHAT TO EXPECT:

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application.

[Note to Supplemental Registrants: Refer to table for appropriate bag sizes.]

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

[Optional Spreader Charts]

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
Scotts Drop	5
Scotts SpeedyGreen® Rotary	3.5
GreenView Drop (Spred II)	4
Central Drop (#205)	5.5
Sears Drop (#19810)	4
Ortho Drop (#5000)	5
Precision Drop (Red Devil)	4

- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, the following language on equipment settings of drop spreaders, rotary spreaders, and other granular applications is optional – it may be presented on the container labels at your discretion]

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	3.0	4,000	25	3.0	8,300
	3.2	3,800		3.2	7,800
	3.4	-		3.4	7,400
	3.6	-		3.6	6,900
	4.0	-		4.0	6,250
13	3.0	4,300	26	3.0	8,700
	3.2	4,100		3.2	8,100
	3.4	3,800		3.4	7,600
	3.6	3,600		3.6	7,200
	4.0	3,250		4.0	6,500
14	3.0	4,700	28	3.0	9,300
	3.2	4,400		3.2	8,800
	3.4	4,100		3.4	8,200
	3.6	3,900		3.6	7,800
	4.0	3,500		4.0	7,000
15	3.0	5,000	36	3.0	12,000
	3.2	4,700		3.2	11,300
	3.4	4,400		3.4	10,500
	3.6	4,200		3.6	10,000
	4.0	3,750		4.0	9,000
16	3.0	5,300	37	3.0	12,300
	3.2	5,000		3.2	11,600
	3.4	4,700		3.4	10,900
	3.6	4,400		3.6	10,300
	4.0	4,000		4.0	9,250
17	3.0	5,700	39	3.0	13,000
	3.2	5,300		3.2	12,200
	3.4	5,000		3.4	11,500
	3.6	4,700		3.6	10,800
	4.0	4,250		4.0	9,750

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
18	3.0	6,000	42	3.0	14,000
	3.2	5,600		3.2	13,100
	3.4	5,300		3.4	12,400
	3.6	5,000		3.6	11,700
	4.0	4,500		4.0	10,500
24	3.0	8,000			
	3.2	7,500			
	3.4	7,100			
	3.6	6,700			
	4.0	6,000			

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]

SINGLE PASS DISTRIBUTION:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

[End optional language]

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

BROADLEAF WEEDS			
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	Elderberry	Plains <i>coreopsis</i>	Tumble mustard
Blessed thistle	False dandelion	Plantain	Tumble pigweed
Bloodflower milkweed	Falseflax	Poison ivy	Velvetleaf
Blue lettuce	False sunflower	Poison oak	Venice mallow
Blue vervain	Fiddleneck	Pokeweed	Virginia buttonweed
Bracted plantain	Florida pusley	Poorjoe	Virginia creeper
Bristly oxtongue	Frenchweed	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Galinsoga	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prickly sida	Western clematis
Broomweed	Ground ivy	Prostrate knotweed	Western salsify
Buckhorn	Gumweed	Prostrate pigweed	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate spurge	Wild aster
Bulbous buttercup	Hawkweed	Prostrate vervain	Wild buckwheat
Bull thistle	Healall	Puncturevine	Wild carrot

Bullnettle	Heartleaf drymary	Purslane, common	Wild four-o'clock
Burclover	Heathaster	Ragweed	Wild garlic
Burdock	Hedge bindweed	Red sorrel	Wild geranium
Burning nettle	Hedge mustard	Redroot pigweed	Wild lettuce
Burweed	Hemp	Redstem filaree	Wild marigold
Buttercup	Henbit	Rough cinquefoil	Wild mustard
Buttonweed	Hoary cress	Rough fleabane	Wild onion
Canada thistle	Hoary plantain	Russian pigweed	Wild parsnip
Carolina geranium	Hoary vervain	Russian thistle	Wild radish
Carpetweed	Horsenettle	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Jimsonweed	Scotch thistle	Wild strawberry
Catnip	Knawel	Sheep sorrel	Wild sweet potato
Catsear	Knotweed	Shepherdspurse	Wild vetch
Chickweed, common	Kochia	Slender plantain	Woodsorrel
Chickweed, mouseear	Lambsquarters	Smallflower galinsoga	Woolly croton
Chicory	Lespedeza	Smooth dock	Woolly morningglory
Cinquefoil	Mallow	Smooth pigweed	Woolly plantain
Clover, crimson	Matchweed	Sorrel	Wormseed
Clover, hop	Mexicanweed	Sowthistle	Yarrow
Clover, red	Milk vetch	Spanishneedles	Yellow rocket
Clover, strawberry	Morningglory	Speedwell	Yellowflower pepperweed
Clover, sweet	Mouseear hawkweed	Spiny amaranth	
	Mugwort		
	Musk thistle		

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

[The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.]

OR

[FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.]

SUBLABEL 2

Precautionary Statements and Directions for Use for occupational users

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves made of any waterproof material.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

<h5>User Safety Recommendations</h5> <ul style="list-style-type: none">• Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.• Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.• Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
--

<h5>First Aid</h5>	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND USE RESTRICTIONS:

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

OTHER USE PRECAUTIONS:

- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product on St. Augustinegrass, dichondra, carpetgrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

WHERE TO USE:

Weed-'N-Feed 6 is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

Weed-'N-Feed 6 is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Weed-'N-Feed 6 may be applied to newly seeded cool season turfgrass. Applications are suitable only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

Weed-'N-Feed 6 may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the use rate for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area.

~~Do not use this product on St. Augustinegrass, dichondra, carpetgrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.~~

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescue spp., perennial ryegrass, bermudagrass, and zoysiagrass:

Apply 130 to 174 pounds of Weed-'N-Feed 6 per acre of established turfgrass. Or the equivalent application rates are 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.5 lb 2,4-D ae, 0.33 lb MCPP-p ae and 0.16 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [3.0 lb 2,4-D ae, 0.66 lb MCPP-p ae and 0.32 lb dicamba ae per acre per year].

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications is 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

APPLICATION SUGGESTIONS:

1. Spring and fall applications are suggested for the best control. However, Weed-'N-Feed 6 can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application.

[Note to Supplemental Registrants: Refer to table for appropriate bag sizes.]

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.

[Optional Spreader Charts]

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
Scotts Drop	5
Scotts SpeedyGreen® Rotary	3.5
GreenView Drop (Spred II)	4
Central Drop (#205)	5.5
Sears Drop (#19810)	4
Ortho Drop (#5000)	5
Precision Drop (Red Devil)	4

- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, the following language on equipment settings of drop spreaders, rotary spreaders, and other granular applications is optional – it may be presented on the container labels at your discretion]

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	3.0	4,000	26	3.0	8,700
	3.2	3,800		3.2	8,100
	3.4	-		3.4	7,600
	3.6	-		3.6	7,200
	4.0	-		4.0	6,500
13	3.0	4,300	28	3.0	9,300
	3.2	4,100		3.2	8,800
	3.4	3,800		3.4	8,200
	3.6	3,600		3.6	7,800
	4.0	3,250		4.0	7,000
14	3.0	4,700	36	3.0	12,000
	3.2	4,400		3.2	11,300
	3.4	4,100		3.4	10,500
	3.6	3,900		3.6	10,000
	4.0	3,500		4.0	9,000
15	3.0	5,000	37	3.0	12,300
	3.2	4,700		3.2	11,600
	3.4	4,400		3.4	10,900
	3.6	4,200		3.6	10,300
	4.0	3,750		4.0	9,250
16	3.0	5,300	39	3.0	13,000
	3.2	5,000		3.2	12,200
	3.4	4,700		3.4	11,500
	3.6	4,400		3.6	10,800
	4.0	4,000		4.0	9,750
17	3.0	5,700	40	3.0	13,300
	3.2	5,300		3.2	12,500
	3.4	5,000		3.4	11,800
	3.6	4,700		3.6	11,100
	4.0	4,250		4.0	10,000
18	3.0	6,000	42	3.0	14,000
	3.2	5,600		3.2	13,100
	3.4	5,300		3.4	12,400
	3.6	5,000		3.6	11,700
	4.0	4,500		4.0	10,500
24	3.0	8,000	45	3.0	15,000
	3.2	7,500		3.2	14,100
	3.4	7,100		3.4	13,200
	3.6	6,700		3.6	12,500
	4.0	6,000		4.0	11,300
25	3.0	8,300	50	3.0	16,700
	3.2	7,800		3.2	15,600
	3.4	7,400		3.4	14,700
	3.6	6,900		3.6	13,900
	4.0	6,250		4.0	12,500

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]

SINGLE PASS DISTRIBUTION:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to ensure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

[End optional language]

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

WEEDS CONTROLLED:

Dandelion, chickweed, plantain, and clover are controlled effectively with Weed-'N-Feed 6. A list of susceptible weeds is shown below.

BROADLEAF WEEDS			
Annual yellow sweetclover	Clover, white	Mustard	Spiny cocklebur
Aster	Cockle	Narrowleaf plantain	Spiny sowthistle
Austrian fieldcress	Cocklebur	Narrowleaf vetch	Spotted catsear
Bedstraw	Common mullein	Nettle	Spotted spurge
Beggarticks	Creeping jenny	Orange hawkweed	Spurweed
Betony, Florida	Cudweed	Oriental cocklebur	Stinging nettle
Bindweed, field	Curly dock	Oxalis	Strawberry, India mock
Bird vetch	Daisy, English	Parsley-piert	Tall nettle
Bitter wintercress	Daisy fleabane	Parsnip	Tall vervain
Bittercress, hairy	Daisy, oxeye	Pearlwort	Tansy ragwort
Bitterweed	Dandelion	Pennycress	Tansy mustard
Black-eyed Susan	Dichondra	Pepperweed	Tanweed
Black medic	Dogfennel	Pigweed	Thistle
Black mustard	Dollarweed	Pineywoods bedstraw	Trailing crownvetch
Blackseed plantain	Elderberry	Plains <i>coreopsis</i>	Tumble mustard
Blessed thistle	False dandelion	Plantain	Tumble pigweed
Bloodflower milkweed	Falseflax	Poison ivy	Velvetleaf
Blue lettuce	False sunflower	Poison oak	Venice mallow
Blue vervain	Fiddleneck	Pokeweed	Virginia buttonweed
Bracted plantain	Florida pusley	Poorjoe	Virginia creeper
Bristly oxtongue	Frenchweed	Prairie sunflower	Virginia pepperweed
Broadleaf dock	Galinsoga	Prickly lettuce	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prickly sida	Western clematis
Broomweed	Ground ivy	Prostrate knotweed	Western salsify
Buckhorn	Gumweed	Prostrate pigweed	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate spurge	Wild aster
Bulbous buttercup	Hawkweed	Prostrate vervain	Wild buckwheat
Bull thistle	Healall	Puncturevine	Wild carrot
Bullnettle	Heartleaf drymary	Purslane, common	Wild four-o'clock
Burclover	Heathaster	Ragweed	Wild garlic
Burdock	Hedge bindweed	Red sorrel	Wild geranium
Burning nettle	Hedge mustard	Redroot pigweed	Wild lettuce
Burweed	Hemp	Redstem filaree	Wild marigold
Buttercup	Henbit	Rough cinquefoil	Wild mustard
Buttonweed	Hoary cress	Rough fleabane	Wild onion
Canada thistle	Hoary plantain	Russian pigweed	Wild parsnip
Carolina geranium	Hoary vervain	Russian thistle	Wild radish
Carpetweed	Horsenettle	Scarlet pimpernel	Wild rape
Catchweed bedstraw	Jimsonweed	Scotch thistle	Wild strawberry
Catnip	Knawel	Sheep sorrel	Wild sweet potato
Catsear	Knotweed	Shepherdspurse	Wild vetch
Chickweed, common	Kochia	Slender plantain	Woodsorrel
Chickweed, mouseear	Lambsquarters	Smallflower galinsoga	Woolly croton
Chicory	Lespedeza	Smooth dock	Woolly morningglory
Cinquefoil	Mallow	Smooth pigweed	Woolly plantain

Clover, crimson Clover, hop Clover, red Clover, strawberry Clover, sweet	Matchweed Mexicanweed Milk vetch Morningglory Mouseear hawkweed Mugwort Musk thistle	Sorrel Sowthistle Spanishneedles Speedwell Spiny amaranth	Wormseed Yarrow Yellow rocket Yellowflower pepperweed
--	--	---	--

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

[The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.]

OR

[FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND

DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.]

APPENDIX

1. Statements which may appear on different label components depending on packaging configuration.

- See next panel for additional Precautionary Statements and First Aid
- Net Weight _____
- EPA Est. No. _____

21. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

Herbicide Benefit Claims:

- Kills Weeds
- Use when weeds are actively growing.
- Controls Over 200 Broadleaf Weeds
- **Kills over 200 Weeds!**
- Controls dandelion, clover, & over [XXX] other broadleaf weeds
- Controls chickweed, dandelion, knotweed, plantain, henbit, spurge and many other weeds as listed
- **Contains Trimec Herbicide**

Fertilizer Benefit Claims:

- Controlled Release Nitrogen Nutralene®
- Controlled release technology feeds 2+ months
- This product contains polymer-coated Enhanced Efficiency Fertilizer technology. As your lawn demands nutrients, nitrogen is released slowly into the soil for plant root uptake. This controlled-release approach eliminates surge growth and provides predictable feeding for over two months.
- All [Brand] fertilizer products contain Enhanced Efficiency Fertilizer which ensures consistent feeding, deep greening and promotes overall lawn health.
- The ultimate timed release fertilizer as used by professional golf courses.
- Fall Lawn Food
- Winterizer
- Prepares Lawn For Winter
- Iron enriched formula
- Quickens spring green-up

General Marketing Claims:

- A granular fertilizer containing Trimec® herbicide which controls: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and 200 other broadleaf weeds which are listed on the (back/side) panel.
- Produce Superior Results...Every Time!
- [100%] Satisfaction Guaranteed Or Your Money Back
- Calculating Total Lawn Area [graphic] Length x Width = Total Sq.Ft. then subtract Non-Lawn Areas (House, Garden, Driveway)
- Greenyard Professional (TM)
- Greenyard Professional Logo is a trademark of Growmark
- EC Grow www.ecgrowproducts.com TIPS – FAQ's – INFO
- 2nd application late spring/early summer [when used in 4 step program]
- Promotes thick & lush lawn
- Menard's Premium 100% Satisfaction Money Back Guarantee
- 4th application late summer/fall [when used in 4 step program]
- **Covers XXX sq. ft.**

Referrals to other products used sequentially:



- [Brand] Lawn Care Program:
 - Allow 8 weeks between applications
 - Menard’s Premium Crabgrass Preventer
 - 1st Application
 - Early Spring
 - Season-long crabgrass control
 - Menard’s Premium Weed & Feed
 - 2nd Application
 - Late Spring/Early Summer
 - Controls dandelions, clover and 200 other broadleaf weeds
 - Menard’s Premium Lawn Food
 - 3rd Application
 - Mid/Late Summer or Anytime
 - Promotes a lush and deep green lawn
 - Menard’s Premium Fall Lawn Food
 - 4th Application
 - Late Summer/Fall
 - Prepares lawn for winter
 - Quickens spring green-up

32. State mandated fertilizer labeling requirements:

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading “3.1 State Restrictions”]

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	

Available Phosphate (P2O5)	X%
Soluble Potash (K2O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Iron (Fe).....	X%
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride. *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen. This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- California, Oregon, Washington: “Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: [Note to reviewer: maximum number of applications in this section is dually based upon maximum ai load as well as fertilizer analysis effect on maximum nitrogen rates in different regions in Florida]:
 - In North Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In Central Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
 - In South Florida: Do not exceed X applications per year on bermudagrass. Do not exceed X applications per year on centipedegrass. Do not exceed X applications per year on St. Augustine grass. Do not exceed X applications per year on zoysiagrass.
- Idaho: “Apply only as directed”
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”
- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the “Soil Erosion and Sediment Control Act,” P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots.”
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: “In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency.”

DOCUMENT CONTROL INFORMATION

1. **Unique Label Identifier:** 002217-00561.20170113.notif-proposed-highlighted

2. **Reason for Issue:** ad claims, formatting, minor revisions

Internal Usage Notes
Associated Container Label: Supplemental Distributors