



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

April 29, 2022

Eric D. Smith  
Director, Regulatory Affairs  
PBI/Gordon Corporation  
22701 W 68<sup>th</sup> Terrace  
Shawnee, KS 66226

Subject: Notification per PRN 98-10 – Add optional label statements  
Product Name: WEED-‘N-FEED 6  
EPA Registration Number: 2217-561  
Application Date: August 11, 2021  
Decision Number: 581712

Dear Eric Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact please contact Derek Corbin at 202-566-2571 or by email at Corbin.Derek@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Mindy Ondish".

Mindy Ondish  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

# WEED-'N-FEED 6

## EPA Reg. No. 2217-561

### NOTIFICATION

2217-561

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

04/29/2022

#### ACTIVE INGREDIENTS:

|  |                |
|--|----------------|
| Dimethylamine salt of 2,4-dichlorophenoxyacetic acid .....                     | 1.040%         |
| Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid ..... | 0.230%         |
| Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) .....               | 0.108%         |
| OTHER INGREDIENTS: .....   | <u>98.622%</u> |
| TOTAL .....  | 100.000%       |

#### THIS PRODUCT CONTAINS:

- 17.20 lb 2, 4-dichlorophenoxyacetic acid equivalent per ton or 0.86%
- 3.80 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.19%
- 1.80 lb 3, 6-dichloro-o-anisic acid equivalent per ton or 0.09%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

### KEEP OUT OF REACH OF CHILDREN

### CAUTION

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

See attached booklet [inside pages] for complete Precautionary Statements and Directions for Use, including Storage and Disposal.

OR

See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

**Net Contents:** \_\_\_\_\_

**EPA Est. No.** \_\_\_\_\_

**Company Name and Address:**

PBI/Gordon Corporation

P.O. Box 860350

Shawnee, Kansas 66286



---

## SUBLABEL 1

### Precautionary Statements and Directions for Use for outdoor domestic use pattern

---

#### **PRECAUTIONARY STATEMENTS**

##### **Hazards to Humans & Domestic Animals**

**CAUTION:** Harmful if swallowed. Harmful if adsorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

| <b>First Aid</b>   |   |
|--|---|
| <b>If swallowed:</b>   | <ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul> |
| <b>If on skin or clothing:</b>   | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| <b>If in eyes:</b>   | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information. |   |

##### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

##### **USE RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product

landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.

- Do not use on dichondra, carpetgrass or St. Augustinegrass.
- Do not use this product where desirable clovers are present. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

#### **INSTRUCTIONS:**

Weed 'N Feed 6 is a Trimec® complex, a combination of three proven weed killers -- 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 28-2-8 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second application may be made in 30 days.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.5 lb 2,4-D ae, 0.33 lb MCP-p ae and 0.16 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [3.0 lb 2,4-D ae, 0.66 lb MCP-p ae and 0.32 lb dicamba ae per acre per year].

#### **WHEN TO APPLY:**

Apply when broadleaf weeds are actively growing in spring or early fall.

#### **WHAT TO EXPECT:**

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

#### **CALIBRATION INSTRUCTIONS:**

To provide proper distribution, calibrate spreader before application.

**[Note to Supplemental Registrants:** Refer to table for appropriate bag sizes.]

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

**[Note to Supplemental Registrants:** Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, the following language on equipment settings of drop spreaders, rotary spreaders, and other granular applications is optional – it may be presented on the container labels at your discretion]

| Bag Size, Lbs. | Application Rate, Lbs./1,000 Sq.Ft. | Treated Area for This Bag Size, Sq.Ft. | Bag Size, Lbs. | Application Rate, Lbs./1,000 Sq.Ft. | Treated Area for This Bag Size, Sq.Ft. |
|----------------|-------------------------------------|--|----------------|-------------------------------------|--|
| 12             | 3.0                                 | 4,000                                  | 25             | 3.0                                 | 8,300                                  |
|                | 3.2                                 | 3,800                                  |                | 3.2                                 | 7,800                                  |
|                | 3.4                                 | -                                      |                | 3.4                                 | 7,400                                  |
|                | 3.6                                 | -                                      |                | 3.6                                 | 6,900                                  |
|                | 4.0                                 | -                                      |                | 4.0                                 | 6,250                                  |
| 13             | 3.0                                 | 4,300                                  | 26             | 3.0                                 | 8,700                                  |
|                | 3.2                                 | 4,100                                  |                | 3.2                                 | 8,100                                  |
|                | 3.4                                 | 3,800                                  |                | 3.4                                 | 7,600                                  |
|                | 3.6                                 | 3,600                                  |                | 3.6                                 | 7,200                                  |
|                | 4.0                                 | 3,250                                  |                | 4.0                                 | 6,500                                  |
| 14             | 3.0                                 | 4,700                                  | 28             | 3.0                                 | 9,300                                  |
|                | 3.2                                 | 4,400                                  |                | 3.2                                 | 8,800                                  |
|                | 3.4                                 | 4,100                                  |                | 3.4                                 | 8,200                                  |
|                | 3.6                                 | 3,900                                  |                | 3.6                                 | 7,800                                  |
|                | 4.0                                 | 3,500                                  |                | 4.0                                 | 7,000                                  |
| 15             | 3.0                                 | 5,000                                  | 36             | 3.0                                 | 12,000                                 |
|                | 3.2                                 | 4,700                                  |                | 3.2                                 | 11,300                                 |
|                | 3.4                                 | 4,400                                  |                | 3.4                                 | 10,500                                 |
|                | 3.6                                 | 4,200                                  |                | 3.6                                 | 10,000                                 |
|                | 4.0                                 | 3,750                                  |                | 4.0                                 | 9,000                                  |
| 16             | 3.0                                 | 5,300                                  | 37             | 3.0                                 | 12,300                                 |
|                | 3.2                                 | 5,000                                  |                | 3.2                                 | 11,600                                 |
|                | 3.4                                 | 4,700                                  |                | 3.4                                 | 10,900                                 |
|                | 3.6                                 | 4,400                                  |                | 3.6                                 | 10,300                                 |
|                | 4.0                                 | 4,000                                  |                | 4.0                                 | 9,250                                  |
| 17             | 3.0                                 | 5,700                                  | 39             | 3.0                                 | 13,000                                 |
|                | 3.2                                 | 5,300                                  |                | 3.2                                 | 12,200                                 |
|                | 3.4                                 | 5,000                                  |                | 3.4                                 | 11,500                                 |
|                | 3.6                                 | 4,700                                  |                | 3.6                                 | 10,800                                 |
|                | 4.0                                 | 4,250                                  |                | 4.0                                 | 9,750                                  |
| 18             | 3.0                                 | 6,000                                  | 42             | 3.0                                 | 14,000                                 |
|                | 3.2                                 | 5,600                                  |                | 3.2                                 | 13,100                                 |
|                | 3.4                                 | 5,300                                  |                | 3.4                                 | 12,400                                 |
|                | 3.6                                 | 5,000                                  |                | 3.6                                 | 11,700                                 |
|                | 4.0                                 | 4,500                                  |                | 4.0                                 | 10,500                                 |
| 24             | 3.0                                 | 8,000                                  |                |                                     |  |
|                | 3.2                                 | 7,500                                  |                |                                     |  |
|                | 3.4                                 | 7,100                                  |                |                                     |  |
|                | 3.6                                 | 6,700                                  |                |                                     |  |
|                | 4.0                                 | 6,000                                  |                |                                     |  |

**Coverage** - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]

**SINGLE PASS DISTRIBUTION:**

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

[Optional Spreader Charts]

| <b>Spreader settings for 1.0 pound of product per 310 square feet</b> |     |
|---|-----|
| Scotts EasyGreen® Rotary .....  | 26  |
| Scotts Drop .....   | 5   |
| Scotts SpeedyGreen® Rotary .....                                      | 3.5 |
| GreenView Drop (Spred II) .....                                       | 4   |
| Central Drop (#205) .....   | 5.5 |
| Sears Drop (#19810) .....   | 4   |
| Ortho Drop (#5000) .....  | 5   |
| Precision Drop (Red Devil) .....                                      | 4   |

[End optional language]

**DISTRIBUTION AND SPREADER PATTERNS:**

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

**BROADLEAF WEEDS**

|                           |                     |                         |
|---------------------------|---------------------|-------------------------|
| Annual yellow sweetclover | Chickweed, mouseear | Hoary cress             |
| Aster                     | Chicory             | Hoary plantain          |
| Austrian fieldcress       | Cinquefoil          | Hoary vervain           |
| Bedstraw                  | Clover, crimson     | Horsenettle             |
| Beggarticks               | Clover, hop         | Jimsonweed              |
| Betony, Florida           | Clover, red         | Knawel                  |
| Bindweed, field           | Clover, strawberry  | Knotweed                |
| Bird vetch                | Clover, sweet       | Kochia                  |
| Bitter wintercress        | Clover, white       | Lambsquarters           |
| Bittercress, hairy        | Cockle              | Lespedeza               |
| Bitterweed                | Cocklebur           | Mallow                  |
| Black-eyed Susan          | Common mullein      | Matchweed               |
| Black medic               | Creeping jenny      | Mexicanweed             |
| Black mustard             | Cudweed             | Milk vetch              |
| Blackseed plantain        | Curly dock          | Morningglory            |
| Blessed thistle           | Daisy, English      | Mouseear hawkweed       |
| Bloodflower milkweed      | Daisy fleabane      | Mugwort                 |
| Blue lettuce              | Daisy, oxeye        | Musk thistle            |
| Blue vervain              | Dandelion           | Mustard                 |
| Bracted plantain          | Dichondra           | Narrowleaf plantain     |
| Bristly oxtongue          | Dogfennel           | Narrowleaf vetch        |
| Broadleaf dock            | Dollarweed          | Nettle                  |
| Broadleaf plantain        | Elderberry          | Orange hawkweed         |
| Broomweed                 | False dandelion     | Oriental cocklebur      |
| Buckhorn                  | Falseflax           | Oxalis                  |
| Buckhorn plantain         | False sunflower     | Parsley-piert           |
| Bulbous buttercup         | Fiddleneck          | Parsnip                 |
| Bull thistle              | Florida pusley      | Pearlwort               |
| Bullnettle                | Frenchweed          | Pennycress              |
| Burclover                 | Galinsoga           | Pepperweed              |
| Burdock                   | Goldenrod           | Pigweed                 |
| Burning nettle            | Ground ivy          | Pineywoods bedstraw     |
| Burweed                   | Gumweed             | Plains <i>coreopsis</i> |
| Buttercup                 | Hairy fleabane      | Plantain                |
| Buttonweed                | Hawkweed            | Poison ivy              |
| Canada thistle            | Healall             | Poison oak              |
| Carolina geranium         | Heartleaf drymary   | Pokeweed                |
| Carpetweed                | Heathaster          | Poorjoe                 |
| Catchweed bedstraw        | Hedge bindweed      | Prairie sunflower       |
| Catnip                    | Hedge mustard       | Prickly lettuce         |
| Catsear                   | Hemp                | Prickly sida            |
| Chickweed, common         | Henbit              | Prostrate knotweed      |

Prostrate pigweed  
Prostrate spurge  
Prostrate vervain  
Puncturevine  
Purslane, common  
Ragweed  
Red sorrel  
Redroot pigweed  
Redstem filaree  
Rough cinquefoil  
Rough fleabane  
Russian pigweed  
Russian thistle  
Scarlet pimpernel  
Scotch thistle  
Sheep sorrel  
Shepherdspurse  
Slender plantain  
Smallflower galinsoga  
Smooth dock  
Smooth pigweed  
Sorrel  
Sowthistle  
Spanishneedles  
Speedwell

Spiny amaranth  
Spiny cocklebur  
Spiny sowthistle  
Spotted catsear  
Spotted spurge  
Spurweed  
Stinging nettle  
Strawberry, India mock  
Tall nettle  
Tall vervain  
Tansy ragwort  
Tansy mustard  
Tanweed  
Thistle  
Trailing crownvetch  
Tumble mustard  
Tumble pigweed  
Velvetleaf  
Venice mallow  
Virginia buttonweed  
Virginia creeper  
Virginia pepperweed  
Wavyleaf bullthistle  
Western clematis  
Western salsify

White mustard  
Wild aster  
Wild buckwheat  
Wild carrot  
Wild four-o'clock  
Wild garlic  
Wild geranium  
Wild lettuce  
Wild marigold  
Wild mustard  
Wild onion  
Wild parsnip  
Wild radish  
Wild rape  
Wild strawberry  
Wild sweet potato  
Wild vetch  
Woodsorrel  
Woolly croton  
Woolly morningglory  
Woolly plantain  
Wormseed  
Yarrow  
Yellow rocket  
Yellowflower pepperweed

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

## LIMITED WARRANTY AND DISCLAIMER

[The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.]

OR

[FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.]



---

## SUBLABEL 2

### Precautionary Statements and Directions for Use for occupational users

---

#### **PRECAUTIONARY STATEMENTS**

##### **Hazards to Humans & Domestic Animals**

**CAUTION:** Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves made of any waterproof material.

##### **User Safety Requirements**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

|  |
|--|
| <h5><b>User Safety Recommendations</b></h5> <ul style="list-style-type: none"><li>• Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.</li><li>• Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li><li>• Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li></ul> |
|--|



|  |   |
|--|---|
| <h5><b>First Aid</b></h5>  |   |
| <b>If swallowed:</b>   | <ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul> |
| <b>If on skin or clothing:</b>   | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| <b>If in eyes:</b>   | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>   |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information. |   |








##### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## **USE RESTRICTIONS:**

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product on St. Augustinegrass, dichondra, carpetgrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

## **WHERE TO USE:**

Weed-'N-Feed 6 is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

Weed-'N-Feed 6 is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Weed-'N-Feed 6 may be applied to newly seeded cool season turfgrass. Applications are suitable only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

Weed-'N-Feed 6 may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the use rate for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area.

## **HOW MUCH TO APPLY:**

**For Kentucky bluegrass, fescue spp., perennial ryegrass, bermudagrass, and zoysiagrass:**

Apply 130 to 174 pounds of Weed-'N-Feed 6 per acre of established turfgrass. Or the equivalent application rates are 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.5 lb 2,4-D ae, 0.33 lb MCP-p ae and 0.16 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [3.0 lb 2,4-D ae, 0.66 lb MCP-p ae and 0.32 lb dicamba ae per acre per year].

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications is 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

**APPLICATION SUGGESTIONS:**

1. Spring and fall applications are suggested for the best control. However, Weed-N-Feed 6 can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

**CALIBRATION INSTRUCTIONS:**

To provide proper distribution, calibrate spreader before application.

**[Note to Supplemental Registrants:** Refer to table for appropriate bag sizes.]

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

**[Note to Supplemental Registrants:** Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, the following language on equipment settings of drop spreaders, rotary spreaders, and other granular applications is optional – it may be presented on the container labels at your discretion]

| Bag Size, Lbs. | Application Rate, Lbs./1,000 Sq.Ft. | Treated Area for This Bag Size, Sq.Ft. | Bag Size, Lbs. | Application Rate, Lbs./1,000 Sq.Ft. | Treated Area for This Bag Size, Sq.Ft. |
|----------------|-------------------------------------|--|----------------|-------------------------------------|--|
| 12             | 3.0                                 | 4,000                                  | 26             | 3.0                                 | 8,700                                  |
|                | 3.2                                 | 3,800                                  |                | 3.2                                 | 8,100                                  |
|                | 3.4                                 | -                                      |                | 3.4                                 | 7,600                                  |
|                | 3.6                                 | -                                      |                | 3.6                                 | 7,200                                  |
|                | 4.0                                 | -                                      |                | 4.0                                 | 6,500                                  |

| Bag Size, Lbs. | Application Rate, Lbs./1,000 Sq.Ft. | Treated Area for This Bag Size, Sq.Ft. | Bag Size, Lbs. | Application Rate, Lbs./1,000 Sq.Ft. | Treated Area for This Bag Size, Sq.Ft. |
|----------------|-------------------------------------|--|----------------|-------------------------------------|--|
| 13             | 3.0                                 | 4,300                                  | 28             | 3.0                                 | 9,300                                  |
|                | 3.2                                 | 4,100                                  |                | 3.2                                 | 8,800                                  |
|                | 3.4                                 | 3,800                                  |                | 3.4                                 | 8,200                                  |
|                | 3.6                                 | 3,600                                  |                | 3.6                                 | 7,800                                  |
|                | 4.0                                 | 3,250                                  |                | 4.0                                 | 7,000                                  |
| 14             | 3.0                                 | 4,700                                  | 36             | 3.0                                 | 12,000                                 |
|                | 3.2                                 | 4,400                                  |                | 3.2                                 | 11,300                                 |
|                | 3.4                                 | 4,100                                  |                | 3.4                                 | 10,500                                 |
|                | 3.6                                 | 3,900                                  |                | 3.6                                 | 10,000                                 |
|                | 4.0                                 | 3,500                                  |                | 4.0                                 | 9,000                                  |
| 15             | 3.0                                 | 5,000                                  | 37             | 3.0                                 | 12,300                                 |
|                | 3.2                                 | 4,700                                  |                | 3.2                                 | 11,600                                 |
|                | 3.4                                 | 4,400                                  |                | 3.4                                 | 10,900                                 |
|                | 3.6                                 | 4,200                                  |                | 3.6                                 | 10,300                                 |
|                | 4.0                                 | 3,750                                  |                | 4.0                                 | 9,250                                  |
| 16             | 3.0                                 | 5,300                                  | 39             | 3.0                                 | 13,000                                 |
|                | 3.2                                 | 5,000                                  |                | 3.2                                 | 12,200                                 |
|                | 3.4                                 | 4,700                                  |                | 3.4                                 | 11,500                                 |
|                | 3.6                                 | 4,400                                  |                | 3.6                                 | 10,800                                 |
|                | 4.0                                 | 4,000                                  |                | 4.0                                 | 9,750                                  |
| 17             | 3.0                                 | 5,700                                  | 40             | 3.0                                 | 13,300                                 |
|                | 3.2                                 | 5,300                                  |                | 3.2                                 | 12,500                                 |
|                | 3.4                                 | 5,000                                  |                | 3.4                                 | 11,800                                 |
|                | 3.6                                 | 4,700                                  |                | 3.6                                 | 11,100                                 |
|                | 4.0                                 | 4,250                                  |                | 4.0                                 | 10,000                                 |
| 18             | 3.0                                 | 6,000                                  | 42             | 3.0                                 | 14,000                                 |
|                | 3.2                                 | 5,600                                  |                | 3.2                                 | 13,100                                 |
|                | 3.4                                 | 5,300                                  |                | 3.4                                 | 12,400                                 |
|                | 3.6                                 | 5,000                                  |                | 3.6                                 | 11,700                                 |
|                | 4.0                                 | 4,500                                  |                | 4.0                                 | 10,500                                 |
| 24             | 3.0                                 | 8,000                                  | 45             | 3.0                                 | 15,000                                 |
|                | 3.2                                 | 7,500                                  |                | 3.2                                 | 14,100                                 |
|                | 3.4                                 | 7,100                                  |                | 3.4                                 | 13,200                                 |
|                | 3.6                                 | 6,700                                  |                | 3.6                                 | 12,500                                 |
|                | 4.0                                 | 6,000                                  |                | 4.0                                 | 11,300                                 |
| 25             | 3.0                                 | 8,300                                  | 50             | 3.0                                 | 16,700                                 |
|                | 3.2                                 | 7,800                                  |                | 3.2                                 | 15,600                                 |
|                | 3.4                                 | 7,400                                  |                | 3.4                                 | 14,700                                 |
|                | 3.6                                 | 6,900                                  |                | 3.6                                 | 13,900                                 |
|                | 4.0                                 | 6,250                                  |                | 4.0                                 | 12,500                                 |

**Coverage** - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]

**SINGLE PASS DISTRIBUTION:**

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to ensure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

[Optional Spreader Charts]

| Spreader settings for 1.0 pound of product per 310 square feet |     |
|--|-----|
| Scotts EasyGreen® Rotary .....                                 | 26  |
| Scotts Drop .....  | 5   |
| Scotts SpeedyGreen® Rotary .....                               | 3.5 |

|                                  |     |
|----------------------------------|-----|
| GreenView Drop (Spred II) .....  | 4   |
| Central Drop (#205) .....        | 5.5 |
| Sears Drop (#19810) .....        | 4   |
| Ortho Drop (#5000) .....         | 5   |
| Precision Drop (Red Devil) ..... | 4   |

**[End optional language]**

**DISTRIBUTION AND SPREADER PATTERNS:**

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

**BROADLEAF WEEDS CONTROLLED:**

Dandelion, chickweed, plantain, and clover are controlled effectively with Weed-'N-Feed 6. A list of susceptible weeds is shown below.

|                           |                    |                         |
|---------------------------|--------------------|-------------------------|
| Annual yellow sweetclover | Clover, crimson    | Knotweed                |
| Aster                     | Clover, hop        | Kochia                  |
| Austrian fieldcress       | Clover, red        | Lambsquarters           |
| Bedstraw                  | Clover, strawberry | Lespedeza               |
| Beggarticks               | Clover, sweet      | Mallow                  |
| Betony, Florida           | Clover, white      | Matchweed               |
| Bindweed, field           | Cockle             | Mexicanweed             |
| Bird vetch                | Cocklebur          | Milk vetch              |
| Bitter wintercress        | Common mullein     | Morningglory            |
| Bittercress, hairy        | Creeping jenny     | Mouseear hawkweed       |
| Bitterweed                | Cudweed            | Mugwort                 |
| Black-eyed Susan          | Curly dock         | Musk thistle            |
| Black medic               | Daisy, English     | Mustard                 |
| Black mustard             | Daisy fleabane     | Narrowleaf plantain     |
| Blackseed plantain        | Daisy, oxeye       | Narrowleaf vetch        |
| Blessed thistle           | Dandelion          | Nettle                  |
| Bloodflower milkweed      | Dichondra          | Orange hawkweed         |
| Blue lettuce              | Dogfennel          | Oriental cocklebur      |
| Blue vervain              | Dollarweed         | Oxalis                  |
| Bracted plantain          | Elderberry         | Parsley-piert           |
| Bristly oxtongue          | False dandelion    | Parsnip                 |
| Broadleaf dock            | Falseflax          | Pearlwort               |
| Broadleaf plantain        | False sunflower    | Pennycress              |
| Broomweed                 | Fiddleneck         | Pepperweed              |
| Buckhorn                  | Florida pusley     | Pigweed                 |
| Buckhorn plantain         | Frenchweed         | Pineywoods bedstraw     |
| Bulbous buttercup         | Galinsoga          | Plains <i>coreopsis</i> |
| Bull thistle              | Goldenrod          | Plantain                |
| Bullnettle                | Ground ivy         | Poison ivy              |
| Burclover                 | Gumweed            | Poison oak              |
| Burdock                   | Hairy fleabane     | Pokeweed                |
| Burning nettle            | Hawkweed           | Poorjoe                 |
| Burweed                   | Healall            | Prairie sunflower       |
| Buttercup                 | Heartleaf drymary  | Prickly lettuce         |
| Buttonweed                | Heathaster         | Prickly sida            |
| Canada thistle            | Hedge bindweed     | Prostrate knotweed      |
| Carolina geranium         | Hedge mustard      | Prostrate pigweed       |
| Carpetweed                | Hemp               | Prostrate spurge        |
| Catchweed bedstraw        | Henbit             | Prostrate vervain       |
| Catnip                    | Hoary cress        | Puncturevine            |
| Catsear                   | Hoary plantain     | Purslane, common        |
| Chickweed, common         | Hoary vervain      | Ragweed                 |
| Chickweed, mouseear       | Horsenettle        | Red sorrel              |
| Chicory                   | Jimsonweed         | Redroot pigweed         |
| Cinquefoil                | Knawel             | Redstem filaree         |

Rough cinquefoil  
Rough fleabane  
Russian pigweed  
Russian thistle  
Scarlet pimpernel  
Scotch thistle  
Sheep sorrel  
Shepherdspurse  
Slender plantain  
Smallflower galinsoga  
Smooth dock  
Smooth pigweed  
Sorrel  
Sowthistle  
Spanishneedles  
Speedwell  
Spiny amaranth  
Spiny cocklebur  
Spiny sowthistle  
Spotted catsear  
Spotted spurge  
Spurweed

Stinging nettle  
Strawberry, India mock  
Tall nettle  
Tall vervain  
Tansy ragwort  
Tansy mustard  
Tanweed  
Thistle  
Trailing crownvetch  
Tumble mustard  
Tumble pigweed  
Velvetleaf  
Venice mallow  
Virginia buttonweed  
Virginia creeper  
Virginia pepperweed  
Wavyleaf bullthistle  
Western clematis  
Western salsify  
White mustard  
Wild aster  
Wild buckwheat

Wild carrot  
Wild four-o'clock  
Wild garlic  
Wild geranium  
Wild lettuce  
Wild marigold  
Wild mustard  
Wild onion  
Wild parsnip  
Wild radish  
Wild rape  
Wild strawberry  
Wild sweet potato  
Wild vetch  
Woodsorrel  
Woolly croton  
Woolly morningglory  
Woolly plantain  
Wormseed  
Yarrow  
Yellow rocket  
Yellowflower pepperweed

## **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Nonrefillable container: Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## **LIMITED WARRANTY AND DISCLAIMER**

[The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.]

OR

[FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.]

## **APPENDIX**

### **1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:**

#### Herbicide Benefit Claims:

- Kills Weeds
- [Apply/Use] when weeds are actively growing.
- [Kills/Controls] Over 200 listed Broadleaf Weeds
- Kills over 200 listed Weeds!
- Controls dandelion, clover, & over [XXX] other broadleaf weeds as listed
- Controls chickweed, dandelion, knotweed, plantain, henbit, spurge and many other weeds as listed
- Controls [insert weed from weed list] [and other listed weeds]
- Contains Trimec Herbicide
- Contains 3 proven weed killers
- 3-way active ingredient for optimum control
- Kills listed weeds to the root
- [Kills][Controls] most common weeds as listed
- Kills listed weeds and thickens lawns
- Best results are obtained when weeds are growing actively in spring or early fall

#### Fertilizer Benefit Claims:

- Controlled Release Nitrogen Nutralene®
- Controlled release technology feeds 2+ months
- This product contains polymer-coated Enhanced Efficiency Fertilizer technology. As your lawn demands nutrients, nitrogen is released slowly into the soil for plant root uptake. This controlled-release approach eliminates surge growth and provides predictable feeding for over two months.
- All [Brand] fertilizer products contain Enhanced Efficiency Fertilizer which ensures consistent feeding, deep greening and promotes overall lawn health.
- The ultimate timed release fertilizer as used by professional golf courses.
- Fall Lawn Food
- Winterizer
- Prepares Lawn For Winter
- Iron enriched formula
- Quickens spring green-up

- Feeds And Thickens Your Lawn
- Kills listed weeds And Thickens [Your] Lawn
- Long-lasting feeding
- Long-lasting greening
- Promotes a rich green lawn
- Contains Iron for Deep Greening
- Promotes color and growth
- Fortified with milorganite [logo/graphic]
- Milorganite. For Better Results. Naturally.
- [Contains] Iron for extra greening
- Stimulates plant growth
- Contains non-burning, slow release nitrogen and iron for deep greening
- Contains non-burning/slow-release nitrogen
- Improves turf quality
- Zero phosphorus
- Phosphorus Free
- [Contains] Controlled/Slow release nitrogen for extended feeding
- Contains [X%] Iron for deep greening
- Feeds Your Lawn for [over] [x] weeks/months
- Deep greening [formula]
- Non-staining [formula]
- Stimulates [rich/deep green] color and growth
- Replaces [Provides] Nutrients for a Thicker, Greener Lawn
- Quick greening/green-up and/then continues to feed your lawn
- Slow release nitrogen feeds your lawn [turfgrass] over [x] months/weeks.
- Allow [6/X] weeks between applications [Note to reviewer: X weeks will be greater than 4 to adhere to minimum retreatment interval of 30 days]
- Feeds And Thickens Your Lawn
- Promotes color and growth
- Lush Green Lawn
- Formulated to promote root building and winter hardiness

General Marketing Claims:

- A granular fertilizer containing Trimec® herbicide which controls: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and 200 other broadleaf weeds which are listed on the (back/side) panel.
- 2-In-1 Kills listed weeds and feed(s) your lawn
- Provides a full feeding of fertilizer and kills over listed 200 weeds
- Produce Superior Results...Every Time!
- [100%] Satisfaction Guaranteed Or Your Money Back
- [100%] Satisfaction Guarantee. If, after using this product, you are dissatisfied with the results, simply mail the sales receipt, a brief description of the problem and your contact information to the address listed below or contact Menards guest services for a full refund.
- Calculating Total Lawn Area [graphic] Length x Width = Total Sq.Ft. then subtract Non-Lawn Areas (House, Garden, Driveway)
- Greenyard Professional (TM)
- Greenyard Professional Logo is a trademark of Growmark
- EC Grow [www.ecgrowproducts.com](http://www.ecgrowproducts.com) TIPS – FAQ's – INFO
- 2<sup>nd</sup> application late spring/early summer [when used in 4 step program]
- Promotes thick & lush lawn
- Menard's Premium 100% Satisfaction Money Back Guarantee
- 4<sup>th</sup> application late summer/fall [when used in 4 step program]
- Covers XXX sq. ft.



- Apply When Weeds Are Actively Growing
- Apply in fall for a greener lawn next spring

Referrals to other products used sequentially:

**MENARDS® PREMIUM LAWN CARE PROGRAM**

| 1 <sup>ST</sup> APPLICATION   | 2 <sup>ND</sup> APPLICATION                              | 3 <sup>RD</sup> APPLICATION           | 4 <sup>TH</sup> APPLICATION                          |
|-------------------------------|--|---------------------------------------|--|
| <b>EARLY SPRING</b>           | <b>LATE SPRING/<br/>EARLY SUMMER</b>                     | <b>MID/LATE SUMMER<br/>OR ANYTIME</b> | <b>LATE SUMMER/FALL</b>                              |
| Season-Long Crabgrass Control | Controls dandelion, clover and 200 other broadleaf weeds | Promotes a lush and deep green lawn   | Prepares lawn for winter<br>Quickens spring green-up |

**ALLOW 8 WEEKS BETWEEN APPLICATIONS**

**1<sup>st</sup> APP** **CRABGRASS PREVENTER**  
APPLY EARLY SPRING  
with LAWN FOOD 26-0-3

---

**2<sup>nd</sup> APP** **WEED & FEED**  
APPLY LATE SPRING/  
EARLY SUMMER  
28-0-3

---

**3<sup>rd</sup> APP** **LAWN FOOD**  
APPLY MID/LATE SUMMER  
OR ANYTIME  
30-0-3

---

**4<sup>th</sup> APP** **FALL LAWN FOOD**  
APPLY LATE SUMMER/FALL  
WINTERIZER 22-0-10

**ALLOW 8 WEEKS BETWEEN APPLICATIONS**



- [Brand] Lawn Care Program:
  - Allow 8 weeks between applications
  - Menard's Premium Crabgrass Preventer
    - 1<sup>st</sup> Application
    - [Apply] Early Spring
    - Season-long crabgrass control
  - Menard's Premium Weed & Feed
    - 2<sup>nd</sup> Application
    - [Apply] Late Spring/Early Summer
    - Controls dandelions, clover and 200 other broadleaf weeds
  - Menard's Premium Lawn Food
    - 3<sup>rd</sup> Application
    - [Apply] Mid/Late Summer or Anytime
    - Promotes a lush and deep green lawn
  - Menard's Premium Fall Lawn Food
    - 4<sup>th</sup> Application
    - [Apply] Late Summer/Fall
    - Prepares lawn for winter
    - Quickens spring green-up
- [Brand] lawn care fertilizer program [insert graphic showing different application steps in the program with name]



- Lawncare [fertilizer] program [early spring, late spring/early summer, summer or anytime, late summer/fall] [insert graphic showing different application steps in the program with name]



- Application/App 1 [2][3][4]
- JRK Brand Lawn Care Program
- JRK Brand A Gertens® family of products.

**2. State mandated fertilizer labeling requirements:**

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading “3.1 State Restrictions”]

| GUARANTEED ANALYSIS XX-X-X  |     |
|---|-----|
| Total Nitrogen (N) .....  | XX% |
| x.x% ammoniacal nitrogen  |     |
| x.x% urea nitrogen*   |     |
| Available Phosphate (P2O5) .....  | X%  |
| Soluble Potash (K2O) .....  | X%  |
| Sulfur (S) .....  | X%  |
| x.x% free sulfur (S)  |     |
| Iron (Fe).....  | X%  |
| Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride.  |     |
| *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen.  |     |
| This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx |     |

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- California, Oregon, Washington: “[When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.] “Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: “Apply only to actively growing turf.”
- Florida: “Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided

the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”

- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the “Soil Erosion and Sediment Control Act,” P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots.”
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: “In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency.”
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]
- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet [and herbicide to control listed weeds]. Please consult your local extension office for questions regarding a regular fertilizer program [suitable for your geographic region].”

## **DOCUMENT CONTROL INFORMATION**

1. **Unique Label Identifier:** 002217-00561.20210811.notif-proposed-highlighted.doc

2. **Reason for Issue:** ad claims

| <b>Internal Usage Notes</b>                           |
|---|
| Associated Container Label: Supplemental Distributors |