



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 9, 2022

Eric D. Smith
VP of Regulatory, Formulation, and GLP
PBI/Gordon Corporation
22701 W 68th Terrace
Shawnee, KS 66226

Subject: Notification per PRN 98-10 – add optional spreader graphics, warranty statements, marketing claims, and other minor edits
Product Name: WEED-‘N-FEED 6
EPA Registration Number: 2217-561
Application Date: August 10, 2022
Decision Number: 586925

Dear Mr. Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped “NOTIFICATION” and placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, please contact me at ondish.mindy@epa.gov or at (202)566-2857.

Sincerely,

A handwritten signature in cursive script that reads "Mindy Ondish".

Mindy Ondish
Product Manager 23
Herbicide Branch
Registration Division (7505T)
Office of Pesticide Programs

NOTIFICATION

2217-561

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

09/09/2022

WEED-'N-FEED 6

EPA Reg. No. 2217-561

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	1.040%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.230%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.108%
OTHER INGREDIENTS:	98.622%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

- 17.20 lb 2, 4-dichlorophenoxyacetic acid equivalent per ton or 0.86%
- 3.80 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.19%
- 1.80 lb 3, 6-dichloro-o-anisic acid equivalent per ton or 0.09%

Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

See attached booklet [inside pages] for complete Precautionary Statements and Directions for Use, including Storage and Disposal.

OR

See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

Net Contents: _____

EPA Est. No. _____

Company Name and Address:

PBI/Gordon Corporation
P.O. Box 860350
Shawnee, Kansas 66286



SUBLABEL 1

Precautionary Statements and Directions for Use for outdoor domestic use pattern

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Harmful if adsorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

First Aid	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product

landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.

- Do not use on dichondra, carpetgrass or St. Augustinegrass.
- Do not use this product where desirable clovers are present. Do not apply in areas underlaid by roots of desirable trees or shrubs. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning.

INSTRUCTIONS:

Weed 'N Feed 6 is a Trimec® complex, a combination of three proven weed killers -- 2,4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a 28-2-8 fertilizer provides a supply of plant nutrients. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth.

Apply 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second application may be made in 30 days.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.5 lb 2,4-D ae, 0.33 lb MCP-p ae and 0.16 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [3.0 lb 2,4-D ae, 0.66 lb MCP-p ae and 0.32 lb dicamba ae per acre per year].

WHEN TO APPLY:

Apply when broadleaf weeds are actively growing in spring or early fall.

WHAT TO EXPECT:

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control. This product will not prevent weed seeds from producing new plants.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application.

[Note to Supplemental Registrants: Refer to table for appropriate bag sizes.]

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, the following language on equipment settings of drop spreaders, rotary spreaders, and other granular applications is optional – it may be presented on the container labels at your discretion]

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	3.0	4,000	25	3.0	8,300
	3.2	3,800		3.2	7,800
	3.4	-		3.4	7,400
	3.6	-		3.6	6,900
	4.0	-		4.0	6,250
13	3.0	4,300	26	3.0	8,700
	3.2	4,100		3.2	8,100
	3.4	3,800		3.4	7,600
	3.6	3,600		3.6	7,200
	4.0	3,250		4.0	6,500
14	3.0	4,700	28	3.0	9,300
	3.2	4,400		3.2	8,800
	3.4	4,100		3.4	8,200
	3.6	3,900		3.6	7,800
	4.0	3,500		4.0	7,000
15	3.0	5,000	36	3.0	12,000
	3.2	4,700		3.2	11,300
	3.4	4,400		3.4	10,500
	3.6	4,200		3.6	10,000
	4.0	3,750		4.0	9,000
16	3.0	5,300	37	3.0	12,300
	3.2	5,000		3.2	11,600
	3.4	4,700		3.4	10,900
	3.6	4,400		3.6	10,300
	4.0	4,000		4.0	9,250
17	3.0	5,700	39	3.0	13,000
	3.2	5,300		3.2	12,200
	3.4	5,000		3.4	11,500
	3.6	4,700		3.6	10,800
	4.0	4,250		4.0	9,750
18	3.0	6,000	42	3.0	14,000
	3.2	5,600		3.2	13,100
	3.4	5,300		3.4	12,400
	3.6	5,000		3.6	11,700
	4.0	4,500		4.0	10,500
24	3.0	8,000			
	3.2	7,500			
	3.4	7,100			
	3.6	6,700			
	4.0	6,000			

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]

SINGLE PASS DISTRIBUTION [SPREADING PATTERNS]:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

[Optional Spreader Charts]

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
Scotts Drop	5
Scotts SpeedyGreen® Rotary	3.5
GreenView Drop (Spred II)	4
Central Drop (#205)	5.5
Sears Drop (#19810)	4
Ortho Drop (#5000)	5
Precision Drop (Red Devil)	4

[End optional language]

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

BROADLEAF WEEDS

Annual yellow sweetclover	Chickweed, mouseear	Hoary cress
Aster	Chicory	Hoary plantain
Austrian fieldcress	Cinquefoil	Hoary vervain
Bedstraw	Clover, crimson	Horsenettle
Beggarticks	Clover, hop	Jimsonweed
Betony, Florida	Clover, red	Knawel
Bindweed, field	Clover, strawberry	Knotweed
Bird vetch	Clover, sweet	Kochia
Bitter wintercress	Clover, white	Lambsquarters
Bittercress, hairy	Cockle	Lespedeza
Bitterweed	Cocklebur	Mallow
Black-eyed Susan	Common mullein	Matchweed
Black medic	Creeping jenny	Mexicanweed
Black mustard	Cudweed	Milk vetch
Blackseed plantain	Curly dock	Morningglory
Blessed thistle	Daisy, English	Mouseear hawkweed
Bloodflower milkweed	Daisy fleabane	Mugwort
Blue lettuce	Daisy, oxeye	Musk thistle
Blue vervain	Dandelion	Mustard
Bracted plantain	Dichondra	Narrowleaf plantain
Bristly oxtongue	Dogfennel	Narrowleaf vetch
Broadleaf dock	Dollarweed	Nettle
Broadleaf plantain	Elderberry	Orange hawkweed
Broomweed	False dandelion	Oriental cocklebur
Buckhorn	Falseflax	Oxalis
Buckhorn plantain	False sunflower	Parsley-piert
Bulbous buttercup	Fiddleneck	Parsnip
Bull thistle	Florida pusley	Pearlwort
Bullnettle	Frenchweed	Pennycress
Burclover	Galinsoga	Pepperweed
Burdock	Goldenrod	Pigweed
Burning nettle	Ground ivy	Pineywoods bedstraw
Burweed	Gumweed	Plains <i>coreopsis</i>
Buttercup	Hairy fleabane	Plantain
Buttonweed	Hawkweed	Poison ivy
Canada thistle	Healall	Poison oak
Carolina geranium	Heartleaf drymary	Pokeweed
Carpetweed	Heathaster	Poorjoe
Catchweed bedstraw	Hedge bindweed	Prairie sunflower
Catnip	Hedge mustard	Prickly lettuce
Catsear	Hemp	Prickly sida
Chickweed, common	Henbit	Prostrate knotweed

Prostrate pigweed
Prostrate spurge
Prostrate vervain
Puncturevine
Purslane, common
Ragweed
Red sorrel
Redroot pigweed
Redstem filaree
Rough cinquefoil
Rough fleabane
Russian pigweed
Russian thistle
Scarlet pimpernel
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smooth dock
Smooth pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell

Spiny amaranth
Spiny cocklebur
Spiny sowthistle
Spotted catsear
Spotted spurge
Spurweed
Stinging nettle
Strawberry, India mock
Tall nettle
Tall vervain
Tansy ragwort
Tansy mustard
Tanweed
Thistle
Trailing crownvetch
Tumble mustard
Tumble pigweed
Velvetleaf
Venice mallow
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Wavyleaf bullthistle
Western clematis
Western salsify

White mustard
Wild aster
Wild buckwheat
Wild carrot
Wild four-o'clock
Wild garlic
Wild geranium
Wild lettuce
Wild marigold
Wild mustard
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild vetch
Woodsorrel
Woolly croton
Woolly morningglory
Woolly plantain
Wormseed
Yarrow
Yellow rocket
Yellowflower pepperweed

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or www.website.com for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

[The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.]

OR

[FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.]

SUBLABEL 2

Precautionary Statements and Directions for Use for occupational users

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves made of any waterproof material.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

<h5>User Safety Recommendations</h5> <ul style="list-style-type: none">• Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.• Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.• Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
--

<h5>First Aid</h5>	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.
- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Do not use this product on St. Augustinegrass, dichondra, carpetgrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

WHERE TO USE:

Weed-'N-Feed 6 is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

Weed-'N-Feed 6 is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Weed-'N-Feed 6 may be applied to newly seeded cool season turfgrass. Applications are suitable only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

Weed-'N-Feed 6 may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the use rate for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area.

HOW MUCH TO APPLY:

For Kentucky bluegrass, fescue spp., perennial ryegrass, bermudagrass, and zoysiagrass:

Apply 130 to 174 pounds of Weed-'N-Feed 6 per acre of established turfgrass. Or the equivalent application rates are 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application [1.5 lb 2,4-D ae, 0.33 lb MCP-p ae and 0.16 lb dicamba ae per acre per application]. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft. [3.0 lb 2,4-D ae, 0.66 lb MCP-p ae and 0.32 lb dicamba ae per acre per year].

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications is 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

APPLICATION SUGGESTIONS:

1. Spring and fall applications are suggested for the best control. However, Weed-N-Feed 6 can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application.

[Note to Supplemental Registrants: Refer to table for appropriate bag sizes.]

- For 3.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 333 square feet and adjust to apply at the rate of 1 pound of product per 333 square feet.
- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 3.4 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 300 square feet and adjust to apply at the rate of 1 pound of product per 300 square feet.
- For 3.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 275 square feet and adjust to apply at the rate of 1 pound of product per 275 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

[Note to Supplemental Registrants: Select the appropriate bag size and present only the information provided for that bag size on your product labeling. Also, the following language on equipment settings of drop spreaders, rotary spreaders, and other granular applications is optional – it may be presented on the container labels at your discretion]

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
12	3.0	4,000	26	3.0	8,700
	3.2	3,800		3.2	8,100
	3.4	-		3.4	7,600
	3.6	-		3.6	7,200
	4.0	-		4.0	6,500

Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	Bag Size, Lbs.	Application Rate, Lbs./1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.
13	3.0	4,300	28	3.0	9,300
	3.2	4,100		3.2	8,800
	3.4	3,800		3.4	8,200
	3.6	3,600		3.6	7,800
	4.0	3,250		4.0	7,000
14	3.0	4,700	36	3.0	12,000
	3.2	4,400		3.2	11,300
	3.4	4,100		3.4	10,500
	3.6	3,900		3.6	10,000
	4.0	3,500		4.0	9,000
15	3.0	5,000	37	3.0	12,300
	3.2	4,700		3.2	11,600
	3.4	4,400		3.4	10,900
	3.6	4,200		3.6	10,300
	4.0	3,750		4.0	9,250
16	3.0	5,300	39	3.0	13,000
	3.2	5,000		3.2	12,200
	3.4	4,700		3.4	11,500
	3.6	4,400		3.6	10,800
	4.0	4,000		4.0	9,750
17	3.0	5,700	40	3.0	13,300
	3.2	5,300		3.2	12,500
	3.4	5,000		3.4	11,800
	3.6	4,700		3.6	11,100
	4.0	4,250		4.0	10,000
18	3.0	6,000	42	3.0	14,000
	3.2	5,600		3.2	13,100
	3.4	5,300		3.4	12,400
	3.6	5,000		3.6	11,700
	4.0	4,500		4.0	10,500
24	3.0	8,000	45	3.0	15,000
	3.2	7,500		3.2	14,100
	3.4	7,100		3.4	13,200
	3.6	6,700		3.6	12,500
	4.0	6,000		4.0	11,300
25	3.0	8,300	50	3.0	16,700
	3.2	7,800		3.2	15,600
	3.4	7,400		3.4	14,700
	3.6	6,900		3.6	13,900
	4.0	6,250		4.0	12,500

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to XXX square feet. [Graphics Optional]

SINGLE PASS DISTRIBUTION [SPREADING PATTERNS]:

First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to ensure even coverage [Graphics Optional]. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes.

[Optional Spreader Charts]

Spreader settings for 1.0 pound of product per 310 square feet	
Scotts EasyGreen® Rotary	26
Scotts Drop	5
Scotts SpeedyGreen® Rotary	3.5

GreenView Drop (Spred II)	4
Central Drop (#205)	5.5
Sears Drop (#19810)	4
Ortho Drop (#5000)	5
Precision Drop (Red Devil)	4

[End optional language]

DISTRIBUTION AND SPREADER PATTERNS:

Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

BROADLEAF WEEDS CONTROLLED:

Dandelion, chickweed, plantain, and clover are controlled effectively with Weed-'N-Feed 6. A list of susceptible weeds is shown below.

Annual yellow sweetclover	Clover, crimson	Knotweed
Aster	Clover, hop	Kochia
Austrian fieldcress	Clover, red	Lambsquarters
Bedstraw	Clover, strawberry	Lespedeza
Beggarticks	Clover, sweet	Mallow
Betony, Florida	Clover, white	Matchweed
Bindweed, field	Cockle	Mexicanweed
Bird vetch	Cocklebur	Milk vetch
Bitter wintercress	Common mullein	Morningglory
Bittercress, hairy	Creeping jenny	Mouseear hawkweed
Bitterweed	Cudweed	Mugwort
Black-eyed Susan	Curly dock	Musk thistle
Black medic	Daisy, English	Mustard
Black mustard	Daisy fleabane	Narrowleaf plantain
Blackseed plantain	Daisy, oxeye	Narrowleaf vetch
Blessed thistle	Dandelion	Nettle
Bloodflower milkweed	Dichondra	Orange hawkweed
Blue lettuce	Dogfennel	Oriental cocklebur
Blue vervain	Dollarweed	Oxalis
Bracted plantain	Elderberry	Parsley-piert
Bristly oxtongue	False dandelion	Parsnip
Broadleaf dock	Falseflax	Pearlwort
Broadleaf plantain	False sunflower	Pennycress
Broomweed	Fiddleneck	Pepperweed
Buckhorn	Florida pusley	Pigweed
Buckhorn plantain	Frenchweed	Pineywoods bedstraw
Bulbous buttercup	Galinsoga	Plains <i>coreopsis</i>
Bull thistle	Goldenrod	Plantain
Bullnettle	Ground ivy	Poison ivy
Burclover	Gumweed	Poison oak
Burdock	Hairy fleabane	Pokeweed
Burning nettle	Hawkweed	Poorjoe
Burweed	Healall	Prairie sunflower
Buttercup	Heartleaf drymary	Prickly lettuce
Buttonweed	Heathaster	Prickly sida
Canada thistle	Hedge bindweed	Prostrate knotweed
Carolina geranium	Hedge mustard	Prostrate pigweed
Carpetweed	Hemp	Prostrate spurge
Catchweed bedstraw	Henbit	Prostrate vervain
Catnip	Hoary cress	Puncturevine
Catsear	Hoary plantain	Purslane, common
Chickweed, common	Hoary vervain	Ragweed
Chickweed, mouseear	Horsenettle	Red sorrel
Chicory	Jimsonweed	Redroot pigweed
Cinquefoil	Knawel	Redstem filaree

Rough cinquefoil
Rough fleabane
Russian pigweed
Russian thistle
Scarlet pimpernel
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smooth dock
Smooth pigweed
Sorrel
Sowthistle
Spanishneedles
Speedwell
Spiny amaranth
Spiny cocklebur
Spiny sowthistle
Spotted catsear
Spotted spurge
Spurweed

Stinging nettle
Strawberry, India mock
Tall nettle
Tall vervain
Tansy ragwort
Tansy mustard
Tanweed
Thistle
Trailing crownvetch
Tumble mustard
Tumble pigweed
Velvetleaf
Venice mallow
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Wavyleaf bullthistle
Western clematis
Western salsify
White mustard
Wild aster
Wild buckwheat

Wild carrot
Wild four-o'clock
Wild garlic
Wild geranium
Wild lettuce
Wild marigold
Wild mustard
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild vetch
Woodsorrel
Woolly croton
Woolly morningglory
Woolly plantain
Wormseed
Yarrow
Yellow rocket
Yellowflower pepperweed

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

[The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.]

OR

[FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.]

APPENDIX

1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:

Herbicide Benefit Claims:

- Kills Weeds
- [Apply/Use] when weeds are actively growing.
- [Kills/Controls] Over 200 listed Broadleaf Weeds
- [Kills/Controls] Over 200 Broadleaf Weeds as listed
- Kills over 200 listed Weeds!
- Controls dandelion, clover, & over [XXX] other broadleaf weeds as listed
- Controls dandelion, clover, & over [XXX] other listed broadleaf weeds
- Controls chickweed, dandelion, knotweed, plantain, henbit, spurge and many other weeds as listed
- Controls [insert weed from weed list] [and other listed weeds]
- Contains Trimec Herbicide
- Contains 3 proven weed killers
- 3-way active ingredient for optimum control
- Kills listed weeds to the root
- [Kills][Controls] most common weeds as listed
- Kills listed weeds and thickens lawns
- Best results are obtained when weeds are growing actively in spring or early fall

Fertilizer Benefit Claims:

- Controlled Release Nitrogen Nutralene®
- Controlled release technology feeds 2+ months
- This product contains polymer-coated Enhanced Efficiency Fertilizer technology. As your lawn demands nutrients, nitrogen is released slowly into the soil for plant root uptake. This controlled-release approach eliminates surge growth and provides predictable feeding for over two months.
- All [Brand] fertilizer products contain Enhanced Efficiency Fertilizer which ensures consistent feeding, deep greening and promotes overall lawn health.
- Promotes overall lawn health
- The ultimate timed release fertilizer as used by professional golf courses.
- Fall Lawn Food
- Winterizer


- Prepares Lawn For Winter
- Prepares Your Lawn For Winter
- Iron enriched formula
- Quickens spring green-up
- Feeds And Thickens Your Lawn
- Kills listed weeds And Thickens [Your] Lawn
- Long-lasting feeding
- Long-lasting greening
- Promotes a rich green lawn
- Contains Iron for Deep Greening
- Promotes color and growth
- Fortified with milorganite [logo/graphic]
- Fortified with Nutromatic® Ingredients
- Fortified with N-VIRO SR® [logo/graphic]
- Blended with N-VIRO SR® [logo/graphic]
- Enhanced with N-VIRO SR® [logo/graphic]
- Milorganite. For Better Results. Naturally.
- Natural lawn fertilizer
- Contains humic and fulvic acids
- [Contains] Iron for extra greening
- Stimulates plant growth
- Contains non-burning, slow release nitrogen and iron for deep greening
- Contains non-burning/slow-release nitrogen
- Improves turf quality
- Improves soil quality
- Improves turf and soil quality
- Zero phosphorus
- Phosphorus Free
- [Contains] Controlled/Slow release nitrogen for extended feeding
- Contains [X%] Iron for deep greening
- Feeds Your Lawn for [over] [x] weeks/months
- Deep greening [formula]
- Non-staining [formula]
- Stimulates [rich/deep green] color and growth
- Promotes root uptake of vital nutrients
- Promotes root uptake of nutrients
- Increases water retention
- Replaces [Provides] Nutrients for a Thicker, Greener Lawn
- Quick greening/green-up and/then continues to feed your lawn
- Slow release nitrogen feeds your lawn [turfgrass] over [x] months/weeks.
- Allow [6/X] weeks between applications [Note to reviewer: X weeks will be greater than 4 to adhere to minimum retreatment interval of 30 days]
- Feeds And Thickens Your Lawn
- Promotes color and growth
- Lush Green Lawn
- Formulated to promote root building and winter hardiness

General Other Marketing Statements Claims:

- A granular fertilizer containing Trimec® herbicide which controls: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and 200 other broadleaf weeds which are listed on the (back/side) panel.
- 2-In-1 Kills listed weeds and feed(s) your lawn
- Provides a full feeding of fertilizer and kills over listed 200 weeds

- Produce Superior Results...Every Time!
- [100%] Satisfaction Guaranteed Or Your Money Back
- [100%] Satisfaction Guarantee. If, after using this product, you are dissatisfied with the results, simply mail the sales receipt, a brief description of the problem and your contact information to the address listed below [or contact Menards guest services for a full refund].
- [100%] Satisfaction Guarantee. If, after using this product, you are dissatisfied with the results, simply mail the sales receipt, a brief description of the problem and your contact information to the address listed below for a full refund of the purchase price of the product.
- Stein's Policy – 100% Satisfaction Guaranteed. If for any reason you are not satisfied with the results after using this premium product, you are entitled to replacement bags of fertilizer or your money back. Simply bring proof of purchase to any Stein Garden Center.
- Calculating Total Lawn Area [graphic] Length x Width = Total Sq.Ft. then subtract Non-Lawn Areas (House, Garden, Driveway)
- Greenyard Professional (TM)
- Greenyard Professional Logo is a trademark of Growmark
- EC Grow www.ecgrowproducts.com TIPS – FAQ's – INFO
- 2nd application late spring/early summer [when used in 4 step program]
- Promotes thick & lush lawn
- Menard's Premium 100% Satisfaction Money Back Guarantee
- 4th application late summer/fall [when used in 4 step program]
- Covers XXX sq. ft.
- Apply When Weeds Are Actively Growing
- Use When Weeds Are Actively Growing
- Apply in fall for a greener lawn next spring
- Reseeding or over seeding areas may occur three (3) weeks after the application of this product.
- Apply when lawn is wet/moist
- Manufactured exclusively for [Stein's Garden & Home]
- For more information on this and other products, see us on the web at [website address].

Optional Spreader Related Images:



Referrals to other products used sequentially:

MENARDS® LAWN CARE PROGRAM
PREMIUM

1ST APPLICATION EARLY SPRING
 Season-Long Crabgrass Control

2ND APPLICATION LATE SPRING/EARLY SUMMER
 Controls dandelion, clover and 200 other broadleaf weeds

3RD APPLICATION MID/LATE SUMMER OR ANYTIME
 Promotes a lush and deep green lawn

4TH APPLICATION LATE SUMMER/FALL
 Prepares lawn for winter
 Quickens spring green-up

ALLOW 8 WEEKS BETWEEN APPLICATIONS

1st APP **CRABGRASS PREVENTER**
 APPLY EARLY SPRING
 with LAWN FOOD 28-0-3

2nd APP **WEED & FEED**
 APPLY LATE SPRING/EARLY SUMMER
 28-0-3

3rd APP **LAWN FOOD**
 APPLY MID/LATE SUMMER OR ANYTIME
 30-0-3

4th APP **FALL LAWN FOOD**
 APPLY LATE SUMMER/FALL
 WINTERIZER 22-0-10

ALLOW 8 WEEKS BETWEEN APPLICATIONS

HANK'S LAWN CARE Program

1ST APPLICATION EARLY SPRING
2ND APPLICATION LATE SPRING/EARLY SUMMER
3RD APPLICATION SUMMER OR ANYTIME
4TH APPLICATION LATE SUMMER/FALL

Allow 6 Weeks Between Applications

- [Brand] Lawn Care Program:
 - Allow 8 weeks between applications
 - Menard's Premium Crabgrass Preventer
 - 1st Application

- [Apply] Early Spring
 - Season-long crabgrass control
- Menard's Premium Weed & Feed

- 2nd Application
- [Apply] Late Spring/Early Summer
- Controls dandelions, clover and 200 other broadleaf weeds

- Menard's Premium Lawn Food
- 3rd Application
 - [Apply] Mid/Late Summer or Anytime
 - Promotes a lush and deep green lawn

Menard's Premium Fall Lawn Food

- 4th Application
- [Apply] Late Summer/Fall
- Prepares lawn for winter
- Quickens spring green-up

- [Brand] lawn care fertilizer program [insert graphic showing different application steps in the program with name]



- Lawncare [fertilizer] program [early spring, late spring/early summer, summer or anytime, late summer/fall] [insert graphic showing different application steps in the program with name]



STEIN LAWN CARE PROGRAM



- Application/App 1 [2][3][4]
- JRK Brand Lawn Care Program
- JRK Brand
- A Gertens® family of products.

2. State mandated fertilizer labeling requirements:

[Note to Supplemental Registrants: Please select the statements relevant to the areas of the country in which your specific product is registered and sold; if any of these statements are relevant to your labeling, please place in the Directions for Use under the heading “3.1 State Restrictions”]

GUARANTEED ANALYSIS XX-X-X	
Total Nitrogen (N)	XX%
x.x% ammoniacal nitrogen	
x.x% urea nitrogen*	
Available Phosphate (P2O5)	X%
Soluble Potash (K2O)	X%
Sulfur (S)	X%
x.x% free sulfur (S)	
Iron (Fe).....	X%
Derived from: polymer-coated sulfur-coated urea, urea, ammonium phosphate, potassium chloride. *A portion of the urea has been coated to provide x.x% coated slow-release nitrogen. This product and/or its manufacture is covered by one or more of the following U.S. patents and international counterparts thereof: US Patent Numbers: xxxxxx	

[Note: Fertilizer grades with nutrient contents expressed as the percentages of N, P2O5 and K2O may include the following: 23-3-7 or x-x-x. The fertilizer grade analyses may contain macronutrients and micronutrients. The nutrient contents expressed will vary among states, and the analyses will differ between supplemental registrations.]

- California, Oregon, Washington: “[When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.] “Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>”
- Florida: “Apply only to actively growing turf.”
- Florida: “Check with your local Cooperative Extension Agency to obtain specific information on local turf best management practices. Check with your county or city government to determine if there are local regulations for fertilizer use.
- Maryland: “This fertilizer contains phosphorus and may not be used on turf in the state of Maryland except when 1) Providing nutrients to specific soils and target vegetation as determined to be

necessary in accordance with a soil test that was conducted by a laboratory identified under § 8-803.7 of the Agriculture Article, Annotated Code of Maryland, performed no more than 3 years before the application; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State; or 3) Reestablishing or repairing a turf area.”

- New Jersey: “In the state of New Jersey, this product may be used only when 1) Providing nutrients to specific soils and target vegetation as determined to be necessary pursuant to a soil test conducted by a laboratory identified pursuant to subsection b. of section 6 of P.L.2010, c.112 (C.58:10A-66) and performed no more than three years before the application, and pursuant to the associated annual fertilizer recommendation issued by the New Jersey Agricultural Experiment Station at Rutgers, the State University; 2) Establishing vegetation for the first time, such as after land disturbance, provided the application is in accordance with the standards and requirements established under the “Soil Erosion and Sediment Control Act,” P.L.1975, c.251 (C.4:24-39 et seq.) and the rules and regulations adopted pursuant thereto; 3) Reestablishing or repairing a turf area; or 4) Delivering liquid or granular fertilizer under the soil surface, directly to the feeder roots.”
- New Hampshire: In the state of New Hampshire, this product may only be used on 1) Establishing new lawns, for repairing a lawn, for seeding, or for use when a soil test indicates a phosphorus deficiency.
- Virginia: “In the state of Virginia, this product may be used only on 1) Newly established turf areas from sod or seed during the first growing season; 2) Turf areas being repaired or renovated, and; 3) Turf areas where soil tests performed within the past three years indicate nutrient deficiency.”
- [Virginia, Florida, Maryland, others] DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN. [AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK INTO YOUR LAWN/GARDEN.] [Note to Supplemental Registrant: The bracketed part of this statement is not required for liquid products.]
- [For products containing Iron:] Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
- [Note to Supplemental Distributor: In Kentucky if the product supplies less than 1lb of Nitrogen per application, include the following:] A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing XX pounds of nitrogen per 1,000 square feet [and herbicide to control listed weeds]. Please consult your local extension office for questions regarding a regular fertilizer program [suitable for your geographic region].”

DOCUMENT CONTROL INFORMATION

1. Unique Label Identifier: 002217-00561.20220810.notif-proposed-highlighted.doc

2. Reason for Issue: ad claims, alternate warranty statements, and other minor updates

Internal Usage Notes
Associated Container Label: Supplemental Distributors