PM-23

1 3 5



1217 W. 12TH STREET P.O. BOX 4090 KANSAS CITY, MO 64101

The Certification of Compliance for the 2.4-D Exposure Reduction Program

"I, being an authorized representative of PBI/Gordon Corporation, certify that all containers of St. Augustinegrass Weed-N-Feed 580*(2217-568) produced by June 15, 1994 for products registered for use only on residential or turf sites, excluding sod farms will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said product sold or distributed by this company by January 1, 1995 for products registered for use only on residential or turf sites, excluding sod farms will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products."

J.A. Almelia.

Date

James A. Armbruster, Ph.D. Mgr. Regulatory Services

ST. AUGUSTINEGRASS WEED & FEED 580

20-10-5

LAWN FERTILIZER PLUS TRIMEC(R)

-COVERS 5,000 SQUARE FEET-

CONTROLS: Chickweed, Dandelion, Knotweed, Plantain, Henbit, Spurge, and many other weeds in St. Augustinegrass, Centipedegrass, Bentgrass and most other grasses.

ACTIVE INGREDIENT:

2, 4-dichlorophenoxyacetic acid		•		•	. 0.1746%
2-(2-methyl-4-chlorophenoxy) propionic acid					
Dicamba (3, 6-dichloro-o-anisic acid)	•	•		•	. 0.0635%
INERT INGREDIENTS	•	٠	•	•	.98.97228
					100.0000%

EQUIVALENT TO:

3.488 lbs of 2,4-dichlorophenoxyacetic acid per ton. 15.775 lbs of 2-(2-methyl-4-chlorophenoxy) propionic acid per ton. 1.268 lbs of (3,6-dichloro-o-anisic acid) per ton.

Isomer Specific by AOAC Methods.

TRIMEC^(R) is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel for Precautionary Statements and Statement of Practical Treatment

NET WEIGHT 20 POUNDS

/1292 APXXXXX

EPA REG. NO. 2217-568 EPA EST. NO. 2217-KS-1

Mfd. by PBI/GORDON CORPORATION
KANSAS CITY, MISSOURI 64101

3 7 5

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid inhaling dust or contact with skin, eyes and clothing. Wear long pants, long-sleeved shirt and shoes when applying this product.

When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

Environmental Hazards

Drift or runoff may adversely affect nontarge, plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

- USE DIRECTIONS -

Do not use this product where desirable clovers are present or allow the product to "wind-drift" on flowers, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. During application close spreader openings when turning. Apply when the air is calm. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species.

20-10-5	
GUARANTEED ANALYSIS	
Total Nitrogen (N)	20%
Water Insoluble Nitrogen (N)	. 5%
Available Phosphoric Acid (P2O5)	10%
Soluble Potash (K ₂ O)	. 5%

GENERAL INFORMATION: This product contains a special formulation low in 2,4-D for maximum safety to St. Augustinegrass. The fertilizer provides needed plant food while the herbicide will control hard-to-kill weeds such as creeping charlie, knotweed, chickweed, dandelions and plantains and most other broadleaf weeds.

If adverse low moisture conditions prevail and weeds are in a state of poor growth, possible retreatment may be necessary with a TRIMEC Herbicide liquid formulation when conditions improve.

This product can be used on such sensitive grasses as St. Augustinegrass, bahiagrass, centipedegrass, bermudagrass, bentgrass and zoysiagrass. Under conditions of adequate moisture, high temperature



495

and high humidity, over application may cause minor transitory discoloration on St. Augustine.

WHEN TO APPLY: For best weed control, application should be made when weeds are growing vigorously. Application can be made at any time but for maximum safety to St. Augustinegrass, make application during spring and fall when temperatures are moderate.

HOW MUCH TO USE: This 20 pound bag should be used on 5,000 square feet. Apply evenly with a lawn spreader either as a one-pass application or a two-pass application by crisscrossing the area at one half the rate each time.

To get maximum weed killing results, do not rake, mow, or water the lawn for at least 24 hours after spreading the product.

Lawns which are heavily overrun by weeds and are in poor condition may require more than one application of this product. A suggested second application could be made in approximately 30 days.

The maximum application rate to turf is 0.4 pounds 2,4-D acid equivalent per acre per application per site, The maximum number of broadcast applications per treatment site is 2 per year. Spreader setting should be adjusted as follows:

The above spreader settings are approximate. Calibrate your spreader on 100 square feet and adjust to apply at the rate of 1/2 pound of product per 100 square feet.

The following is a partial list of weeds controlled by this product:

Bedstraw Knotweed Black medic Lambsquarter Buckhorn Lespedeza Burdock Mallow Chicory Morningglory Chickweed Oxalis Clover Peppergrass Dandelion Pigweed Dock **Plantains** Ground by Poison Ivy Heal-All Poison Oak Henbit Purslane

Ragweed
Sheep sorrel
Sheepherdspurse
Speedwell
Spurge
Thistle
Wild Carrot
Wild Garlic
Wild Lettuce
Wild Onion
Yarrow

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential.



damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.