

For the control of Fruit decay (*Monilia*) and *Rhizopus* fruit rot 50  
lbs. per acre. Apply 10 days and 1 day prior to harvest.

**GRAPES:**

For the control of *Botrytis* rot, Powdery mildew, apply 30 lbs. per acre. Begin application while clusters are still loose and repeat application at two week intervals. Allow one day between application and harvest. Caution: Concord and other labrusca type grapes may be injured by sulfur.

**CONDITION OF SALE:** Britz Fertilizers, Inc. (BRITZ) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Britz neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Britz's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants and crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Britz under 1. above) resulting from handling, storage, and the use of this material.

FORMULATED BY

*Britz Fertilizers, Inc.*

P.O. BOX 366  
FIVE POINTS, MISSOURI 63624  
(209) 241-2421 Made in U.S.A.

E.P.A. Reg. No.

E.P.A. Reg. No. 10091 CA-5

Registered Trademark of NOR-AM Chemical Company  
U.S. Pat. No. 3,119,236

*"Britz for Results"*

NET WEIGHT 50 POUNDS

LOT NO. \_\_\_\_\_

**BEST COPY AVAILABLE**

Dear Dr. Armbruster:

SUBJECT: Label Amendment  
Gordon's Trimec St. Augustine Grass Broadleaf Herbicide  
EPA Registration No. 2217-570  
Your Application Dated January 6, 1995

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, is acceptable provided the following changes are made:

1. Under OPTION I/ SUPPLEMENTAL LABELING FOR DIFFERENT BRANDS, TYPES, OR SIZES OF HOSE-END APPLICATORS/OPTION B: Correct the typographical error in the last line of the table at the top of page 5 ("For St. Augustinegrass, Bentgrass and Buffalograss:"). The Area to Treat using 6 Fluid Ounces of product should be changed from "240" to "2400" square feet.
2. Under OPTION II/ SUPPLEMENTAL LABEL FOR ORTHO HOSE-END SPRAYERS: Correct the Tank Sprayer use directions "For Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Hahiagrass and Centipedegrass" at the top of page 7. The area to cover using 1 fluid ounce is incorrectly stated as "1000 square feet." This sentence should be revised to read:

"For Tank Sprayers: Use 1 fluid ounce (2 Tablespoons) per gallon of water to cover 200 square feet of lawn."

CONCURRENCES							
SYMBOL ▶	7505C						
SURNAME ▶	S. Stanton						
DATE ▶	Mar 30, 1995						

A stamped copy is enclosed for your records. Please submit (one) final printed copy for the above mentioned label before releasing the product for shipment.

Sincerely yours,

Joanne I. Miller  
Product Manager (23)  
Fungicide-Herbicide Branch  
Registration Division (7505C)

Enclosure

RD:STANTON:CM-2:Rm. 235:305-5218:March 30, 1995:Disk #1:  
S479836.LET

2

# GORDON'S TRIMEC ST. AUGUSTINE GRASS BROADLEAF HERBICIDE

## ACTIVE INGREDIENTS:

* Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid.....	10.60%
** Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid.....	3.05%
*** Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid). ....	1.30%
INERT INGREDIENTS.....	<u>85.05%</u>
TOTAL	100.00%

## THIS PRODUCT CONTAINS:

- \*0.75 lb. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 8.76%
  - \*\*0.22 lb. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 2.53%
  - \*\*\*0.093 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.08%
- Isomer Specific by AOAC Method 6.DO1-5.  
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side panels for additional Precautionary Statements.

NET CONTENTS [ONE PINT, ONE QUART, or ONE U.S. GALLON]

880/ AP122094  
EPA REG NO. 2217-570  
EPA EST. NO. 2217-KS-1

Mfd. by PBI/GORDON CORPORATION  
KANSAS CITY, MISSOURI 64101

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

MAR 31 1995  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
2217-570

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals:**

**CAUTION:** Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wear face shield or goggles when mixing, loading or applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

**Personal Hygiene Statements:**

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

**Statement of Practical Treatment**

**In case of eye contact:** Immediately flush eyes with plenty of water. If irritation develops, get medical attention.

**In case of skin contact:** Thoroughly wash hands and exposed skin with soap and water. If irritation develops, get medical attention.

**ENVIRONMENTAL HAZARDS:** This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

=====

[OPTION I. Directions For Use Permitted to the Acme Division of PBI/Gordon Corporation.]

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

**STORAGE & DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide, or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent) Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**OTHER USE PRECAUTIONS:**

Do not apply this product through any type of irrigation system

Do not allow spray to drift onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants since injury may result. Do not pour spray solutions near these plants.

Whether using a hose-end sprayer or pressure-sprayer, do not use fine sprays because they can drift and cause damage to non-target plants. Coarse spray droplets are less likely to wind-drift. Use only lawn-type sprayers.

Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Do not spray roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful with dripline of trees and other ornamental species.

Avoid spraying St. Augustinegrass when temperature is above 90°F and under conditions of high humidity. Under these conditions, some slight temporary discoloration of the St. Augustinegrass foliage may occur. This product may be injurious to St. Augustinegrass, bentgrass and centipedegrass in cases of accidental overdose.

**WHAT IT IS, WHERE & HOW TO USE IT:**

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide is a TRIMEC® turf herbicide formulation, a combination of three proven weed killers: 2,4-D, mecoprop and dicamba. Together, they control a wide range of broadleaf lawn weeds. This particular formulation is low in 2,4-D for maximum safety to 2,4-D sensitive grasses such as St. Augustinegrass, bentgrass and buffalograss. Spraying should be done when broadleaf weeds are small and tender for best results.

The maximum application rate to turf is 0.4 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Irrigate the lawn 2 or 3 days prior to spraying. Do not spray lawns under drought conditions.
2. Do not water for 24 hours after treatment.
3. To avoid damage do not apply to St. Augustinegrass or centipedegrass while emerging from winter dormancy. Also, damage may occur to these grasses if temperatures drop below 45°F within 72 hours of application.

**ST. AUGUSTINEGRASS, BENTGRASS & BUFFALOGRASS.** Mix 1 fluid ounce in 1 gallon water and uniformly spray 400 square feet. If necessary, repeat treatment in 2 to 3 weeks if hard-to-kill weeds persist. Do not use on Floratam or other improved varieties of St. Augustinegrass in Florida.

**BERMUDAGRASS, BAHIAGRASS, BLUEGRASS, CENTIPEDEGRASS, FESCUES & RYEGRASS:** Mix 1 1/2 fluid ounces in 1 gallon water and uniformly spray over 400 square feet. On hard-to-kill species, or on older weeds, mix 2 fluid ounces in 1 gallon water and spray on 400 square feet.

- WEEDS -			
Bedstraw	Dock	Mustard	Speedwell
Black Medic	Galinsoga	Oxalis	Tansy Mustard
Buckhorn	Ground Ivy	Peppergrass	Thistle
Bull Thistle	Hawkweed	Pigweed	Spurge
Burdock	Heal-All	Plantain	Velvetleaf
Buttercup	Henbit	Poison Ivy	Wild Carrot
Chicory	Knotweed	Poison Oak	Wild Garlic
Chickweed	Lambsquarters	Purslane	Wild Lettuce
Clover	Lespedeza	Ragweed	Wild Onion
Cocklebur	Mallow	Sheep Sorrel	Yarrow and other
Dandelion	Morningglory	Shepherdspurse	broadleaf weeds

10/10

**SUPPLEMENTAL LABELING FOR DIFFERENT BRANDS, TYPES, OR SIZES OF HOSE-END APPLICATORS.**

See primary label on container for Precautionary Statements, the Statement of Practical Treatment, and Full Directions For Use.

**OPTION A:**

**(INSERT BRAND NAME) HOSE-END SPRAYER INSTRUCTIONS:** Use a hose-end sprayer designed for lawns. Put recommended amount of product (Column 1) in bottle and add water to indicated level (Column 2) and apply spray uniformly to area indicated (Column 3).

For Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, Centipedegrass:

Product	Amount to Use:	Water	Area to Treat:
(Tbsp)	(Fluid Ounces)	Fill Bottle to:	(Square Feet)
8	4	8	1000
16	8	16	2000
24	12	24	3000
32	16	32	4000

For St. Augustinegrass, Bentgrass and Buffalograss:

Product	Amount to Use:	Water	Area to Treat:
(Tbsp)	(Fluid Ounces)	Fill Bottle to:	(Square Feet)
5.0	2.5	5.0	1000
10.0	5.0	10.0	2000
15.0	7.5	15.0	3000
20.0	10.0	20.0	4000
25.0	12.5	25.0	5000

**OPTION B:**

**(INSERT BRAND NAME) HOSE-END SPRAYER INSTRUCTIONS:** Use a hose-end sprayer designed for lawns. Put recommended amount of product (Column 1) in bottle and add water to indicated level (Column 2) and apply spray uniformly to area indicated (Column 3).

For Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, Centipedegrass:

Product Amount to Use:	Water	Area to Treat:
(Fluid Ounces)	Fill Bottle to:	(Square Feet)
1 1/2	1	400
3	2	800
6	4	1600
9	6	2400

7/2/10

Product Amount to Use: (Fluid Ounces)	Water Fill Bottle to: (Gallons)	Area to Treat: (Square Feet)
1	1	400
2	2	800
4	4	1600
6	6	240

End of Option I:-----

**[OPTION II. Proposed Directions For Use. This supplemental product registration will be intended for use by homeowners only on ornamental lawns and turfgrass established in residential sites.]**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

**STORAGE & DISPOSAL**

**STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse container (bottle, can, jar). Rinse thoroughly before discarding in trash.

**WHAT IT IS:** Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide is a Trimec® turf herbicide formulation, a combination of three proven weed killers 2,4-D mecoprop and dicamba. Together, they control a wide range of broadleaf lawn weeds in residential lawns.

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide kills all major broadleaf lawn weeds such as dandelion, chickweed, knotweed, plantains in lawns without harming lawn grasses. Other weeds include the following:

- WEEDS -			
Bedstraw	Cinquefoil	Lespedeza	Ragweed
Beggarweed	Clover	Mallow	Sheep Sorrel
Bindweed	Cocklebur	Matchweed	Shepherdspurse
Bittercress	Dandelion	Morningglory	Speedwell
Black Medic	Dock	Mustard	Tansy Mustard
Buckhorn	English daisy	Oxalis	Thistle
Bull Thistle	Galinsoga	Pennywort	Spurge
Burdock	Geranium	Peppergrass	Velvetleaf
Burweed	Ground Ivy	Pigweed	Wild Carrot
Buttercup	Hawkweed	Plantain	Wild Garlic
Buttonweed	Heal-All	Poison Ivy	Wild Lettuce
Carpetweed	Henbit	Poison Oak	Wild Onion
Catsear	Knawel	Prickly lettuce	Yarrow and other
Chicory	Knotweed	Purslane	broadleaf weeds
Chickweed	Lambsquarters		

7



7-7-10

**WHERE TO USE:** Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide is intended for use on most ornamental lawns including Kentucky bluegrass, perennial ryegrass, fescues, bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, and St. Augustinegrass.

- Do not use this product on Floratam, carpetgrass, dichondra, or lawns containing desirable clovers. Do not spray roots of trees and ornamentals.

**HOW TO TIPS:**

- Spray when broadleaf weeds are small and actively growing for best results.
- Be sure lawn is well-watered. If dry, water 1 to 2 days before spraying. To reduce risk of lawn injury, avoid spraying drought stressed lawns or when temperatures are above 90°F.
- Spray when air is calm to avoid spray drift onto landscape plants as contact with foliage may injure flowers, shrubs, trees, and other desirable plants.
- Rain or watering within 24 hours after treatment may reduce effectiveness.
- For best results, use hose-end sprayers or tank-type (pump-up pressure) sprayers. Spray evenly over area at recommended rates.

**HOW MUCH TO USE:** The maximum application rate to turf is 0.4 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year. Do not exceed specified dosages for any area.

For St. Augustinegrass, Bentgrass, and Buffalograss

For Tank Sprayers: Use 1/2 fluid ounce (1 Tablespoon) per gallon of water to cover 200 square feet of lawn. Do not use on Floratam or other improved varieties of St. Augustinegrass in Florida.

Do not apply to St. Augustinegrass or Centipedegrass emerging from winter dormancy. Temperatures below 45°F within 72 hours of application may cause damage.

For Bluegrass, Fescues, Perennial Ryegrass, Bahiagrass, Bermudagrass, and Centipedegrass

Use 1 fluid ounce (2 Tablespoons) per gallon of water to cover 200 square feet.

Repeat treatment for hard-to-control weeds in 2 to 3 weeks if they regrow.

Spot treat new weeds as they appear with Weed-B-Gon® Weed Killer Ready-To-Use or use this product as directed in a tank-type sprayer. Do not use a hose-end sprayer to spray small areas or spot treat weeds.

CONVERSION TABLE	
1 Tablespoon (Tbsp)	= 3 Teaspoons (Tsp.)
1 Fluid Ounce (Fl. Oz.)	= 2 Tablespoons (Tbsp.)
1 Cup	= 8 Fluid Ounces (Fl. Oz.)

=====
**SUPPLEMENTAL LABEL FOR ORTHO HOSE-END SPRAYERS**

For St. Augustinegrass, Bentgrass and Buffalograss:

- For Tank Sprayers: Use 1/2 fluid ounce (1 Tablespoon) per gallon of water to cover 200 square feet of lawn.
- For the Dial 'N spray Hose-End Sprayer: Measure 2 1/2 fluid ounces (5 Tablespoons) into sprayer jar for every 1,000 square feet of lawn. Do not add water to jar. Set dial to 1 Tbsp. and spray.
- For the Ortho Plant & Weed Spray-ette® Hose-End Sprayer: Put 2 1/2 fluid ounces (5 Tablespoons) into sprayer jar. Fill sprayer jar to the 5 gallon mark with water and spray to cover 1,000 square feet.

6

For Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass and Centipedegrass

- For Tank Sprayers: Use 1 fluid ounce (2 Tablespoons) per gallon of water to cover 1000 square feet of lawn.
- For Dial 'N Spray: Measure 5 fluid ounces (10 Tablespoons) into sprayer jar for every 1,000 square feet of lawn. Do not add water to the container jar. Set dial on 1 oz. and spray.
- For Ortho Plant & Weed Spray-ette® Hose-End Sprayer: Measure 5 fluid ounces (10 Tablespoons) into sprayer jar. Fill with water to the 5 gallon mark and spray to cover 1,000 square feet.

**OR TABLE FORM**

Lawn Grass	Sprayer Type	Product To Measure	Area to Treat	Comments
St. Augustinegrass*, Bentgrass and Buffalograss	Tank Sprayer	1/2 fl. oz. (1 Tbsp) into 1 gallon of water and spray.	200 sq. ft.	*Do not use on Floratam or other improved varieties of St. Augustinegrass in Florida.
	Dial 'N Spray	2 1/2 fl. ozs. (5 Tbsp) into sprayer jar. DO NOT ADD WATER. Set dial on 1 Tbsp. and spray.	1000 sq. ft.	
	Ortho Plant & Weed Spray-ette®	2 1/2 fl. ozs. (5 Tbsp) into sprayer jar. Fill with water to the 5 gallon mark and spray.	1000 sq. ft.	

Lawn Grass	Sprayer Type	Product to Measure	Area To Treat
Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, and Centipedegrass	Tank Sprayer:	1 fl. oz. (2 Tbsp) into 1 gallon of water and spray.	200 sq. ft.
	Dial 'N Spray:	5 fl. ozs. (10 Tbsp) into sprayer jar. DO NOT ADD WATER. Set dial on 1 oz. and spray.	1000 sq. ft.
	Ortho Plant & Weed Sprayette®	5 fl. ozs. (10 Tbsp) into sprayer jar. Fill with water to the 5 gallon mark and spray.	1000 sq. ft.

End of Option II.=====

**LIMITED WARRANTY AND DISCLAIMER.**

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.