

2217-570

8/19/2010

10423



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

AUG 19 2010

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

James L. Kunstman, Ph.D.  
PBI/Gordon Corporation  
PO Box 014090  
Kansas City, MO 64101

Subject: Notification: Revised Disposal Instructions (PR Notice 2007-4); and Reinstatement of  
Zoysiagrass on Label (PR Notice 98-10)  
Gordon's Trimec St. Augustinegrass Broadleaf Herbicide  
EPA Reg. No. 2217-570  
Application, Dated July 15, 2010

Dear Dr. Kunstman:

The Agency is in receipt of your Application for Pesticide Notification under  
Pesticide Registration Notices (PRN) 2007-4 and 98-10 for the subject product.

The Registration Division (RD) has conducted a review of this request for applicability  
under PRN 2007-4, 2008-1, and 98-10 and finds that the label changes requested fall within the  
scope of the PR Notices. The label has been date-stamped "Notification" and will be placed in our  
records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other  
code identifying the batch of the pesticide distributed and sold be placed on nonrefillable  
containers. The code may appear either on the label (and can be added by non-notification/PR  
Notice 98-10) or durably marked on the container itself.

If you have any questions, please call me directly at (703) 305-1243 or Susan Stanton of my staff  
at (703) 305-5218.

Sincerely,

A handwritten signature in cursive script that reads "Susan L. Stanton, for".

Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs



United States  
**Environmental Protection Agency**  
 Washington, DC 20460

Registration  
 Amendment  
 Other

OPP Identifier Number

**Application for Pesticide - Section I**

1. Company/Product Number <b>2217-570</b>	2. EPA Product Manager <b>Kathryn V. Montague</b>	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) <b>Gordon's Trimec St. Augustinegrass Broadleaf Herbicide</b>	PM# <b>Product Manager—Team 23</b>	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ <b>NOTIFICATION</b> Product Name _____ <b>2217-570</b>	

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

**Labeling notification per Pesticide Registration Notices (PRN) 2007-4 and 98-10.**

Please refer to cover letter for details and certification statements

e-mail to [jkunstman@pbigordon.com](mailto:jkunstman@pbigordon.com) FAX: 816-421-2731

**Section - III**

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per Container
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 1 pint to 1 gallon		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name <b>James L. Kunstman, Ph.D.</b>		Title <b>Director of Regulatory Services</b>		Telephone No. (Include Area Code) <b>816-460-6292</b>	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
2. Signature 		3. Title <b>Director of Regulatory Services</b>			
4. Typed Name <b>James L. Kunstman, Ph.D.</b>		5. Date <b>July 15, 2010</b>			







Language Common to All Sublabels

GORDON'S TRIMEC® ST.  
AUGUSTINEGRASS BROADLEAF  
HERBICIDE

NOTIFICATION

AUG 19 2010

EPA Reg. No. 2217-570

ACTIVE INGREDIENT:

Mecoprop-p, dimethylamine salt .....	5.30%
2,4-D, dimethylamine salt .....	3.05%
Dicamba, dimethylamine salt .....	1.29%

INERT INGREDIENTS: ..... 90.36%

TOTAL ..... 100.00%

THIS PRODUCT CONTAINS:

0.37 lb. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 4.38%

0.22 lb. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 2.53%

0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.08%

Isomer Specific by AOAC Method 6.DO1-5.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

First Aid	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

**Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

80923

---

## SUBLABEL 1

---

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **GENERAL PRECAUTIONS RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

#### **USE PRECAUTIONS:**

Do not apply this product through any type of irrigation system.

Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Do not exceed specified dosages for any area.

Avoid spraying St. Augustinegrass when temperature is above 90F and under conditions of high humidity. Under these conditions, some slight temporary discoloration of the St. Augustinegrass foliage may occur. This product may be injurious to St. Augustinegrass, bentgrass and centipede grass in cases of accidental overdose.

#### **Other Use Precautions:**

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

#### **WHAT IT IS, WHERE & HOW TO USE IT:**

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide is a Trimec® turf herbicide formulation, a combination of three proven weed killers: 2,4-D, mecoprop-p and dicamba. Together, they control a wide range of broadleaf lawn weeds. This particular formulation is low in 2,4-D for sensitive grasses such as St. Augustinegrass, bentgrass and buffalograss. Spraying should be done when broadleaf weeds are small and growing actively for best results.

1. Irrigate the lawn 2 or 3 days prior to spraying. Do not spray lawns under drought conditions.
2. Do not water for 24 hours after treatment.
3. To avoid damage do not apply to St. Augustinegrass or centipede grass while emerging from winter dormancy. Also, damage may occur to these grasses if temperatures drop below 45°F within 72 hours of application.

#### **HAND-OPERATED SPRAYER INSTRUCTIONS**

**St. Augustinegrass, Bentgrass and Buffalograss:** Mix 1 fl.oz. in 1 gallon water and uniformly spray 400 sq.ft. If necessary, repeat treatment in 30 days if hard-to-kill weeds persist. Do not use on 'Floritam' St. Augustinegrass in Florida.

**Bermudagrass, Bahiagrass, Kentucky Bluegrass, Centipede grass, Fescues, Perennial Ryegrass, and Zoysiagrass:** Mix 1.5 fl.oz. in 1 gallon water and uniformly spray 400 sq.ft. On hard-to-kill species, or on older weeds, mix 2 fl.oz. in 1 gallon water and spray on 400 sq.ft.



**Limitations on spot treatments to residential turfgrass:**

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 4.0 fl.oz. per 1,000 sq.ft. per application [0.50 lbs MCPP-p acid equivalent per acre]. The maximum number of spot treatment applications is 2 per year with a minimum of 30 days between applications.

<b>WEEDS</b>			
Bedstraw	Dock	Mustard	Speedwell
Black medic	Galinsoga	Oxalis	Spurge
Buckhorn	Ground ivy	Peppergrass	Tansy mustard
Bull thistle	Hawkweed	Pigweed	Thistle
Burdock	Healall	Plantain	Velvetleaf
Buttercup	Henbit	Poison ivy	Wild carrot
Chicory	Knotweed	Poison oak	Wild garlic
Chickweed	Lambsquarters	Purslane	Wild lettuce
Clover	Lespedeza	Ragweed	Wild onion
Cocklebur	Mallow	Sheep sorrel	Yarrow and other
Dandelion	Morningglory	Shepherdspurse	broadleaf weeds

**For Different Brands, Types, or Sizes of Hose End Sprayers**

**A.**

**(INSERT BRAND NAME) HOSE END SPRAYER INSTRUCTIONS:** Use a hose end sprayer designed for lawns. Pour recommended amount of product (Column 1) in bottle and add water to indicated level (Column 2) and apply spray uniformly to area indicated (Column 3).

<b>For Kentucky Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, Centipedegrass, and Zoysiagrass:</b>			
<b>Product, Amount to Use:</b>		<b>Water, Fill Bottle to: (Gallons)</b>	<b>Area to Treat: (Square Feet)</b>
<b>(Tablespoons)</b>	<b>(Fluid Ounces)</b>		
8	4	8	1,000
16	8	16	2,000
24	12	24	3,000
32	16	32	4,000

<b>For St. Augustinegrass, Bentgrass and Buffalograss:</b>			
<b>Product, Amount to Use:</b>		<b>Water, Fill Bottle to: (Gallons)</b>	<b>Area to Treat: (Square Feet)</b>
<b>(Tablespoons)</b>	<b>(Fluid Ounces)</b>		
5	2.5	5	1,000
10	5.	10	2,000
15	7.5	15	3,000
20	10.	20	4,000
25	12.5	25	5,000

**B.**

**(INSERT BRAND NAME) HOSE END SPRAYER INSTRUCTIONS:** Use a hose end sprayer designed for lawns. Pour recommended amount of product (Column 1) in bottle and add water to indicated level (Column 2) and apply spray uniformly to area indicated (Column 3).

For Kentucky Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, Centipedegrass, and Zoysiagrass:		
Product, Amount to Use: (Fluid Ounces)	Water, Fill Bottle to: (Gallons)	Area to Treat: (Square Feet)
1.5	1	400
3	2	800
6	4	1,600
9	6	2,400

For St. Augustinegrass, Bentgrass and Buffalograss:		
Product, Amount to Use: (Fluid Ounces)	Water, Fill Bottle to: (Gallons)	Area to Treat: (Square Feet)
1	1	400
2	2	800
4	4	1,600
6	6	2,400

CONVERSION TABLE
1 Tablespoon (Tbs.) = 3 teaspoons (tsp.)
1 Fluid Ounce (fl.oz.) = 2 Tablespoons (Tbs.)
1 Cup = 8 Fluid Ounces (fl.oz.)

**Limitations on broadcast applications to residential turfgrass:**  
 The maximum application rate is 4.0 fl.oz. of product per 1,000 sq.ft. per application (0.30 lbs 2,4-D ae, 0.50 lb MCPP-p ae and 0.12 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl.oz. of product per 1,000 sq.ft. (0.60 lbs 2,4-D ae, 1.0 lb MCPP-p ae and 0.25 lb dicamba ae per acre).

**STORAGE & DISPOSAL [Trashcan Illustration Optional]**  
 Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**  
 FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

## SUBLABEL 2

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

#### **STORAGE & DISPOSAL** [Trashcan Illustration Optional]

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### WHAT IT IS:

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide is a Trimec® turf herbicide formulation, a combination of three proven weed killers 2,4-D, mecoprop-p and dicamba. Together, they control a wide range of broadleaf lawn weeds in residential lawns.

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide kills major northern and southern broadleaf lawn weeds such as dandelion, chickweed, knotweed, plantain in lawns without harming lawn grasses when used as directed. Other weeds include the following:

#### **[Weed Illustration Optional]**

WEEDS:			
Annual aster	Ragweed	Mallow	Red clover
Bedstraw	Cocklebur	Matchweed	Red henbit
Beggarweed	Common chickweed	Morningglory	Scarlet pimpernel
Bindweed	Cudweed	Mouseear chickweed	Sheep sorrel
Bittercress	Curly dock	Musk thistle	Shepherdspurse
Black medic	Dandelion	Mustard	Sowthistle
Brassbuttons	Dock	Narrowleaf plantain	Speedwell
Broadleaf plantain	English daisy	Oxalis	Spotted spurge
Buckhorn	False dandelion	Pennsylvania smartweed	Spurge
Bull thistle	Florida pusley	Pennywort(creeping Charlie)	Tansy mustard
Burdock	Galinsoga	Peppergrass	Thistle
Burclover	Geranium	Pigweed	Velvetleaf
Burweed	Ground ivy	Piney woods bedstraw	White clover
Buttercup	Hairy fleabane	Plantain	Wild carrot
Buttonweed	Hawkweed	Poison ivy	Wild garlic
Carolina geranium	Healall	Poison oak	Wild lettuce
Carpetweed	Henbit	Prickly lettuce	Wild onion
Catsear	Knawel	Prostrate spurge	Wild strawberry
Chicory	Knotweed	Puncturevine	Yarrow
Chickweed	Lambsquarters	Purslane	and other broadleaf weeds
Cinquefoil	Lespedeza		
Clover			

**WHERE TO USE:**

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide is intended for use on most ornamental lawns including Kentucky bluegrass, perennial ryegrass, fescues, bermudagrass, bahiagrass, bentgrass, buffalograss, centipede grass, zoysiagrass, and St. Augustinegrass.

Do not use this product on 'Floritam' St. Augustinegrass, carpetgrass, dichondra, or lawns containing desirable clovers. Do not spray roots of trees and ornamentals.

**HOW TO TIPS:**

- Spray when broadleaf weeds are small and actively growing for best results.
- Be sure lawn is well watered. If dry, water 1 to 2 days before spraying. To reduce risk of lawn injury, avoid spraying drought stressed lawns or when temperatures are above 90°F.
- Rain or watering within 24 hours after treatment may reduce effectiveness.
- For best results, use hose end sprayers or tank type (pump-up pressure) sprayers. Spray evenly over area at recommended rates.

**Other Use Precautions:**

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

**HOW MUCH TO USE:**

Spot treatments:

**For St. Augustinegrass, Bentgrass, Buffalograss, and Centipede grass:**

For Tank Sprayers - Use 1/2 fluid ounce (1 Tablespoon) per gallon of water to cover 200 square feet of lawn. Do not use on Floritam or other improved varieties of St. Augustinegrass.

Do not apply to St. Augustinegrass or centipede grass emerging from winter dormancy. Temperatures below 45°F within 72 hours of application may cause damage.

**For Bluegrass, Fescues, Perennial Ryegrass, Bahiagrass, Bermudagrass, and Zoysiagrass:** Use 1 fluid ounce (2 Tablespoons) per gallon of water to cover 200 square feet.

Repeat treatment for hard-to-control weeds in 30 days if they regrow.


Do not use a hose end sprayer to spray small areas or spot treat weeds.

**Limitations on spot treatments to residential turfgrass:**

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 5.0 fl.oz. per 1,000 sq.ft. per application [0.63 lbs MCP-p acid equivalent per acre]. The maximum number of spot treatment applications is 2 per year with a minimum of 30 days between applications.

CONVERSION TABLE
1 Tablespoon (Tbs.) = 3 teaspoons (tsp.)
1 Fluid Ounce (fl.oz.) = 2 Tablespoons (Tbs.)
1 Cup = 8 Fluid Ounces (fl.oz.)

[Note to supplemental registrant. Select the appropriate container size and present only the information provided for that size on your product labeling.]

<b>PRODUCT FACTS:</b>	
• Treats up to 6,400 square feet (approximate size of 3 tennis courts). Pint Size	
• Treats up to 12,800 square feet (approximate size of 6 tennis courts). Quart Size	
• Treats up to 21,200 square feet (approximate size of 10 tennis courts). 42 Gallon Size	
• Treats up to 51,200 square feet (approximate size of 24 tennis courts). Gallon Size	
<b>KILLS WEEDS:</b>	Dandelion, Clover, Chickweed <b>plus over</b> [Optional "No" Weed Icon] 80 additional broadleaf weeds
<b>WHERE TO USE:</b>	<b>ON LAWNS</b> [Optional Lawn Illustration] without harming lawn grasses when used as directed.
<b>AMOUNT TO USE:</b>	Bent, Buffalo, Centipede, and St. Augustine (Except 'Floritam'): 1 fluid ounce (2 Tablespoons) / gallon of water / 400 square feet
<b>AMOUNT TO USE:</b>	Bahia, Bermuda, Fescues, Kentucky Blue, Perennial Rye and Zoysia: 2 fluid ounces (4 Tablespoons) / gallon of water / 400 square feet
	Questions, Comments or Medical Information? Call: 1-800-xxx-xxxx • <a href="http://www.xxxx.com">http://www.xxxx.com</a>

**FOR ORTHO HOSE-END SPRAYERS.**

**Narrative format:**

**For St. Augustinegrass, Bentgrass, Buffalograss, and Centipedeagrass:**

- **For Tank Sprayers:** Use 1/2 fl.oz. (1 Tbs.) per gallon of water to cover 200 sq.ft. of lawn.
- **For the Dial 'N Spray Hose-End Sprayer:** Measure 2.5 fl.ozs. (5 Tbs.) into sprayer jar for every 1,000 sq.ft. of lawn. Do not add water to jar. Set dial to 1 Tbs. and spray.
- **For the Ortho Plant & Weed Sprayette® Hose-End Sprayer:** Put 2.5 fl.ozs. (5 Tbs.) into sprayer jar. Fill sprayer jar to the 5 gallon mark with water and spray to cover 1,000 sq.ft.

**For Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, and Zoysiagrass:**

- **For Tank Sprayers:** Use 1 fl.oz. (2 Tbs.) per gallon of water to cover 200 sq.ft. of lawn.
- **For Dial 'N Spray:** Measure 5 fl.ozs. (10 Tbs.) into sprayer jar for every 1,000 sq.ft. of lawn. Do not add water to the container jar. Set dial on 1 oz. and spray.
- **For Ortho Plant & Weed Sprayette® Hose-End Sprayer:** Measure 5 fl.ozs. (10 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray to cover 1,000 sq.ft.

**Tabular Format**

**Table 1.**

Lawn Grass	Sprayer Type	Product To Measure	Area to Treat	Comments
St. Augustinegrass*, Bentgrass, Buffalograss, and Centipedeagrass	Tank Sprayer	1/2 fl.oz. (1 Tbs.) into 1 gallon of water and spray.	200 sq.ft. (10' x 20')	*Do not use on 'Floritam' St. Augustinegrass
	Dial 'N Spray®	2.5 fl.ozs. (5 Tbs.) into sprayer jar. DO NOT ADD WATER. Set dial on 1 Tbs. and spray.	1,000 sq.ft.	Do not apply to St. Augustinegrass and centipedeagrass emerging from winter dormancy.
	Ortho Plant & Weed Sprayette®	2.5 fl.ozs. (5 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray.	1,000 sq.ft.	Temperatures below 45°F within 72 hours of application may cause damage

**Table 2.**

Lawn Grass	Sprayer Type	Product to Measure	Area To Treat
Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, and Zoysiagrass	Tank Sprayer:	1 fl.oz. (2 Tbs.) into 1 gallon of water and spray.	200 sq.ft. (10' x 20')
	Dial 'N Spray®	5 fl.ozs. (10 Tbs.) into sprayer jar. DO NOT ADD WATER. Set dial on 1 oz. and spray.	1,000 sq.ft.
	Ortho Plant & Weed Sprayette®	5 fl.ozs. (10 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray.	1,000 sq.ft.

**Table 3. MEASURING CAP CONTAINER**

MEASURING CAP: To open: squeeze cap side while unscrewing.

**Chart 1: DILUTION TABLE**

Gallons of Spray	Area to Treat (sq.ft.)	Hardy Lawns (bluegrass, zoysiagrass, fescues, ryegrass, bermudagrass, bahiagrass)	Sensitive Lawns** (St. Augustinegrass, bentgrass, buffalograss, centipedegrass)
1	200	Fill cap to level 3 (1 fl.oz.)	Fill cap to level 2 (1/2 fl.oz.)
5	1,000	Fill cap to level 3 five times (5 fl.oz.)	Fill cap to level 2 five times (2.5 fl.oz.)

The measuring cap has 4 levels indicated inside the cap. Fill cap to appropriate level for fast, easy accurate measuring. Close cap tightly after use — no need to rinse.

- Level 1 = 1 teaspoon (tsp.)
- Level 2 = 1/2 fl.oz. = 1 Tablespoon (Tbs.) = 3 tsp.
- Level 3 = 1 fl.oz. (2 Tbs.)
- Level 4 = 2 fl.oz. ( 4 Tbs.)

\*\*See specific directions for St. Augustinegrass and other sensitive lawns.

**Limitations on broadcast applications to residential turfgrass:**

The maximum application rate is 5.0 fl.oz. of product per 1,000 sq.ft. per application (0.37 lbs 2,4-D ae, 0.63 lb MCP-P ae and 0.15 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.0 fl.oz. of product per 1,000 sq.ft. (0.75 lbs 2,4-D ae, 1.26 lb MCP-P ae and 0.31 lb dicamba ae per acre).

**Instructions For Use With The Ortho Dial'N Spray and Measuring Cap Containers. THREE EASY STEPS TO MIX AND APPLY**

1. Measure the lawn area to be sprayed. Calculate square feet by multiplying length by width.
2. Use a clean sprayer, such as a tank type or hose end sprayer. Carefully mix the proper amount of concentrate with water. See below for directions for the Ortho Dial 'N Spray Hose End Sprayer.
3. Spray evenly over the lawn.

**DIRECTIONS FOR ORTHO DIAL 'N SPRAY HOSE END SPRAYER**

1. Measure the proper amount of product (see Chart 1) into the sprayer jar for the number of square feet you need to cover. Do not add water to the sprayer jar.
2. Set the dial to 1 Tbs. if you have St. Augustinegrass, Bentgrass, Buffalograss, or Centipedegrass lawns. Set dial to 1 oz. for Bluegrass, Fescue, Ryegrass, Bermudagrass, Bahiagrass, or Zoysiagrass lawns. Spray evenly, wetting weeds and grasses.

WON'T HARM LAWN GRASSES	KILLS WEEDS
Bahiagrass, Bentgrass, Bermudagrass, Buffalograss, Kentucky Bluegrass, Centipedegrass, Fescues, Perennial Ryegrass, Zoysiagrass and St. Augustinegrass (EXCEPT 'Floritam')	Dandelion, Chickweed, Knotweed, Plantains <b>PLUS OVER 80</b> additional broadleaf weeds. See inside label for complete list of weeds.

**Equal Measures:**

1 Tablespoon (Tbs.) = 3 teaspoons (tsp.)  
 1 Fluid Ounce (fl.oz.) = 2 Tablespoons (Tbs.)  
 1 Cup = 8 Fluid Ounces (fl.oz.)

**[Measuring Container Illustration Optional]****MIXING INSTRUCTIONS:**

**For Sensitive/Southern Lawns** - Bentgrass, Buffalograss, Centipedegrass and St. Augustinegrass (EXCEPT Floratam\*).

**Amount to Use:** 1 fl.oz. (2 Tbs.) per gallon of water for each 400 sq.ft.

**[Hose End Sprayer Illustration Optional]**

- When using **Ortho Dial'N Spray**: Set dial to 1 oz. Add 2.5 fl.oz. (5 Tbs.) to sprayer jar. DO NOT add water. Spray evenly over 1,000 sq.ft. until jar is empty.

\*Floratam is a variety of St. Augustinegrass widely grown in Florida

**MIXING INSTRUCTIONS:**

**For Hardy/Northern Lawns** - Bahiagrass, Bermudagrass, Kentucky Bluegrass, Fescues, Perennial Ryegrass, and Zoysiagrass.

**Amount to Use:** 2 fl.oz. (4 Tbs.) per gallon of water for each 400 sq.ft.

**[Hose End Sprayer Illustration Optional]**

- When using **Ortho Dial'N Spray**: Set dial to 2 oz. Add 5 fl.oz. to sprayer jar. DO NOT add water. Spray evenly over 1,000 sq.ft. until jar is empty.

**FOR BEST RESULTS:**

- Measure lawn. Calculate square feet by multiplying length times width.
- Spray when weeds are small and actively growing.
- To reduce risk of lawn injury, avoid spraying drought stressed lawns or when temperature is above 90°F. If dry, water 1 to 2 days before spraying.
- Reapplication can be made in 30 days if necessary.

**Note:** St. Augustinegrass, Centipedegrass or Bermudagrass lawns may have delayed green-up if this product is used when lawns are emerging from dormancy or when temperature drops below 45°F within 72 hours of application. Do not spray Carpetgrass, Dichondra, or desirable clovers. Do not use on or around fruits, vegetables, or flowers.

**Limitations on broadcast applications to residential turfgrass:**

The maximum application rate is 5.0 fl.oz. of product per 1,000 sq.ft. per application (0.37 lbs 2,4-D ae, 0.63 lb MCP-P ae and 0.15 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.0 fl.oz. of product per 1,000 sq.ft. (0.75 lbs 2,4-D ae, 1.26 lb MCP-P ae and 0.31 lb dicamba ae per acre).

**Other Use Precautions:**

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

**Alternate Instructions For Use With The Ortho Dial'N Spray and Measuring Cap Containers.**

**HOW TO APPLY: [MEASURE - Alternative heading]**  
**[Sprayer Illustration Optional] &/or [Walking Man Illustration Optional]**

- To treat entire lawn (broadcast application); Use a tank or hose end sprayer. Measure lawn. Calculate square feet by multiplying length times width. Spray evenly over measured area.
- To spot treat individual weeds in lawns: Use a tank sprayer.
- Rain or watering 6 hours after application will not wash away effectiveness.

**WHEN TO MOW AND WATER: [MOW AND WATER - Alternative heading]**  
**[Lawn Mower Illustration Optional] &/or [Watering Hose Illustration Optional]**

- Do not mow for 1 to 2 days before or after application. [OR - Do not mow for 1 to 2 days after application.]
- If soil is dry, water 1 to 2 days before application.

**WHEN TO APPLY: [Weather Illustration Optional]**

- Spray when weeds are small and actively growing.
- Spray when temperature is below 90°F.
- Reapplication can be made in 30 days if necessary.
- Newly seeded lawn can be sprayed after first mowing.
- Bare spots can be planted with grass seed 1 week after application.

**[Check Mark Illustration Optional] Note: [IMPORTANT - Alternative heading]** St. Augustinegrass, Centipedegrass or Bermudagrass lawns may have delayed green-up if this product is used when lawns are emerging from dormancy or when temperature drops below 45°F within 72 hours of application. Do not spray Carpetgrass, Dichondra, or desirable clovers.

**Other Use Precautions:**

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

**HOW IT WORKS**  
 Because broadleaf weeds and grasses grow differently, (Brand Name) kills broadleaf weeds without harming lawns when used as directed. (Brand Name) singles out weeds, enters through leaves and moves inside the weed to the roots.

**Table 3. MEASURING CAP CONTAINER**

MEASURING CAP: To open: squeeze cap side while unscrewing.



**Chart 1: DILUTION TABLE**

Gallons of Spray	Area to Treat (sq.ft.)	Hardy Lawns (bluegrass, zoysiagrass, fescues, ryegrass, bermudagrass, bahiagrass)	Sensitive Lawns** (St. Augustinegrass, bentgrass, buffalograss, centipedegrass)
1	400	Fill cap to level 4 (2 fl.oz.)	Fill cap to level 3 (1 fl.oz.)
5	2,000	Fill cap to level 4 five times (10 fl.oz.)	Fill cap to level 3 five times (5 fl.oz.)

The measuring cap has 4 levels indicated inside the cap. Fill cap to appropriate level for fast, easy accurate measuring. Close cap tightly after use — no need to rinse.

- Level 1 = 1 teaspoon (tsp.)
- Level 2 = 1/2 fl.oz. = 1 Tablespoon (Tbs.) = 3 tsp.
- Level 3 = 1 fl.oz. (2 Tbs.)
- Level 4 = 2 fl.oz. ( 4 Tbs.)

\*\*See specific directions for St. Augustinegrass and other sensitive lawns.

**Limitations on broadcast applications to residential turfgrass:**

The maximum application rate is 5.0 fl.oz. of product per 1,000 sq.ft. per application (0.37 lbs 2,4-D ae, 0.63 lb MCP-P ae and 0.15 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.0 fl.oz. of product per 1,000 sq.ft. (0.75 lbs 2,4-D ae, 1.26 lb MCP-P ae and 0.31 lb dicamba ae per acre).

**Limitations on spot treatments to residential turfgrass:**

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 5.0 fl.oz. per 1,000 sq.ft. per application [0.63 lbs MCP-P acid equivalent per acre]. The maximum number of spot treatment applications is 2 per year with a minimum of 30 days between applications.

**INSTRUCTIONS FOR USE WITH THE Lock 'N Spray(TM) SYSTEM.**

**HOW TO USE THE Lock 'N Spray(TM) SYSTEM [Walking Man Illustration Optional]**


1. Calculate square feet of lawn by multiplying length times width.
2. Connect sprayer to garden hose by turning the sprayer hose connector.
3. Peel off sticker from the top of cartridge.
4. Point sprayer away from you. Hold cartridge so label faces you. Attach cartridge to sprayer.
5. Turn cartridge in either direction until it clicks into place.
6. Make sure sprayer nozzle tip is down.
7. Turn on water.
8. Extend garden hose to farthest section of lawn to be treated. Squeeze (color) trigger and begin spraying.
9. Spray evenly over entire lawn following the listed coverage rates.
10. When finished, release (color) trigger then turn cartridge to unlock and remove from sprayer. (No need to turn off water.)
11. Flush sprayer with water, squeezing and releasing the (color) trigger in short bursts 2 or more times.
11. Turn off water and squeeze the (color) trigger to relieve water pressure.
12. Disconnect sprayer, shake off excess water, and store in a safe place.
13. When finished spraying, the cartridge reseals itself automatically. No spills. No leaks.




[Container to be marked with 5 Lines for 16 oz. size (or 1/2 liter) or 5 Lines for 32 oz. size (or 1 liter)]

**FOR USE WITH THE Lock 'N Spray(TM) SYSTEM**


Lock 'N Spray(TM) makes lawn and garden care easy. Simply attach this cartridge to the Lock 'N Spray(TM) sprayer (sold separately). Eliminates mixing, measuring, and messy clean-up. Prevents spills and leaks. Automatically mixes the right amount of product and water.



Connect sprayer to hose.



Attach and lock cartridge to sprayer. Make sure sprayer nozzle tip is down.



Turn on water and spray. When finished spraying the cartridge reveals itself automatically. No spills. No leaks.

For complete Lock'N Spray(TM) System directions see inside this label.

**WHERE TO USE AND COVERAGE RATES**

**IMPORTANT:** Use (Brand Name) to kill broadleaf weeds in the hardy and sensitive lawns listed below without harming when used as directed. Do not spray Carpetgrass, Dichondra, or lawns containing desirable clovers.

**HARDY LAWNS**

**Bahiagrass, Bermudagrass, (Kentucky) Bluegrass, Fescues, and Perennial Ryegrass and Zoysiagrass:**

- (16 oz. size) Each line on cartridge represents the amount of product needed to cover approximately 500 square feet of lawn; or
- (32 oz. size) Each line on cartridge represents the amount of product needed to cover approximately 1,000 square feet of lawn.

**SENSITIVE LAWNS**

**Bentgrass, Buffalograss, Centipedegrass, and St. Augustinegrass (Do not use on 'Floritam' St. Augustinegrass that is widely grown in Florida):**

- (16 oz. size) Each line on cartridge represents the amount of product needed to cover approximately 1,000 square feet of lawn; or
- (32 oz size) Each line on cartridge represents the amount of product needed to cover approximately 2,000 square feet of lawn.

Note: If applied to St. Augustinegrass, Centipedegrass or Bermudagrass lawns when they are emerging from winter dormancy, or when temperatures drop below 45°F within 72 hours of application, green-up may be delayed.

**FOR BEST RESULTS**

1. Water 1 to 2 days before application if soil is dry.
2. Do not mow 1 to 2 days before and after application. [OR - Do not mow for 1 to 2 days after application.]
3. Spray when weeds are small and actively growing.
4. Spray only when daytime temperature is below 90°F.
5. Hard-to-control weeds may require a repeat application in 30 days.
6. Treated lawn can be planted with grass seed or sod after 1 week.
7. Newly planted lawn can be treated after first mowing.

**Other Use Precautions:**

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.


**HOW IT WORKS**

Because broadleaf weeds and grasses grow differently, (Brand Name) kills broadleaf weeds without harming lawn grasses when used as directed. (Brand Name) singles out broadleaf weeds, enters through the leaves and moves inside the weed to the roots. Leaves begin to curl within a week with complete kill in 2 to 4 weeks.


**Limitations on broadcast applications to residential turfgrass:**

The maximum application rate is 4.0 fl.oz. of product per 1,000 sq.ft. per application (0.30 lbs 2,4-D ae, 0.50 lb MCP-P ae and 0.12 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl.oz. of product per 1,000 sq.ft. (0.60 lbs 2,4-D ae, 1.0 lb MCP-P ae and 0.25 lb dicamba ae per acre).

**For Lock’N Spray(TM) Pint Size:**

<b>PRODUCT FACTS:</b>	
• Treats up to 6,400 square feet (approximate size of 3 tennis courts).	
<b>KILLS WEEDS:</b>	Dandelion, Clover, Chickweed <b>plus over</b> [Optional “No” Weed Icon] 80 additional broadleaf weeds
<b>WHERE TO USE:</b>	<b>ON LAWNS [Optional Lawn Illustration]</b> without harming lawn grasses when used as directed.
<b>AMOUNT TO USE[SPRAY]:</b>	Bent, Buffalo, Centipede, and St. Augustine (Except ‘Floratom’): Each line on cartridge covers approximately 1,000 square feet.
<b>AMOUNT TO USE:</b>	Bahia, Bermuda, Fescues, Kentucky Blue, Perennial Rye and Zoysia: Each line on cartridge covers approximately 500 square feet.
	Questions, Comments or Medical Information? Call: 1-800-xxx-xxxx • <a href="http://www.xxxx.com">http://www.xxxx.com</a>

**For Lock’N Spray(TM) Quart Size:**

<b>PRODUCT FACTS:</b>	
• Treats up to 12,800 square feet (approximate size of 6 tennis courts).	
<b>KILLS WEEDS:</b>	Dandelion, Clover, Chickweed <b>plus over</b> [Optional “No” Weed Icon] 80 additional broadleaf weeds
<b>WHERE TO USE:</b>	<b>ON LAWNS [Optional Lawn Illustration]</b> without harming lawn grasses when used as directed.
<b>AMOUNT TO USE[SPRAY]:</b>	Bent, Buffalo, Centipede, and St. Augustine (Except ‘Floratom’): Each line on cartridge covers approximately 2,000 square feet.
<b>AMOUNT TO USE:</b>	Bahia, Bermuda, Fescues, Kentucky Blue, Perennial Rye and Zoysia: Each line on cartridge covers approximately 1,000 square feet.
	Questions, Comments or Medical Information? Call: 1-800-xxx-xxxx - <a href="http://www.xxxx.com">http://www.xxxx.com</a>

**DIRECTIONS FOR USE WITH THE Lock 'N Spray(TM) SYSTEM.**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**GENERAL PRECAUTIONS AND RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.

**AMOUNT TO SPRAY**

**[Walking Man Illustration Optional]**

- Measure lawn. Calculate square feet by multiplying length times width.
- Use the amount indicated for your lawn variety:  
**Bentgrass, Buffalograss, Centipedegrass and St. Augustinegrass (EXCEPT Floratam, a variety of St. Augustinegrass widely grown in Florida.):**
  - (Quart size) Each line on cartridge covers approximately 2,000 square feet.
  - (Pint size) Each line on cartridge covers approximately 1,000 square feet.**Bahiagrass, Bermudagrass, Fescues, Kentucky Bluegrass, Perennial Ryegrass and Zoysiagrass:**
  - (Quart size) Each line on cartridge covers approximately 1,000 square feet.
  - (Pint size) Each line on cartridge covers approximately 500 square feet.

<b>CONNECT</b>	
(Illustration #1)	Connect sprayer to hose.
(Illustration #2)	Attach cartridge to sprayer. Turn cartridge until it clicks into place.
<b>SPRAY</b>	
(Illustration #3)	To begin spraying, make sure sprayer nozzle tip is down, turn on water, and squeeze trigger. Spray evenly over measured area.
<b>FINISH</b>	
(Reverse Illustration #2)	<ul style="list-style-type: none"> <li>• To stop spraying, release trigger, unlock cartridge and remove from sprayer. (No need to turn off water.)</li> <li>• To clean sprayer, flush with water, squeezing and releasing trigger in 2 short bursts. Turn off water.</li> <li>• To disconnect sprayer, relieve water pressure by squeezing trigger.</li> </ul>
<b>Cartridge reseals itself automatically - No spills, no leaks!</b>	

**Limitations on broadcast applications to residential turfgrass:**

The maximum application rate is 4.0 fl.oz. of product per 1,000 sq.ft. per application (0.30 lbs 2,4-D ae, 0.50 lb MCP-P ae and 0.12 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl.oz. of product per 1,000 sq.ft. (0.60 lbs 2,4-D ae, 1.0 lb MCP-P ae and 0.25 lb dicamba ae per acre).

**WHEN TO MOW AND WATER**

**[Lawn Mower Illustration Optional] &/or [Watering Hose Illustration Optional]**

- Do not mow for 1 to 2 days before or after application. [OR - Do not mow for 1 to 2 days after application.]
- If soil is dry, water 1 to 2 days before application.
- Rain or watering 6 hours after application will not wash away effectiveness.

**WHEN TO APPLY: [Weather Illustration Optional]**

- Spray when:
  - weeds are small and actively growing.
  - temperature is below 90°F.
- Reapplication can be made in 30 days if necessary.
- Newly seeded lawn can be sprayed after first mowing.

- Bare spots can be planted with grass seed 1 week after application.

Other Use Precautions:

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

**IMPORTANT [Check Mark Illustration Optional]**

If an ornamental shrub or tree is in the lawn, keep spray off foliage and exposed roots.

St. Augustinegrass, Centipedegrass or Bermudagrass lawns may have delayed green-up if this product is used when lawns are emerging from dormancy or when temperature drops below 45°F within 72 hours of application. Do not spray Carpetgrass, Dichondra, or desirable clovers.

**HOW IT WORKS**

Because broadleaf weeds and grasses grow differently, (Brand Name) kills broadleaf weeds without harming lawns when used as directed. (Brand Name) singles out weeds, enters through leaves and moves inside the weed to the roots.

**LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

**APPENDIX**


**1. Statements which may appear on different label components depending on packaging configuration.**

- See next panel for additional Precautionary Statements and First Aid
- Net Contents: \_\_\_\_\_
- EPA Est. No. \_\_\_\_\_

**2. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:**

- Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds.
- Kills weeds - not lawn grasses.
- Kills weeds - not lawns.

- For lawns containing Kentucky bluegrass, fescues, bentgrass, perennial ryegrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass, and buffalograss.
- Can be applied with hose-end or pump-up (tank) sprayers.
- Kills (broadleaf) weeds - roots and all.
- For northern and southern grasses.
- Won't harm lawn grasses shown on container label.
- Won't harm lawn grasses.
- Kills weeds in your lawn.
- Formulated for use on northern and sensitive southern lawns.
- Kills major broadleaf lawn weeds: List of the weed species.
- Helps provide a beautiful lawn.
- For cool and warm season lawn grasses
- (Brand Name) kills major broadleaf lawn weeds including dandelion, chickweed, knotweed, in lawns without harming lawn grasses when used as directed.
- For use on grass lawns especially Kentucky bluegrass, fescues, perennial ryegrass, bentgrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass, and buffalograss.
- Can be used on St. Augustinegrass, bentgrass, and buffalograss.
- A Trimec® Turf Herbicide Formulation.
- Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- One quart covers up to 12,800 square feet (of southern turfgrass)" or "One pint covers up to 6,400 square feet (of southern turfgrass)".
- Treats up to (Number) square feet.
- Kills over 80/84 broadleaf lawn weeds including dandelion and others.
- Makes up to (Number) gallons of spray.
- For northern and southern lawns.
- [Brand Name] kills over 100 more weeds.
- Kills over 2 times/twice as many weeds.
- Water based.
- Measuring cap & pour spout.
- [With] measuring cap.
- Easy pour spout.
- Fast & easy measuring cup.
- Low odor formula.
- Low odor.
- Formula for northern and southern grasses/lawns.
- Kills major broadleaf weeds including dandelion, clover, oxalis, chickweed, knotweed plantain, henbit, spurge and others.
- Concentrate
- Fast acting
- Rain-Proof in hours
- Rain-Fast in hours
- For control that doesn't/won't wash away/off when used as directed.
- Effective even with rain in 6 hours.
- Specially formulated for residential use
- [Text for Lock 'N Spray(TM) System.]
  - Lock 'N Spray - Use with the Lock 'N Spray(TM) System/Sprayer.
  - An innovative technical breakthrough in sprayer. For Ortho's Lock 'N Spray(TM) smart versatile sprayer system.
  - Kills weeds - Won't harm lawns - Kills weeds without harming lawn grasses when used as directed.
  - Kills major broadleaf lawn weeds including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge and many others.

- One easy step to a weed free lawn - Helps promote a beautiful lawn.
- No mixing, no measuring, no mess.
- No mix, No Measure, No Messy Clean-Up.
- Easy to use, premeasured, accurate dosage.
- Simple easy to use/one step application
- No spill, clean and convenient to use
- Closed system prevents spills
- Simple and easy.
- Cartridge sprayer refill, self sealing refill.
- Automatic locking closure, automatically mixes the right amount of product.
- Reusable, disposable cartridges.
- Just lock and spray - Snap 'N Spray
- Cartridge for the Lock 'N Spray(TM) System/Sprayer Concentrate.
- Use (Brand Name) to kill broadleaf weeds in lawns including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge, and many others. Won't harm bahiagrass, bentgrass, bermudagrass, bluegrass, buffalograss, centipedegrass, fescues, rye or zoysiagrass. Can also be used on St. Augustinegrass except Floratam.
- 1 pint covers 3,000 to 6,000 sq.ft. (1 quart covers 6,000 to 12,000 sq.ft.) depending on lawn variety. See inside booklet for complete instructions. Specially formulated for residential use.
- Lock 'N Spray(TM) Starter Kit, Reusable Sprayer
- No Measuring, No Mixing
- Simply lock in your selected cartridge and you're ready!
- CLEAN - The unique system mixes the precise dilution of product and water. Lock 'N Spray(TM) cartridges automatically seal for easy, leak-free storage.
- CONVENIENT - Fingertip control for comfortable, precise application.
- ECONOMICAL - Use to cover small or large areas. Cartridges available in a wide selection of lawn and garden products.
- Adjustable for 3 spray patterns (up, down or detach for jet spray).
- Swivel hose connector.
- On-Off trigger.
- Interchangeable cartridge (also sold separately).
- **NOTE TO PHYSICIAN:** Emergency information call 1-800-xxx-xxxx or 1-800-xxx-xxxx.
- Medical Information call 1-800-454-2333
- Product Information call 1-800-xxx-xxxx
- Questions, Comments or Medical Information? 
- Call: 1-800-xxx-xxxx • <http://www.xxxxx.com>
- OPEN. Resealable Label for Directions & Precautions. Press to Reseal.
- Notice: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.
- Guarantee: If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.
- TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.
- [Non-notifications per PRN 95-2]
  - 25% More Free
  - 40 fluid ounces at the 32 fluid ounce price
  - Covers up to an extra 3,200 square feet
  - Treats up to 16,000 square feet (approximate size of 7.5 tennis courts)

## **DOCUMENT CONTROL INFORMATION**

1. **Unique Label Identifier:** 002217-00570.20100715.notif-proposed-clean.doc

2. **Reason for Issue:** PRN 2007-4