



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

July 25, 2017

Eric D. Smith  
Federal Registration Manager  
PBI/Gordon Corporation  
P.O. Box 014090  
Kansas City, MO 64101-0090

Subject: Label Amendment  
Product Name: Gordon's Trimec St. Augustinegrass Broadleaf Herbicide  
EPA Registration Number: 2217-570  
Application Date: March 31, 2017  
Decision Number: 528230

Dear Mr. Smith:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2  
EPA Reg. No. 2217-570  
Decision No. 528230

with FIFRA section 6. If you have any questions, please contact Mindy Ondish by phone at 703-605-0723, or via email at [ondish.mindy@epa.gov](mailto:ondish.mindy@epa.gov).


Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague". The signature is written in a cursive style with a long horizontal flourish extending to the right.

Kathryn Montague, Product Manager 23  
Herbicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure

# GORDON'S TRIMEC® ST. AUGUSTINEGRASS BROADLEAF HERBICIDE



## EPA Reg. No. 2217-570

### ACTIVE INGREDIENTS:

Mecoprop-p, dimethylamine salt .....	5.30%
2,4-D, dimethylamine salt .....	3.05%
Dicamba, dimethylamine salt .....	1.29%

OTHER INGREDIENTS: .....	<u>90.36%</u>
TOTAL .....	100.00%

### THIS PRODUCT CONTAINS:

- 0.37 lb. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 4.38%
- 0.22 lb. 2,4-Dichlorophenoxyacetic acid equivalent per gallon or 2.53%
- 0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.08%

Isomer Specific by AOAC Method 6.DO1-5.

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

See attached booklet for complete [Precautionary Statements and] Directions for Use, including Storage and Disposal, and back panel for [complete Precautionary Statements, including] First Aid.

Net Contents: \_\_\_\_\_

EPA Est. No. \_\_\_\_\_

### Company Name and Address:

PBI/Gordon Corporation  
1217 West 12<sup>th</sup> Street  
Kansas City, Missouri 64101



## **PRECAUTIONARY STATEMENTS**

### **Hazards to Humans and Domestic Animals**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and rubber gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

<b>First Aid</b>	
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	

### **Environmental Hazards**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

---

## SUBLABEL 1

### Standard Directions for Use

---

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

##### **USE RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply this product through any type of irrigation system.
- Do not spray on carpetgrass, dichondra, nor on lawns or turf where desirable clovers are present. Do not exceed specified dosages for any area.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

[Note to Reviewer: The highlighted bullet points above were moved from Use Precautions below]

##### **USE PRECAUTIONS:**

- Avoid spraying St. Augustinegrass when temperature is above 90°F and under conditions of high humidity. Under these conditions, some slight temporary discoloration of the St. Augustinegrass foliage may occur. This product may be injurious to St. Augustinegrass, bentgrass and centipedegrass in cases of accidental overdose.

##### **WHAT IT IS, WHERE & HOW TO USE IT:**

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide is a Trimec® turf herbicide formulation, a combination of three proven weed killers: 2,4-D, mecoprop-p and dicamba. Together, they control a wide range of broadleaf lawn weeds as listed. This particular formulation is low in 2,4-D for sensitive grasses such as St. Augustinegrass, bentgrass and buffalograss. Spraying should be done when broadleaf weeds are small and growing actively for best results.

1. Irrigate the lawn 2 or 3 days prior to spraying. Do not spray lawns under drought conditions.
2. Do not water for 24 hours after treatment.
3. To avoid damage do not apply to St. Augustinegrass or centipedegrass while emerging from winter dormancy. Also, damage may occur to these grasses if temperatures drop below 45°F within 72 hours of application.

##### **HAND-OPERATED SPRAYER INSTRUCTIONS**

**St. Augustinegrass, Bentgrass and Buffalograss:** Mix 1 fl.oz. in 1 gallon water and uniformly spray 400 sq.ft. If necessary, repeat treatment in 30 days if hard-to-kill weeds persist. Do not use on 'Floritam' St. Augustinegrass in Florida.

**Bermudagrass, Bahiagrass, Kentucky Bluegrass, Centipedegrass, Fescues, Perennial Ryegrass, and Zoysiagrass:** Mix 1.5 fl.oz. in 1 gallon water and uniformly spray 400 sq.ft. On hard-to-kill species, or on older weeds, mix 2 fl.oz. in 1 gallon water and spray on 400 sq.ft.

**Limitations on spot treatments to residential turfgrass:**

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 4.0 fl.oz. per 1,000 sq.ft. per application [0.50 lbs MCP-P acid equivalent, 0.30 lbs 2,4-D acid equivalent, and 0.12 lbs dicamba acid equivalent per acre]. The maximum number of spot treatment applications is 2 per year with a minimum of 30 days between applications.

Broadleaf Weeds Controlled [82]			
Aster, annual	Clover	Lespedeza	Ragweed
Bedstraw	Clover, red	Mallow	Scarlet pimpernel
Beggarweed	Clover, white	Matchweed	Shepherd's purse
Bindweed	Cocklebur	Morningglory	Smartweed,
Bittercress	Cudweed	Mustard	Pennsylvania
Black medic	Daisy, English	Mustard, tansy	Sorrel, sheep
Brass buttons	Dandelion	Oxalis	Sowthistle
Buckhorn	Dandelion, false	Pennywort(creeping	Speedwell
Burclover	Dock	Charlie)	Spurge
Burdock	Dock, curly	Peppergrass	Spurge, prostrate
Burweed	Fleabane, hairy	Pigweed	Spurge, spotted
Buttercup	Galinsoga	Piney woods bedstraw	Thistle
Buttonweed	Geranium	Plantain	Thistle, bull
Carolina geranium	Ground ivy	Plantain, broadleaf	Thistle, musk
Carpetweed	Hawkweed	Plantain, narrowleaf	Velvetleaf
Catsear	Healall	Poison ivy	Wild carrot
Chickweed	Henbit	Poison oak	Wild garlic
Chickweed, common	Henbit, red	Prickly lettuce	Wild lettuce
Chickweed, mouseear	Knawel	Puncturevine	Wild onion
Chicory	Knotweed	Purslane	Wild strawberry
Cinquefoil	Lambsquarters	Pusley, Florida	Yarrow

**For Different Brands, Types, or Sizes of Hose End Sprayers**

**A.**

**(INSERT BRAND NAME) HOSE END SPRAYER INSTRUCTIONS:** Use a hose end sprayer designed for lawns. Pour recommended amount of product (Column 1) in bottle and add water to indicated level (Column 2) and apply spray uniformly to area indicated (Column 3).

For Kentucky Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, Centipedegrass, and Zoysiagrass:			
Product, Amount to Use:		Water, Fill Bottle to: (Gallons)	Area to Treat: (Square Feet)
(Tablespoons)	(Fluid Ounces)		
8	4	8	1,000
16	8	16	2,000
24	12	24	3,000
32	16	32	4,000

For St. Augustinegrass, Bentgrass and Buffalograss:			
Product, Amount to Use:		Water, Fill Bottle to: (Gallons)	Area to Treat: (Square Feet)
(Tablespoons)	(Fluid Ounces)		
5	2.5	5	1,000
10	5.	10	2,000
15	7.5	15	3,000
20	10.	20	4,000
25	12.5	25	5,000

**B.**

**(INSERT BRAND NAME) HOSE END SPRAYER INSTRUCTIONS:** Use a hose end sprayer designed for lawns. Pour recommended amount of product (Column 1) in bottle and add water to indicated level (Column 2) and apply spray uniformly to area indicated (Column 3).

<b>For Kentucky Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, Centipedegrass, and Zoysiagrass:</b>		
<b>Product, Amount to Use: (Fluid Ounces)</b>	<b>Water, Fill Bottle to: (Gallons)</b>	<b>Area to Treat: (Square Feet)</b>
1.5	1	400
3	2	800
6	4	1,600
9	6	2,400

<b>For St. Augustinegrass, Bentgrass and Buffalograss:</b>		
<b>Product, Amount to Use: (Fluid Ounces)</b>	<b>Water, Fill Bottle to: (Gallons)</b>	<b>Area to Treat: (Square Feet)</b>
1	1	400
2	2	800
4	4	1,600
6	6	2,400

<b>CONVERSION TABLE</b>
1 Tablespoon (Tbs.) = 3 teaspoons (tsp.)
1 Fluid Ounce (fl.oz.) = 2 Tablespoons (Tbs.)
1 Cup = 8 Fluid Ounces (fl.oz.)

**Limitations on broadcast applications to residential turfgrass:**

The maximum application rate is 4.0 fl.oz. of product per 1,000 sq.ft. per application (0.30 lbs 2,4-D ae, 0.50 lb MCP-p ae and 0.12 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 fl.oz. of product per 1,000 sq.ft. (0.60 lbs 2,4-D ae, 1.0 lb MCP-p ae and 0.25 lb dicamba ae per acre).

**STORAGE & DISPOSAL [Trashcan Illustration Optional]**

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or [www.website.com](http://www.website.com) for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL

DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.



---

## SUBLABEL 2

### Extended Sprayer Options

---

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### **USE RESTRICTIONS:**

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

[Note to Reviewer: The above bullet points were moved from page 8: Other Use Precautions to create one cohesive list of restrictions]

#### **WHAT IT IS:**

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide is a Trimec® turf herbicide formulation, a combination of three proven weed killers 2,4-D, mecoprop-p and dicamba. Together, they control a wide range of listed broadleaf lawn weeds in residential lawns.

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide kills major northern and southern broadleaf lawn weeds as listed, such as dandelion, chickweed, knotweed, plantain in lawns without harming lawn grasses when used as directed. :

#### **[Weed Illustration Optional]**

Broadleaf Weeds Controlled [82]			
Aster, annual	Clover	Lespedeza	Ragweed
Bedstraw	Clover, red	Mallow	Scarlet pimpernel
Beggarweed	Clover, white	Matchweed	Shepherd's purse
Bindweed	Cocklebur	Morningglory	Smartweed,
Bittercress	Cudweed	Mustard	Pennsylvania
Black medic	Daisy, English	Mustard, tansy	Sorrel, sheep
Brass buttons	Dandelion	Oxalis	Sowthistle
Buckhorn	Dandelion, false	Pennywort(creeping	Speedwell
Burclover	Dock	Charlie)	Spurge
Burdock	Dock, curly	Peppergrass	Spurge, prostrate
Burweed	Fleabane, hairy	Pigweed	Spurge, spotted
Buttercup	Galinsoga	Piney woods bedstraw	Thistle
Buttonweed	Geranium	Plantain	Thistle, bull
Carolina geranium	Ground ivy	Plantain, broadleaf	Thistle, musk
Carpetweed	Hawkweed	Plantain, narrowleaf	Velvetleaf
Catsear	Healall	Poison ivy	Wild carrot
Chickweed	Henbit	Poison oak	Wild garlic
Chickweed, common	Henbit, red	Prickly lettuce	Wild lettuce

Chickweed, mouseear Chicory Cinquefoil	Knawel Knotweed Lambsquarters	Puncturevine Purslane Pusley, Florida	Wild onion Wild strawberry Yarrow
--	-------------------------------------	---	---

**WHERE TO USE:**

Gordon's Trimec® St. Augustinegrass Broadleaf Herbicide is intended for use on most ornamental lawns including Kentucky bluegrass, perennial ryegrass, fescues, bermudagrass, bahiagrass, bentgrass, buffalograss, centipedegrass, zoysiagrass, and St. Augustinegrass.

Do not use this product on 'Floritam' St. Augustinegrass, carpetgrass, dichondra, or lawns containing desirable clovers. Do not spray roots of trees and ornamentals.

**HOW TO TIPS:**

- Spray when broadleaf weeds are small and actively growing for best results.
- Be sure lawn is well watered. If dry, water 1 to 2 days before spraying. To reduce risk of lawn injury, avoid spraying drought stressed lawns or when temperatures are above 90°F.
- Rain or watering within 24 hours after treatment may reduce effectiveness.
- For best results, use hose end sprayers or tank type (pump-up pressure) sprayers. Spray evenly over area at specified rates.

**HOW MUCH TO USE:**

Spot Treatments:

**For St. Augustinegrass, Bentgrass, Buffalograss, and Centipedegrass:**

For Tank Sprayers - Use 1/2 fluid ounce (1 Tablespoon) per gallon of water to cover 200 square feet of lawn. Do not use on Floritam or other improved varieties of St. Augustinegrass.

Do not apply to St. Augustinegrass or centipedegrass emerging from winter dormancy. Temperatures below 45°F within 72 hours of application may cause damage.

**For Bluegrass, Fescues, Perennial Ryegrass, Bahiagrass, Bermudagrass, and Zoysiagrass:** Use 1 fluid ounce (2 Tablespoons) per gallon of water to cover 200 square feet.

Repeat treatment for hard-to-control weeds in 30 days if they regrow.

Do not use a hose end sprayer to spray small areas or spot treat weeds.

**Limitations on spot treatments to residential turfgrass:**

Spot treatment is defined as a treatment area no greater than 1,000 sq.ft. per acre. The maximum application rate is 5.0 fl.oz. per 1,000 sq.ft. per application [0.63 lbs MCP-P acid equivalent, 0.37 lbs 2,4-D acid equivalent, and 0.15 lbs dicamba acid equivalent per acre]. The maximum number of spot treatment applications is 2 per year with a minimum of 30 days between applications.

Broadcast Treatments

**FOR ORTHO HOSE-END SPRAYERS.**

**For St. Augustinegrass, Bentgrass, Buffalograss, and Centipedegrass:**

- **For Tank Sprayers:** Use 1/2 fl.oz. (1 Tbs.) per gallon of water to cover 200 sq.ft. of lawn.
- **For the Dial 'N Spray Hose-End Sprayer:** Measure 2.5 fl.oz. (5 Tbs.) into sprayer jar for every 1,000 sq.ft. of lawn. Do not add water to jar. Set dial to 1 Tbs. and spray.
- **For the Ortho Plant & Weed Sprayette® Hose-End Sprayer:** Put 2.5 fl.oz. (5 Tbs.) into sprayer jar. Fill sprayer jar to the 5 gallon mark with water and spray to cover 1,000 sq.ft.

**For Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Bahiagrass, and Zoysiagrass:**

- **For Tank Sprayers:** Use 1 fl.oz. (2 Tbs.) per gallon of water to cover 200 sq.ft. of lawn.
- **For Dial 'N Spray:** Measure 5 fl.oz. (10 Tbs.) into sprayer jar for every 1,000 sq.ft. of lawn. Do not add water to the container jar. Set dial on 1 oz. and spray.

- **For Ortho Plant & Weed Sprayette® Hose-End Sprayer:** Measure 5 fl.oz. (10 Tbs.) into sprayer jar. Fill with water to the 5 gallon mark and spray to cover 1,000 sq.ft.

**Limitations on broadcast applications to residential turfgrass:**

The maximum application rate is 5.0 fl.oz. of product per 1,000 sq.ft. per application (0.37 lbs 2,4-D ae, 0.63 lb MCP-P ae and 0.15 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.0 fl.oz. of product per 1,000 sq.ft. (0.75 lbs 2,4-D ae, 1.26 lb MCP-P ae and 0.31 lb dicamba ae per acre).

**STORAGE & DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

**LIMITED WARRANTY AND DISCLAIMER**

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

## **APPENDIX**

### **1. Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:**

- Kills dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds.
- Kills weeds - not lawn grasses.
- Kills weeds - not lawns.
- For lawns containing Kentucky bluegrass, fescues, bentgrass, perennial ryegrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass, and buffalograss.
- Can be applied with hose-end or pump-up (tank) sprayers.
- Kills (broadleaf) weeds - roots and all.
- For northern and southern grasses.
- Won't harm lawn grasses shown on container label.
- Won't harm lawn grasses.
- Kills weeds in your lawn.
- Formulated for use on northern and sensitive southern lawns.
- Kills major broadleaf lawn weeds: List of the weed species.
- Helps provide a beautiful lawn.
- For cool and warm season lawn grasses
- (Brand Name) kills major broadleaf lawn weeds including dandelion, chickweed, knotweed, in lawns without harming lawn grasses when used as directed.
- For use on grass lawns especially Kentucky bluegrass, fescues, perennial ryegrass, bentgrass, bermudagrass, bahiagrass, centipedegrass, zoysiagrass, St. Augustinegrass, and buffalograss.
- Can be used on St. Augustinegrass, bentgrass, and buffalograss.
- A Trimec® Turf Herbicide Formulation.
- Controls dandelion, chickweed, knotweed, plantain, henbit, spurge and many other broadleaf lawn weeds including those shown on the side panel.
- One quart covers up to 12,800 square feet (of southern turfgrass)" or "One pint covers up to 6,400 square feet (of southern turfgrass)".
- Treats up to (Number) square feet.
- Kills over 80/84 broadleaf lawn weeds including dandelion and others.
- Makes up to (Number) gallons of spray.
- For northern and southern lawns.
- [Brand Name] kills over 100 more weeds.
- Kills over 2 times/twice as many weeds.
- Water based.
- Measuring cap & pour spout.
- [With] measuring cap.
- Easy pour spout.
- Fast & easy measuring cup.
- Low odor formula.
- Low odor.
- Formula for northern and southern grasses/lawns.
- Kills major broadleaf weeds including dandelion, clover, oxalis, chickweed, knotweed plantain, henbit, spurge and others.
- Concentrate
- Fast acting
- Rain-Proof in hours
- Rain-Fast in hours
- For control that doesn't/won't wash away/off when used as directed.
- Effective even with rain in 6 hours.
- Formulated for residential use
- [Text for Lock 'N Spray(TM) System.]
  - Lock 'N Spray - Use with the Lock 'N Spray(TM) System/Sprayer.

- An innovative technical breakthrough in sprayer. For Ortho's Lock 'N Spray(TM) smart versatile sprayer system.
- Kills weeds - Won't harm lawns - Kills weeds without harming lawn grasses when used as directed.
- Kills major broadleaf lawn weeds including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge and many others.
- One easy step to a weed free lawn - Helps promote a beautiful lawn.
- No mixing, no measuring, no mess.
- No mix, No Measure, No Messy Clean-Up.
- Easy to use, premeasured, accurate dosage.
- Simple easy to use/one step application
- No spill, clean and convenient to use
- Closed system prevents spills
- Simple and easy.
- Cartridge sprayer refill, self sealing refill.
- Automatic locking closure, automatically mixes the right amount of product.
- Reusable, disposable cartridges.
- Just lock and spray - Snap 'N Spray
- Cartridge for the Lock 'N Spray(TM) System/Sprayer Concentrate.
- Use (Brand Name) to kill broadleaf weeds in lawns including dandelion, clover, oxalis, chickweed, knotweed, plantain, henbit, spurge, and many others. Won't harm bahiagrass, bentgrass, bermudagrass, bluegrass, buffalograss, centipedegrass, fescues, rye or zoysiagrass. Can also be used on St. Augustinegrass except Floratam.
- 1 pint covers 3,000 to 6,000 sq.ft. (1 quart covers 6,000 to 12,000 sq.ft.) depending on lawn variety. See inside booklet for complete instructions. Formulated for residential use.
- Lock 'N Spray(TM) Starter Kit, Reusable Sprayer
- No Measuring, No Mixing
- Simply lock in your selected cartridge and you're ready!
- CLEAN - The unique system mixes the precise dilution of product and water. Lock 'N Spray(TM) cartridges automatically seal for easy, leak-free storage.
- CONVENIENT - Fingertip control for comfortable, precise application.
- ECONOMICAL - Use to cover small or large areas. Cartridges available in a wide selection of lawn and garden products.
- Adjustable for 3 spray patterns (up, down or detach for jet spray).
- Swivel hose connector.
- On-Off trigger.
- Interchangeable cartridge (also sold separately).
- **NOTE TO PHYSICIAN:** Emergency information call 1-800-xxx-xxxx or 1-800-xxx-xxxx.
- Medical Information call 1-800-454-2333
- Product Information call 1-800-xxx-xxxx
- Questions, Comments or Medical Information?
- Call: 1-800-xxx-xxxx • <http://www.xxxxx.com>
- OPEN. Resealable Label for Directions & Precautions. Press to Reseal.
- Notice: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.
- Guarantee: If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.
- TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.
- [Non-notifications per PRN 95-2]
  - 25% More Free
  - 40 fluid ounces at the 32 fluid ounce price
  - Covers up to an extra 3,200 square feet
  - Treats up to 16,000 square feet (approximate size of 7.5 tennis courts)




- For use on Bahiagrass, Bermudagrass, Bentgrass, Buffalograss, Bluegrass, Centipedegrass, Ryegrass, Fescue, Zoysiagrass and St. Augustinegrass. (Do not use on Floratam St. Augustinegrass in Florida)
- **For information and questions concerning this product, please contact us by email at support@southernag.com**

CONVERSION TABLE
1 Tablespoon (Tbs.) = 3 teaspoons (tsp.)
1 Fluid Ounce (fl.oz.) = 2 Tablespoons (Tbs.)
1 Cup = 8 Fluid Ounces (fl.oz.)

**[Note to Reviewer: This language was previously approved, just moved into the appendix from sublabel 2]**

**[Note to supplemental registrant.** Select the appropriate container size and present only the information provided for that size on your product labeling.]

PRODUCT FACTS:	
* Treats up to 6,400 square feet (approximate size of 3 tennis courts). Pint Size	
* Treats up to 12,800 square feet (approximate size of 6 tennis courts). Quart Size	
* Treats up to 21,200 square feet (approximate size of 10 tennis courts). 42 Gallon Size	
* Treats up to 51,200 square feet (approximate size of 24 tennis courts). Gallon Size	
<b>KILLS WEEDS:</b>	Dandelion, Clover, Chickweed <b>plus over</b> [Optional "No" Weed Icon] 80 additional broadleaf weeds
<b>WHERE TO USE:</b>	<b>ON LAWNS [Optional Lawn Illustration]</b> without harming lawn grasses when used as directed.
<b>AMOUNT TO USE:</b>	Bent, Buffalo, Centipede, and St. Augustine (Except 'Floratam'): 1 fluid ounce (2 Tablespoons) / gallon of water / 400 square feet
<b>AMOUNT TO USE:</b>	Bahia, Bermuda, Fescues, Kentucky Blue, Perennial Rye and Zoysia: 2 fluid ounces (4 Tablespoons) / gallon of water / 400 square feet
	Questions, Comments or Medical Information? Call: 1-800-xxx-xxxx • <a href="http://www.xxxx.com">http://www.xxxx.com</a>

**[Note to Reviewer: This language was previously approved, just moved into the appendix from sublabel 2]**

## **DOCUMENT CONTROL INFORMATION**

**1. Unique Label Identifier:** 002217-00570.20170719.amend-proposed-clean

**2. Reason for Issue:** EPA Comments #1 + combine restrictions, combine weeds lists, ad claims, minor corrections, formatting, delete obsolete sprayer instructions and duplications in sublabel 2

Internal Usage Notes
Sublabel 1. Associated Container Label: Supplemental Distributors
Sublabel 2. Associated Container Label: Supplemental Distributors